Zeitschrift: Helvetica Physica Acta

Band: 65 (1992)

Heft: 2-3

Artikel: Frustrated XY model : continuity of the ground state energy in function

of the frustration

Autor: Valat, A. / Beck, H.

DOI: https://doi.org/10.5169/seals-116492

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 29.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Frustrated XY Model: Continuity of the Ground State Energy in Function of the Frustration

A. Vallat and H. Beck Institut de Physique rue Bréguêt 1, CH 2000 Neuchâtel, Switzerland

Abstract. The frustrated XY model can describe Josephson-junction arrays in transverse magnetic fields. It is shown that the ground state energy is uniformly continuous when the frustration runs through all the real numbers. A consequence of this fact is that the critical current of the Josephson-junction array is non-zero at T=0 for any frustration.

Introduction

This contribution concerns the uniformly frustrated XY model (illustrated by the case of an infinite square lattice). Since the periodicity of the ground state configurations for a rational frustration f = p/q depends essentially on the denominator q, one could have doubts about the continuity (over all the real numbers) of the ground state energy E(f) (energy per bond) in function of the frustration f. Indeed one can read in several papers[1,2] that E(f) is not continuous. This question is relevant to study several quantities, particularly at T = 0, such as the helicity modulus or the critical current of a Josephson-junction array. The behaviour of E(f) is also relevant for the discussion of the stability of commensurate flux phase in the tJ model[3].

The model

The hamiltonian of the system in site variables θ_r is: $H = -\sum_{\langle rr' \rangle} \cos(\theta_r - \theta_{r'} - A_{rr'})$ such that: $\theta_r \in (-\pi, \pi]$ and $\sum_{\square R} A_{rr'} = 2\pi f$ where $\sum_{\square R}$ means that the sum is taken in clockwise direction over the bonds surrounding the plaquette R.

This system is exactly equivalent to the following, written in the bond variables $\phi_{rr'}$: $H = -\sum_{\langle rr' \rangle} \cos(\phi_{rr'})$ with the constraints: $\phi_{rr'} = -\phi_{r'r} \in (-\pi, \pi]$ and $\sum_{\square R} \phi_{rr'} = -2\pi f + 2\pi m_R$ for each plaquette where m_R is any integer corresponding to a topological charge.

Uniform continuity of E(f)

Definition: E(f) is uniformly continuous in the real numbers if $\forall \epsilon > 0, \exists \ \delta(\epsilon)$ such that for all real f, \bar{f} , one has: $|\bar{f} - f| < \delta \Rightarrow |E(\bar{f}) - E(f)| < \epsilon$.

From a ground state $\Phi_f \equiv \{\phi_{rr'}\}$ corresponding to the frustration f, we are going to construct some states $\bar{\Phi}_{fn} \equiv \{\bar{\phi}_{rr'}\}$ for a frustration \bar{f} of energy $E_{fn}(\bar{f})$ close to the energy $E(\bar{f})$:

(a): Choosing a positive integer n, one cuts out the lattice into vertical strips containing horizontal line of 2n + 1 sites (Fig.1).




Figure 1: The infinite lattice is cut out into vertical strips. We call a plaquette (and a bond) "joint plaquette" (and "joint bond") if a frontier crosses it and "internal plaquette (and "internal bond") if it doesn't.

(b): To the internal bonds, the following transformation is applied:

$$\phi_{rr'} \mapsto \bar{\phi}_{rr'} = \begin{cases} \phi_{rr'} & \text{on the horizontal bonds} \\ \phi_{rr'} + 2\pi(f - \bar{f})d_{rr'} & \text{on the vertical bonds} \end{cases}$$
(1)

where the vertical bond rr' is considered such that r is under r' and $d_{rr'}$ is the algebraic distance between the bond and the central vertical line of the strip belonging to it. So all internal plaquette constraints are satisfied for \bar{f} .

(c): The $\phi_{rr'}$ of joint bonds crossing one frontier are determinated by the plaquette constraints after choosing freely one of them. Then this choice is done such that the average of $\cos \bar{\phi}_{rr'}$ crossing each frontier is positive (it is possible by adding π to each $\bar{\phi}_{rr'}$ if the average would be negative). One notes $E_{fn}(\bar{f})$ the energy of the state $\bar{\Phi}_{fn} \equiv \{\bar{\phi}_{rr'}\}$ thus constructed.

Now the mean variation over any set of bonds corresponding to 2n + 1 sites of one strip in a horizontal line (see bold bonds in Fig.1) is:

$$\frac{1}{2(2n+1)} \sum_{\langle rr' \rangle} \left[-\cos(\bar{\phi}_{rr'}) + \cos(\phi_{rr'}) \right] \le \frac{1}{2(2n+1)} \left[\sum_{vert.bds} 2\pi \mid \bar{f} - f \mid\mid d_{rr'} \mid + 2 \right]$$
(2)

The sum on right hand side puts an upper bound on the variation on the vertical bonds and the variation due to the horizontal joint bond is bounded by 2. Moreover the mean of the variation on the joint bonds over a frontier is bounded by 1 (according to (c)). So one can write:

$$E_{fn}(\bar{f}) - E(f) \le \frac{1}{2(2n+1)} \left[\sum_{vert.bds} 2\pi \mid \bar{f} - f \mid \mid d_{rr'} \mid +1 \right]$$

$$= \frac{1}{2(2n+1)} \left[n(n+1)2\pi \mid \bar{f} - f \mid +1 \right]. \tag{3}$$

Let ε such that $0 < \varepsilon < 1/2$. One can choose $n(\varepsilon) \in [\frac{1}{2\varepsilon} - 1, \frac{1}{2\varepsilon}]$ and $\delta(\varepsilon) = 2\varepsilon^2/\pi$, so the right hand side of (Eq.3) will be less than ε (when $|\bar{f} - f| < \delta(\varepsilon)$). Then one has found $\delta(\varepsilon)$ such that, for a certain $n(\varepsilon)$, one has $|\bar{f} - f| < \delta(\varepsilon) \Rightarrow E_{fn}(\bar{f}) - E(f) < \varepsilon$. As, by definition, $E(\bar{f}) \leq E_{fn}(\bar{f})$, one has: $E(\bar{f}) - E(f) < \varepsilon$.

Since this statement depends only on $|f - \bar{f}|$ one has also by exchanging f and \bar{f} : $E(f) - E(\bar{f}) < \varepsilon$. Therefore $\forall \varepsilon > 0, \exists \delta$ such that for all real numbers f, \bar{f} , one has $|\bar{f} - f| < \delta \Rightarrow |E(\bar{f}) - E(f)| < \varepsilon$.

Thus the ground state energy is uniformly continuous when f runs through the real numbers. Moreover by knowing E(0) = -1 and $E(1/2) = -1/\sqrt{2}$, and using the dependence $\delta(\varepsilon) = 2\varepsilon^2/\pi$, it is easy to see that E(f) is always strictly negative.

Critical current of Josephson-junction array

Following Teitel and Jayaprakash[4], the T=0 critical current of the array can be defined by:

$$i_c(f) = \max_{\delta} \{ (i_{c0}/N) \sum_{vert,bds} \sin(\phi_{rr'} + \delta) \}$$
(4)

where i_{c0} is the single junction critical current, N is the number of sites and $\{\phi_{rr'}\}$ is a ground state configuration. Since $\sin x = \cos(x - \pi/2)$, we have:

$$i_c(f) = \max_{\delta} \{ (i_{c0}/N) \sum_{vert,bds} \cos(\phi_{rr'} + \delta - \pi/2) \}$$
 (5)

The sum of cosines reaches its maximum, the absolute value of the ground state energy (-NE(f)), at $\delta = \pi/2$. So $i_c(f) = -i_{c0}E(f)$ and $i_c \neq 0$ even if f is irrational.

References

- [1] T.C.Halsey, Phys.Rev. B 31, 5728 (1985).
- [2] S. Teitel, Physica B 152, 30 (1988).
- [3] For instance: P.Lederer, D.Poilblanc, T.M.Rice, Phys.Rev.Lett. 63, 1519 (1989).
- [4] S. Teitel, C. Jayaprakash, Phys. Rev. Lett. 51, 1999 (1983).