

Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 2-3

Artikel: Crossover from two to three dimensions in the layered xy-model
Autor: Schmidt, A. / Schneider, T.
DOI: <https://doi.org/10.5169/seals-116493>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Crossover from two to three dimensions in the layered *xy*-model

A. Schmidt and T. Schneider

IBM Research Division, Zurich Research Laboratory, Säumerstr. 4
CH-8803 Rüschlikon, Switzerland

Abstract. Using Monte-Carlo simulations we studied the layered *xy*-model. The decrease of the transition temperature T_c with reduced thickness and the behavior of the specific heat are estimated. In an approximate Ginzburg-Landau treatment we find that the drop of T_c and the associated dimensional crossover is a combined effect of fluctuations and boundaries.

Introduction

As a model to study the crossover between two and three dimensions we investigate the layered *xy*-model defined by

$$H = -J_{\parallel} \sum_{\langle ij \rangle_{\parallel}} \vec{S}_i \cdot \vec{S}_j - J_{\perp} \sum_{\langle ij \rangle_{\perp}} \vec{S}_i \cdot \vec{S}_j, \quad (1)$$

where J_{\parallel} (J_{\perp}) is the intra- (inter-) layer coupling strength, $\vec{S}_i = (\cos \varphi_i, \sin \varphi_i)$ are the *xy*-spins on a simple cubic lattice of $N = L^2 M$ spins and $\langle ij \rangle_{\parallel, \perp}$ denote nearest neighbors within and perpendicular to the planes. To mimic a real system of finite thickness we adopt periodic boundary conditions parallel to the layers and free boundaries perpendicular to them.

Numerical method

Using a Metropolis algorithm and the single-cluster algorithm [1] together with the multihistogram technique [2], we estimate the transition temperature $T_c(M)$ by the intersection point of the fourth-order cumulant [3]

$$U_L(T) = 1 - \frac{\langle (\vec{m}^2)^2 \rangle_T}{3 \langle \vec{m}^2 \rangle_T^2} \quad (2)$$

for different sizes L of the layers. Here $\vec{m} = \frac{1}{N} \sum_{i=1}^N \vec{S}_i$ denotes the magnetization and $\langle \rangle_T$ the thermal average.

Approximate treatment

By using a Hubbard-Stratanovich transformation and a mean-field description perpendicular to the layers we can partly take the 2d fluctuations in the layers into account and approximate the free energy of model (1) up to quartic terms in a Ginzburg-Landau form as

$$\begin{aligned} F \approx & \int d^2 r \left\{ T_c^o \left(1 - \frac{T_c^o}{T} \right) |\psi|^2 + J_{\parallel} \left(\frac{T_c^o}{T} - \frac{1}{2} \right) |\nabla_{\parallel} \psi|^2 \right. \\ & \left. + \frac{1}{(M+1)^2} \sum_{n=1}^M \sin^4 \left(\frac{\pi}{M+1} n \right) \frac{(T_c^o)^4}{T^3} |\psi|^4 \right\} \end{aligned} \quad (3)$$

with the mean-field transition temperature $T_c^o(M) = 2J_{\parallel} + J_{\perp} \cos(\frac{\pi}{M+1})$. Using the Kosterlitz-Thouless transition of this effective 2d *xy*-model we obtain an upper bound $T_{KT}^>(M)$ for the transition temperature [4]

$$T_{KT}^>(M) = T_c^o(M) \left(\frac{3}{2} - \frac{1}{2} \sqrt{1 + 16 \frac{M}{(M+1)^2} \frac{\langle S^4 \rangle}{\pi J_{\parallel}} T_c^o(M)} \right), \quad (4)$$

where $\langle S^4 \rangle \equiv \frac{1}{M} \sum_{n=1}^M \sin^4 \left(\frac{\pi}{M+1} n \right)$.

Results and Conclusions

Figure 1 depicts the numerical estimates $T_c(M)$. The limits for $M = 1$ and $M \rightarrow \infty$ agree with previous estimates [5], but we notice a steep increase of $T_c(M)$ for small M values. This can be understood by invoking finite-size scaling to our system: the correlation length $\xi = b \left(\frac{T_c^{3d} - T}{T_c^{3d}} \right)^{-\nu_{3d}}$ of the 3d system cannot exceed the finite thickness M , so that

$$T_c(M) = T_c^{3d} \left(1 - \left(\frac{b}{M} \right)^{\frac{1}{\nu_{3d}}} \right), \quad (5)$$

where b is a measure of the interaction range corresponding to the zero temperature correlation length. Eq. (5) fits the data very well for $M \geq 4$ and yields $\nu_{3d} = 0.70 \pm 0.08$ and $b = 1.3 \pm 0.5$, in agreement with previous estimates of ν_{3d} [5,6].

Because the interaction range b is of the order 1, the drop of T_c is appreciable only for small M . For comparison we also included the results of our approximate treatment. Here the decrease of T_c with reduced thickness is traced back to the presence of boundaries incorporated in $T_c^o(M)$ and a further reduction due to the 2d fluctuations, becoming dominant for small M values. A related crossover phenomenon occurs in the specific heat: a broad peak well above T_c for $M = 1$, which sharpens and shifts towards T_c with increasing M to approach a sharp cusp at T_c in the bulk.

To summarize: our Monte-Carlo study of the layered isotropic xy -model revealed the drop of the transition temperature as a function of thickness for rather small M . This can be understood in terms of the short interaction range b of order one. With our approximate treatment we find two important contributions to this behavior: the free boundary conditions and the 2d fluctuations.

References

- [1] U. Wolff, Phys. Rev. Lett. **62**, 361 (1989); Nucl. Phys. B **322**, 759 (1989)
- [2] A. M. Ferrenberg and R. H. Swendsen, Phys. Rev. Lett. **63**, 1195 (1989)
- [3] K. Binder, Phys. Rev. Lett. **47**, 693 (1981); Z. Phys. B **43**, 119 (1981)
- [4] A. Schmidt and T. Schneider: preprint
- [5] R. Gupta et al., Phys. Rev. Lett. **61**, 1996 (1988) W. Janke, Phys. Lett. A **148**, 306 (1990)
- [6] J. C. Le Guillou, J. Zinn-Justin, Phys. Rev. B, **21**, 3976 (1980)




Figure 1. $T_c^o(M)$, $T_{KT}(M)$ and the numerical estimates $T_c(M)$. The short dashed line indicates the transition temperature of the 3d xy -model.