

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 65 (1992)  
**Heft:** 2-3

**Artikel:** Long-time dynamics of two-dimensional fluid binary mixture in cellular automata models  
**Autor:** Perera, A. / Penson, K.A. / Schultze, U.  
**DOI:** <https://doi.org/10.5169/seals-116501>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 22.02.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Long-Time Dynamics of Two-Dimensional Fluid Binary Mixture in Cellular Automata Models

A. Perera, K. A. Penson, U. Schultze and M. Moreau

Université Pierre et Marie Curie

Laboratoire de Physique Théorique des Liquides

Tour 16, 5eme et. Boîte 121

F-75252 Paris Cedex 05, France.

**Abstract.** We compute exponents of long-time tails of the velocity auto-correlation functions in binary mixtures in cellular automata models in 2D using numerical simulations. The exponents are found to vary with densities as opposed to universal values for the one component system.

The idea of applying the model system in which both space and time are discrete, to obtain properties of fluids, had proven highly successful. Following the seminal work of Frisch, Hasslacher and Pomeau (FHP) [1] very large number of problems have been treated with such cellular automata (CA) models. For recent references consult an exhaustive review article [2] and Ref. [3]. The problem of long-time tails of the velocity correlation function  $\phi(t) = \langle \vec{v}(0)\vec{v}(t) \rangle$  (VACF) has received increased attention. Recently, with a newly conceived rapid simulation method by van der Hoef and Frenkel [4], the prediction of mode-coupling theory  $\lim_{t \rightarrow \infty} \phi(t) \sim t^{-D/2}$  has been for the first time confirmed for a single component CA fluid in 2D and 3D [4].

Here we report simulation results applying the methods of ref.[4] on a CA representation of fluid binary mixture of unequal species, called heavy (H) and light (L) on a 2D triangular lattice.

As usual in CA model the interactions are specified by a set of collision rules : a) the state of H particles is not affected by the presence of L particles. H particles interact between themselves according to FHP-III collision rules [1],[2]; b) if a lattice node is occupied only by L particles, they interact also according to FPH-III rules; c) if a node is occupied by at least one H particle, the L particles are scattered off H particles randomly; d) no two particles can propagate on the same link and the same direction (exclusion principle).


This model is a generalization of the Lorentz gas [5] : the scattering centers for L particles are H particles. The latter can move and display their own dynamics governed by collisions

with other H particles. Obviously the dynamics of L particles is our main interest here. When the density of L particles is low and that of H particles quite high the diffusion of the former will be analogous to a random walk. Indeed we find numerically that the VACF decays exponentially.

The most interesting case arises when either the two densities  $\rho_H$  and  $\rho_L$  of respectively heavy and light particles are either comparable or when  $\rho_H \leq \rho_L$ . We find in fact that, in the case of our model, the VACF for L particles has a long-time tail with an exponent which depends on  $\rho_H$  and  $\rho_L$  in a non trivial way.

Fig.1 shows a plot of  $\log \phi(t)$  versus  $\log t$  for the particular case of  $\rho_H = 0.4$  and for  $\rho_L = 0.2$  and  $\rho_L = 0.4$ . We see that in both cases the exponent of the long-time tail is clearly smaller than unity. This is an opposite behaviour of the static Lorentz gas where the exponent was found to be 2 [6]. The full parameter space of  $\rho_H$  and  $\rho_L$  ( $\rho_H + \rho_L \leq 1$ ) is being currently explored.

Fig.1 : Log-log plot of the VACF's for densities specified in the inset.



### References

- 1) U. Frisch, B. Hasslacher and Y. Pomeau, Phys. Rev. Lett. **56**, 1505 (1986).
- 2) M. H. Ernst, in "Liquid, Freezing and the Glass Transition.", Les Houches, Session LI, D. Levesque, J. P. Hansen and J. Zinn-Justin, Eds. (North Holland, 1991).
- 3) "Lattice Gas Methods for Partial differential equations", G. D. Dolan, Ed. (Addison-Wesley, 1990)
- 4) M. A. van der Hoef and D. Frenkel, Phys. Rev. A **41**, 4277 (1990).
- 5) A. Perera, B. Gaveau, M. Moreau and K. A. Penson, Phys. Lett. A, in press.
- 6) P.-M. Binder and D. Frenkel, Phys. Rev. A **42**, 2463 (1990).