Zeitschrift: Helvetica Physica Acta
Band: 65 (1992)
Heft: 8

Artikel: Algebraic treatment of the Morse potential
Autor: Chetouani, L. / Guechi, L. / Hammann, T.F.
DOI: https://doi.org/10.5169/seals-116523

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 16.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Helv Phys Acta
Vol. 65 (1992)

0018-0238/92/081069-08$1.50+0.20/0
(c) 1992 Birkhäuser Verlag, Basel

ALGEBRAIC TREATMENT OF THE MORSE POTENTIAL

L. CHETOUANI *, L. GUECHI * and T.F. HAMMANN † #

* Département de Physique Théorique, Institut de Physique,
Université de Constantine, Constantine, Algeria

† Laboratoire de Mathématiques, Physique Mathématique et Informatique,
Faculté des Sciences et Techniques, Université de Haute Alsace,
4, rue des Frères Lumière, F68093 Mulhouse, France

(19. III. 1992)

The Green's function for the Morse potential is calculated in the so(2,1) algebraic approach, using the Baker-Campbell-Hausdorff formulas.

Ever since the first notable success of the algebraic method in the calculation of wave functions and of hydrogen atom transition amplitudes [1], renewed interest for the algebraic approach has been emerging. Hence a certain number of potentials have been studied in the algebraic approach [2,3] and their Green's functions have been calculated. This algebraic method consists mainly in the transformation of the Schrödinger equation via a change of variables, in order to introduce generators satisfying a Lie algebra. When the evolution operator is expressed in terms of these generators in the configuration space, it is calculable for a certain class of potentials. In this paper, this algebraic approach is used to study the Morse potential defined as :

$$V(x) = A\, e^{-2ax} - B\, e^{-ax}, \tag{1}$$

where A, B and a are positive constants.

To whom requests for reprints should be addressed.

This Morse potential has been very useful in molecular and nuclear physics. In a generalized separable and nonlocal form, it has once been used as a model for Nucleon-Nucleon [4] and P ion-Nucleon [5] interactions. The path integral solution to the problem can be found in the literature [6,7]. Two solutions used to be agreed on via the introduction of an auxiliary time-variable. A disagreement about expressions of the Green's function related to this potential and requiring a mathematical clarification [8] can nevertheless be noticed. This problem has been solved very recently [9].

The energy spectrum, propagator and Green's function have also been obtained in the phase-space approach of Weyl-Wigner-Moyal [10] . In the algebraic so (2,1) approach, the Green's function can be obtained in a direct and nice manner. Indeed, let G(x,x';E) be the Green's function which is solution of the differential equation :

$$(H-E)G(x,x';E) = -\hbar i\delta(x-x'), \tag{2}$$

where $H(x) = -\frac{\hbar^2}{2m}\frac{d^2}{dx^2} + V(x)$, is the hamiltonian of the particle and E its energy.

The transformation $\xi = \exp\left(-\frac{ax}{2}\right)$, then gives

$$\left(\widetilde{H}-\widetilde{E}\right)\widetilde{G}\left(\xi,\xi';\widetilde{E}\right) = -\hbar i\delta\left(\xi-\xi'\right), \tag{3}$$

where

$$\widetilde{G}\left(\xi,\xi';\widetilde{E}\right) = \frac{2}{a}\left[\xi\xi'\right]^{1/2} G\left(x,x';E\right). \tag{4}$$

The dynamics of the physical system is then governed by the new hamiltonian

$$\widetilde{H}(\xi,) = -\frac{\hbar^2}{2M}\left(\frac{d^2}{d\xi^2} - \frac{\mu(\mu-1)}{\xi^2}\right) + \frac{1}{2}M\widetilde{\omega}^2\xi^2, \tag{5}$$

where $M = \frac{4m}{a^2}$, $\tilde{\omega}^2 = \frac{2A}{M}$, $\mu = \frac{1}{2} + \left(-\frac{2ME}{\hbar^2}\right)^{1/2}$, (6)

and $B = \tilde{E}$ is the pseudo-energy.

Expression (5) is the hamiltonian of a harmonic oscillator with a constant frequency, constrained to a centrifugal repulsion.

Besides, it is well known that the radial Coulombian system can be shown equivalent to the radial oscillator [9], with the help of a simple $r = \xi^2$ transformation. Consequently the Morse potential and the radial Coulombian system are equivalent and accept the same group dynamics.

It is easy to see that one can introduce the three following generators

$$T_1(\xi) = -\frac{\hbar^2}{2M}\left[\frac{\partial^2}{\partial\xi^2} - \frac{\mu(\mu-1)}{\xi^2}\right] \quad , \quad T_2(\xi) = -\frac{i}{2}\left(\xi\frac{\partial}{\partial\xi} + \frac{1}{2}\right), \; T_3(\xi) = \frac{M}{4\hbar^2}\xi^2 , \tag{7}$$

satisfying the Lie algebra :

$$[T_1,T_2] = -iT_1 , [T_2,T_3] = -iT_3 , \text{and} [T_1,T_3] = -iT_2 \tag{8}$$

The operator $\tilde{H}(\xi)$ can readily be expressed in terms of these generators :

$$\tilde{H}(\xi) = T_1(\xi) + 2\tilde{\omega}^2\hbar^2 T_3(\xi) . \tag{9}$$

Being a linear combination of the generators T_i, $\tilde{H}(\xi)$ shows, as expected, a dynamical symmetry so (2,1) [11]. Expressed in Schwinger's integral representation [12] the solution of the differential equation (3), can be written as follows

$$G(\xi,\xi';\tilde{E}) = \int_0^\infty ds \exp\left[-\frac{i}{\hbar}(\tilde{H} - \tilde{E} - io)s\right]\delta(\xi-\xi') =$$

$$= \int_0^\infty ds \exp\left[\frac{i}{\hbar}\tilde{E}s\right] \exp\left\{\frac{-is}{\hbar}\left[T_1(\xi) + 2\hbar^2\tilde{\omega}^2 T_3(\xi)\right]\right\} \delta(\xi-\xi') . \tag{10}$$

The derivation of the Green's function in the algebraic approach, has been replaced by the calculation of the Kernel

$$P(\xi,\xi';s) = \exp\left\{\frac{-is}{\hbar}\left[T_1(\xi) + 2\hbar^2\tilde{\omega}^2 T_3(\xi)\right]\right\} \delta(\xi-\xi') . \qquad (11)$$

We can achieve this calculation thanks to the following two Baker-Campbell-Hausdorff (BCH) formulas :

$$\exp\left\{\frac{-is}{\hbar}\left[T_1 + 2\hbar^2\tilde{\omega}^2 T_3\right]\right\} = \exp(-iaT_3)\exp(-ibT_2)\exp(-icT_1) , \qquad (12)$$

where

$$a = 2\hbar\tilde{\omega}\tan(\tilde{\omega}s) , \qquad (13a)$$

$$b = 2\mathrm{Ln}\left(\cos(\tilde{\omega}s)\right) , \qquad (13b)$$

$$c = \frac{1}{\hbar\tilde{\omega}}\tan(\tilde{\omega}s) , \qquad (13c)$$

and

$$\exp(-i\alpha T_3)\exp(-i\beta T_2)\exp(-i\gamma T_1) = \exp(-icT_1)\exp(\lambda T_3) , \qquad (14)$$

with

$$\alpha = \frac{i\lambda}{1-\frac{i\lambda c}{2}} , \quad \beta = 2\mathrm{Ln}\left(1-\frac{i\lambda c}{2}\right) , \quad \gamma = \frac{c}{1-\frac{i\lambda c}{2}} \qquad (15)$$

Both formulas can be easily checked within the frame of another realization of the algebra, which may well be chosen as finite, for instance expressing T_i generators versus the Pauli matrices σ_i :

$$T_1 = \frac{\sigma_1 - i\sigma_2}{2\sqrt{2}} , \quad T_2 = -\frac{i\sigma_3}{2} , \quad T_3 = \frac{\sigma_1 + i\sigma_2}{2\sqrt{2}} . \qquad (16)$$

In order to calculate (11), we set

$$\delta(\xi-\xi') = \frac{M}{2\hbar^2}\frac{\xi^{\mu}\xi'^{1-\mu}}{2i\pi}\int_{-i\infty+\delta}^{+i\infty+\delta} d\lambda \exp\left\{\frac{M\lambda}{4\hbar^2}\left(\xi^2-\xi'^2\right)\right\} , \quad \delta<0 . \qquad (17)$$

By using (12, 14, 17), the kernel (11) now reads

$$P(\xi,\xi';s) = \frac{M}{2\hbar^2}\xi'^{1-\mu}\exp(-iaT_3)\exp(-ibT_2)\frac{1}{2i\pi}\int_{-i\infty+\delta}^{+i\infty+\delta} d\lambda\, \exp\left(-\frac{M}{4\hbar^2}\xi'^2\lambda\right)\exp(-icT_1)\exp\left(\frac{M}{4\hbar^2}\xi^2\lambda\right)\xi^{\mu} =$$

$$= \frac{M}{2\hbar^2}\xi'^{1-\mu}\exp(-iaT_3)\exp(-ibT_2)\frac{1}{2i\pi}\int_{-i\infty+\delta}^{+i\infty+\delta} d\lambda\, \exp\left(-\frac{M}{4\hbar^2}\xi'^2\lambda\right)\exp(-icT_1)\exp(\lambda T_3)\,\xi^{\mu} = \qquad (18a)$$

$$= \frac{M}{2\hbar^2}\xi'^{1-\mu}\exp(-iaT_3)\exp(-ibT_2)\frac{1}{2i\pi}\int_{-i\infty+\delta}^{+i\infty+\delta} d\lambda\, \exp\left(\frac{-M}{4\hbar^2}\xi'^2\lambda\right)\exp(-i\alpha T_3)\exp(-i\beta T_2)\exp(-i\gamma T_1)\,\xi^{\mu} = \qquad (18b)$$

$$= \frac{M}{2\hbar^2}\xi'^{1-\mu}\exp(-iaT_3)\exp(-ibT_2)\xi^{\mu}\frac{1}{2i\pi}\int_{-i\infty+\delta}^{+i\infty+\delta} d\lambda\, \frac{\exp\left\{\frac{M}{2\hbar^2}\left(\frac{-\lambda\xi'^2}{2}+\frac{\lambda\xi^2}{2-i\lambda c}\right)\right\}}{\left(1-\frac{i\lambda c}{2}\right)^{\mu+1/2}} = \qquad (18c)$$

$$= \frac{-iM}{\hbar^2 c}\exp(-iaT_3)\exp(-ibT_2)\left[(\xi'\xi)^{1/2} I_{\mu-1/2}\left(\frac{M\xi'\xi}{i\hbar^2 c}\right)\exp\left\{\frac{iM}{2\hbar^2 c}\left(\xi^2+\xi'^2\right)\right\}\right] = \qquad (18d)$$

$$= \frac{-iM}{\hbar^2 c}\exp\left(-\frac{iaM}{4\hbar^2}\xi^2\right)\exp\left(-\frac{b}{2}\right)(\xi'\xi)^{1/2} I_{\mu-1/2}\left(\frac{M\xi'\xi}{i\hbar^2 c e^{b/2}}\right)\exp\left\{\frac{iM}{2\hbar^2 c}\left(e^{-b}\xi^2+\xi'^2\right)\right\} = \qquad (18e)$$

$$= \frac{-iM\tilde{\omega}}{\hbar\sin(\tilde{\omega}s)}(\xi'\xi)^{1/2} I_{\mu-1/2}\left(\frac{M\tilde{\omega}\xi'\xi}{i\hbar\sin(\tilde{\omega}s)}\right)\exp\left\{\frac{iM\tilde{\omega}}{2\hbar}\left(\xi^2+\xi'^2\right)\cot(\tilde{\omega}s)\right\}. \qquad (18f)$$

The choice of the form of the Dirac delta distribution is dictated by the desire to obtain the simple following result :

$$\exp(-i\gamma T_1)\xi^{\mu} = \left(1 - i\gamma T_1 + \frac{1}{2!}(-i\gamma T_1)^2 + \ldots\right)\xi^{\mu} = \xi^{\mu}\ .$$

Moreover we have used the formula

$$\exp(-i\beta T_2)\,f(\xi) = \exp\left\{-\frac{\beta}{2}\left[\xi\frac{\partial}{\partial\xi}+\frac{1}{2}\right]\right\}f(\xi) = \exp\left(-\frac{\beta}{4}\right)\exp\left[-\frac{\beta}{2}\frac{\partial}{\partial \mathrm{Ln}\,\xi}\right]f(\xi) =$$

$$= \exp\left(-\frac{\beta}{4}\right) \exp\left[-\frac{\beta}{2}\frac{\partial}{\partial u}\right] f(e^u) =$$

$$= \exp\left(-\frac{\beta}{4}\right) f(e^{u-\beta/2}) = \exp\left(-\frac{\beta}{4}\right) f(e^{-\beta/2}\xi) , \qquad (19)$$

when going over from eq. (18b) to (18c) and (18d) to (18e).

We calculate the integral contained within the eq. (18c) thanks to the Residue Theorem, by decomposing the $\frac{\lambda}{2-i\lambda c}$ factor and by expanding in a series the term in ξ^2 from the exponential function.

We used the following definitions for the first and second kind Bessel functions $J_\mu(x)$ and $I_\mu(x)$ [13, 14],

$$J_\mu(x) = \left(\frac{x}{2}\right)^\mu \sum_{n=0}^{\infty} \frac{(-x^2/4)^n}{n!\Gamma(n+\mu+1)} = \exp\left(i\frac{\pi}{2}\mu\right) I_\mu(-ix) . \qquad (20)$$

By inserting (18f) into (10) and then into (4), and by setting $\nu = \mu - 1/2$, the Green's function is then given by

$$G(x,x';E) = \frac{-iM\tilde{\omega}a}{2\hbar} \int_0^\infty ds \exp\left[\frac{i}{\hbar}\tilde{E}s\right] \frac{1}{\sin(\tilde{\omega}s)} I_\nu\left(\frac{M\tilde{\omega}\xi\xi'}{i\hbar\sin(\tilde{\omega}s)}\right) \exp\left[\frac{iM\tilde{\omega}}{2\hbar}\left(\xi^2+\xi'^2\right)\cot(\tilde{\omega}s)\right]. \qquad (21)$$

By using Gradshteyn's formula [15], with $v > u$,

$$\int_0^\infty dq \frac{e^{-2\lambda q}}{\sinh(q)} I_\nu\left(\frac{(uv)^{1/2}}{\sinh(q)}\right) \exp\left[-\frac{1}{2}(u+v)\coth(q)\right] = \frac{\Gamma\left(\lambda+\frac{\nu}{2}+\frac{1}{2}\right)}{(uv)^{1/2}\Gamma(\nu+1)} M_{-\lambda,\frac{\nu}{2}}(u)\, W_{-\lambda,\frac{\nu}{2}}(v) , \qquad (22)$$

where $M_{-\lambda,\frac{\nu}{2}}(u)$ and $W_{-\lambda,\frac{\nu}{2}}(v)$ are the standard Whittaker functions, the Green's function becomes

$$G(x,x';E) = \frac{-a}{2i\tilde{\omega}} \frac{\Gamma\left(p+\frac{\nu}{2}+\frac{1}{2}\right)}{\Gamma(\nu+1)} \exp\left[\frac{1}{2}a(x+x')\right] M_{-p,\frac{\nu}{2}}\left(De^{-ax'}\right) W_{-p,\frac{\nu}{2}}\left(De^{-ax}\right), \qquad (23)$$

with $x > x'$ and $p = -\frac{\tilde{E}}{2\hbar\tilde{\omega}}$.

This result (21) agrees with the one given in ref. [6].

We succeeded in showing that the Green's function of the Morse potential can be calculated by the so(2,1) algebraic approach. Energies and wave functions may be inferred from the poles of this Green's function in the complex plane.

Acknowledgments

The authors would like to thank Dr. J. Richert, Centre de Recherches Nucléaires, Strasbourg, France, for careful reading of the typescript and useful comments.

References

[1] H. Kleinert, Group Dynamics of the Hydrogen Atom, Lectures presented at the 1967 Boulder Summer School, published in Lectures in Theoretical Physics, vol XB, ed. by A.O. Barut and W.E. Brezin (Gordon and Breach, New York, 1968).

[2] A.I. Milshtein and V.M. Strakhovenko, Phys. Lett. A 90 (1982) 447.

[3] H. Boschi-Filho and A.N. Vaidya, Phys. Lett. A 145 (1990) 69.

[4] T.F. Hammann, G. Oberlechner, G. Trapp, J. Yoccoz, J. Physique 28 (1967) 755 ;
T.F. Hammann and Q. Ho-Kim, Nuovo Cimento 64 B (1969) 367 ;
T.F. Hammann, P.Desgrolard and L. Chetouani, Nuovo Cimento 29 A (1975) 199.

[5] P. Desgrolard and T.F. Hammann, Phys. Rev. C 6 (1972) 482.

[6] J.H. Duru, Phys. Rev. D 28 (1983) 2689.

[7] P.Y. Cai, A. Inomata and R. Wilson, Phys. Lett. A 96 (1983) 117.

[8] H. Kleinert, Phys. Lett. A 120 (1987) 361.

[9] H. Kleinert, Path Integrals in Quantum Mechanics, Statistics and Polymer Physics (World Scientific, Singapore, 1990).

[10] L. Chetouani and T.F. Hammann, Nuovo Cimento 92 B (1986) 106.

[11] B.G. Wybourne, Classical Groups for Physicists (Wiley, New-York, 1974).

[12] J. Schwinger, Phys. Rev. 82 (1951) 664.

[13] I.S. Gradshteyn and I.M. Ryzhik, Table of Integrals, Series and Products (Academic, New-York, N.Y., 1965), p. 959, Eq. (8.440).

[14] Reference [13], p. 952, Eq. (8.406) 1.

[15] Reference [13], p. 729, Eq. (6.699) 4.