

Zeitschrift: Helvetica Physica Acta
Band: 68 (1995)
Heft: 2

Artikel: The three-dimensional BF model with cosmological term in the axial gauge
Autor: Brandhuber, A. / Emery, S. / Landsteiner, K.
DOI: <https://doi.org/10.5169/seals-116732>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The Three-Dimensional BF Model with Cosmological Term in the Axial Gauge

By A. Brandhuber^{#1}, S. Emery^{*2}, K. Landsteiner^{#1} and M. Schweda^{*}

[#] CERN, CH-1211 Genève 23 (Switzerland)

^{*} Institut für Theoretische Physik, Technische Universität Wien
Wiedner Hauptstraße 8-10, A-1040 Wien (Austria)

Abstract. We quantize the three-dimensional BF -model using axial gauge conditions. Exploiting the rich symmetry-structure of the model we show that the Green-functions correspond to tree graphs and can be obtained as the unique solution of the Ward-Identities. Furthermore, we will show that the theory can be uniquely determined by symmetry consideration without the need of an action principle.

1 Introduction

Topological field models [1] of the Schwarz-type [2] have been the subject of continuous investigations over the recent years. These theories are characterized by an invariant action which does not depend on the metric structure of the manifold. Therefore, they are devoid of any local observables. Nevertheless, the metric appears in the gauge-fixing term which is itself a BRST-variation. The variation of the gauge-fixing term with respect to the metric is a BRST-exact quantity implying the existence of a linear vector-like supersymmetry [4] ν_μ in an elegant manner. Together with the BRST-symmetry, the symmetry ν_μ forms an algebra of the form

$$\{s, \nu_\mu\} = \partial_\mu \quad (1.1)$$

¹Supported by the BMWF

²On leave from the Département de Physique Théorique, Université de Genève. Supported by the "Fonds Turrettini" and the "Fonds F. Wurth"

stating that translations are no physical operations and thus reflecting the topological nature of the theory. Thus one might say that this relation lies at the heart of their topological properties.

The most prominent example of these theories is of course the three-dimensional Chern-Simons theory which has led to the powerful connection between link-invariants and the vacuum expectation value of Wilson lines [3]. The Chern-Simons theory has also been studied extensively from a purely field theoretical point of view. It turns out to be a completely ultraviolet finite theory and that this finiteness is a direct consequence of the topological supersymmetry (1.1). Originally, this supersymmetry has been found using the Landau-gauge [4]. In a serie of papers it has been generalized to other gauge-conditions as well [5, 6, 7]. Of particular interest was the case of the axial gauge where it turned out that the topological supersymmetry is not only responsible for the finiteness of the theory, but also allows to compute the Green functions without the use of an action principle [8]. Let us also mention that it has been shown that the topological supersymmetry exists also in string theories [9] and in two-dimensional chiral W_3 -gravity [10] and that it turned out to be an extremely useful tool for solving the descent equations associated with the integrated BRST-cohomology [11].

Another class of Schwarz-type Topological theories [2] are the BF models. Despite of their simple form they reveal a surprisingly rich symmetry structure. Indeed, they allow for reducible invariances [12]. In the particular case of three dimensions [13], another interesting feature is that the B -field is a 1-form and therefore allows the addition of a cubic term in B into the action. This term is usually referred to as a cosmological constant term since then the model is related to three-dimensional Einstein-Hilbert gravity with such a term. A detailed investigation of the symmetry stucture and finiteness properties of this model in the Landau-gauge has been given in [14].

In the case of the Chern-Simons model in the axial gauge, we found that the supersymmetry has two main consequences. Indeed, it turned out to be strong enough for fixing all the Green functions of the theory (in this sense, one can say that it can be substituted to the action principle) and it also imposes the principal value prescription for the propagators. Therefore, it would be desirable to know wether this remains valid for the BF system in the axial gauge [13]. This is precisely the question we will address in this paper and we will show that the answer is positive.

The work is organized as follows. In section 2 we introduce the action and fix the notation and conventions. Section 3 presents the symmetries and all the functional identities. We investigate their consequences for the equations of motion in section 4.1 and for the calculation of the propagators in section 4.2. At the end, we propose some conclusion.

2 The 3D BF model with cosmological term in the axial gauge

The complete action of the 3D BF model containing a cosmological term with an axial gauge fixing is given by

$$S = S_{inv} + S_{gf} , \quad (2.1)$$

with

$$\begin{aligned} S_{inv} &= -\frac{1}{2} Tr \int d^3x \left[\epsilon^{\mu\nu\rho} (F_{\mu\nu} B_\rho + \frac{2\alpha}{3} B_\mu B_\nu B_\rho) \right] , \\ S_{gf} &= Tr \int d^3x \left[b n^\mu A_\mu + d m^\mu B_\mu + \bar{c} n^\mu (D_\mu c + \alpha [B_\mu, \phi]) + \bar{\phi} m^\mu (D_\mu \phi + [B_\mu, c]) \right] . \end{aligned} \quad (2.2)$$

and $D_\mu \dots = \partial_\mu \dots + g[A_\mu, \dots]$ denoting the gauge covariant derivative. $F_{\mu\nu}$ is the field strength of the gauge field A_μ . Further b, d are the Langrange multipliers imposing the gauge-conditions $n^\mu A_\mu = 0$ and $m^\mu B_\mu = 0$ where n^μ and m^μ are it a priori two independent gauge fixing directions. \bar{c}, c and $\bar{\phi}, \phi$ are the anti-ghost and ghost fields corresponding to the two gauge symmetries of S_{inv}

$$\begin{aligned} \delta^1 A_\mu &= -D_\mu \theta , \quad \delta^1 B_\mu = -[B_\mu, \theta] , \\ \delta^2 A_\mu &= -\alpha [B_\mu, \lambda] , \quad \delta^2 B_\mu = -D_\mu \lambda . \end{aligned} \quad (2.3)$$

We choose the gauge group to be simple, all fields belong to the adjoint representation and are written as Lie algebra matrices $\varphi(x) = \varphi^a(x) t_a$, with

$$[t_a, t_b] = f_{ab}^c t_c , \quad Tr(t_a t_b) = \delta_{ab} . \quad (2.4)$$

Finally α is some numerical constant. We summarize the canonical dimensions and the ghost numbers of the various fields in Table 1.

	A	B	b	d	c	\bar{c}	ϕ	$\bar{\phi}$
Dimension	1	1	2	2	0	2	0	2
Ghost number	0	0	0	0	1	-1	1	-1

Table 1: Dimensions and ghost numbers.

3 Symmetries of the action and Ward identities

The action (2.1) is invariant under the BRST transformation s :

$$\begin{aligned}
sA_\mu &= -D_\mu c - \alpha[B_\mu, \phi], & sB_\mu &= -D_\mu \phi - [B_\mu, c], \\
sc &= c^2 + \alpha\phi^2, & s\phi &= \{\phi, c\}, \\
s\bar{c} &= b, & s\bar{\phi} &= d, \\
sb &= 0, & sd &= 0.
\end{aligned} \tag{3.1}$$

Since we are dealing with a topological field theory of Schwarz type, the only metric dependence arises from the gauge fixing part of the action. Therefore, the energy momentum tensor is BRST exact:

$$T_{\alpha\beta} = s\Lambda_{\alpha\beta} \tag{3.2}$$

with

$$\Lambda_{\alpha\beta} = Tr(\eta_{\alpha\beta}\bar{c}n^\rho A_\rho - \bar{c}n_\alpha A_\beta - \bar{c}n_\beta A_\alpha + \eta_{\alpha\beta}\bar{\phi}n^\rho B_\rho - \bar{\phi}n_\alpha B_\beta - \bar{\phi}n_\beta B_\alpha). \tag{3.3}$$

Using the equations of motion, one gets for the divergence of (3.3) the following expression

$$\begin{aligned}
\partial^\alpha \Lambda_{\alpha\beta} &= Tr(\partial_\beta \bar{c} \frac{\delta S}{\delta b} - A_\beta \frac{\delta S}{\delta c} - n^\alpha \bar{c} \varepsilon_{\rho\beta\alpha} \frac{\delta S}{\delta B_\rho} + \partial_\beta \bar{\phi} \frac{\delta S}{\delta b} - B_\beta \frac{\delta S}{\delta \phi} - \\
&\quad - m^\alpha \bar{\phi} \varepsilon_{\rho\beta\alpha} \frac{\delta S}{\delta A_\rho} + n^\alpha \bar{c} \varepsilon_{\rho\beta\alpha} m^\rho d - n^\alpha \bar{\phi} \varepsilon_{\rho\beta\alpha} m^\rho b) + \text{tot. der.}
\end{aligned} \tag{3.4}$$

Integrating (3.4) allows to derive the usual form for the topological supersymmetry only for the case where¹ $n^\mu = m^\mu$ which we will assume for the rest of the paper. Thus we have the following form for the vector supersymmetry transformations ν_α :

$$\begin{aligned}
\nu_\alpha A_\mu &= -\varepsilon_{\alpha\beta\mu} n^\beta \bar{\phi}, & \nu_\alpha B_\mu &= -\varepsilon_{\alpha\beta\mu} n^\beta \bar{c}, \\
\nu_\alpha c &= A_\alpha, & \nu_\alpha \phi &= B_\alpha, \\
\nu_\alpha \bar{c} &= 0, & \nu_\alpha \bar{\phi} &= 0, \\
\nu_\alpha b &= -\partial_\alpha \bar{c}, & \nu_\alpha d &= -\partial_\alpha \bar{\phi}.
\end{aligned} \tag{3.5}$$

The transformations s and the supersymmetry transformations ν_μ form an algebra which closes on-shell:

$$s^2 = \{\nu_\mu, \nu_\nu\} = 0, \quad \{s, \nu_\mu\} = \partial_\mu + \text{Eq. of motion.} \tag{3.6}$$

¹Actually one could also insist in keeping different gauge vectors since the breaking term is BRST exact. This breaking could be controlled by coupling it to an additional source and adding it to the action [7]

In addition there exist two discrete symmetries ² which leave the action (2.1) invariant:

$$c \longleftrightarrow \alpha \bar{c} \quad , \quad \phi \longleftrightarrow \bar{\phi} \quad (3.7)$$

and

$$c \longleftrightarrow \bar{\phi} \quad , \quad \bar{c} \longleftrightarrow \phi \quad (3.8)$$

At the level of the generating functional of the connected Green functions Z_C , all these symmetries leads to a set of WI. The one which correspond to the vector supersymmetry takes the form

$$\begin{aligned} \mathcal{V}_\alpha Z_C = & \text{Tr} \int d^3x \left(J_b \partial_\alpha \frac{\delta Z_C}{\delta J_{\bar{c}}} + \varepsilon_{\alpha\mu\nu} n^\mu J_B^\nu \frac{\delta Z_C}{\delta J_{\bar{c}}} + J_c \frac{\delta Z_C}{\delta J_A^\alpha} + \right. \\ & \left. + J_d \partial_\alpha \frac{\delta Z_C}{\delta J_{\bar{\phi}}} + \varepsilon_{\alpha\mu\nu} n^\mu J_A^\nu \frac{\delta Z_C}{\delta J_{\bar{\phi}}} + J_\phi \frac{\delta Z_C}{\delta J_B^\alpha} \right) = 0. \end{aligned} \quad (3.9)$$

In this formalism, the axial gauge is imposed by the two gauge conditions :

$$\begin{aligned} J_b + n^\mu \frac{\delta Z_C}{\delta J_A^\mu} &= 0, \\ J_d + n^\mu \frac{\delta Z_C}{\delta J_B^\mu} &= 0. \end{aligned} \quad (3.10)$$

As in any linear gauge there exist antighost equations which in the case of the axial gauge are local [5]. In our case we have two of them and they take the following form:

$$\begin{aligned} J_{\bar{c}} - n^\mu \partial_\mu \frac{\delta Z_C}{\delta J_c} + \left[J_b, \frac{\delta Z_C}{\delta J_c} \right] + \alpha \left[J_d, \frac{\delta Z_C}{\delta J_\phi} \right] &= 0, \\ J_{\bar{\phi}} - n^\mu \partial_\mu \frac{\delta Z_C}{\delta J_\phi} + \left[J_b, \frac{\delta Z_C}{\delta J_\phi} \right] + \left[J_d, \frac{\delta Z_C}{\delta J_c} \right] &= 0. \end{aligned} \quad (3.11)$$

Finally, it is well known that in the axial gauge, due to the decoupling of the ghosts, the Slavnov identity which express the invariance of the theory under the BRST-transformations (3.1) takes the form of a dralocal WI. Therefore, one get the two following local gauge WI's

$$\begin{aligned} \partial_\mu J_A^\mu - \left[J_A^\mu, \frac{\delta Z_C}{\delta J_A^\mu} \right] - \left[J_B^\mu, \frac{\delta Z_C}{\delta J_B^\mu} \right] - \left[J_b, \frac{\delta Z_C}{\delta J_b} \right] - \left[J_d, \frac{\delta Z_C}{\delta J_d} \right] - \\ - \left\{ J_c, \frac{\delta Z_C}{\delta J_c} \right\} - \left\{ J_{\bar{c}}, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right\} - \left\{ J_\phi, \frac{\delta Z_C}{\delta J_\phi} \right\} - \left\{ J_{\bar{\phi}}, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right\} + (n \cdot \partial) \frac{\delta Z_C}{\delta J_b} &= 0 \end{aligned} \quad (3.12)$$

and

$$\begin{aligned} \partial_\mu J_B^\mu - \alpha \left[J_A^\mu, \frac{\delta Z_C}{\delta J_B^\mu} \right] - \left[J_B^\mu, \frac{\delta Z_C}{\delta J_A^\mu} \right] - \left[J_b, \frac{\delta Z_C}{\delta J_d} \right] - \alpha \left[J_d, \frac{\delta Z_C}{\delta J_b} \right] - \\ - \alpha \left\{ J_c, \frac{\delta Z_C}{\delta J_\phi} \right\} - \left\{ J_{\bar{c}}, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right\} - \left\{ J_\phi, \frac{\delta Z_C}{\delta J_c} \right\} - \alpha \left\{ J_{\bar{\phi}}, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right\} + (n \cdot \partial) \frac{\delta Z_C}{\delta J_d} &= 0. \end{aligned} \quad (3.13)$$

²In fact, (3.7) and (3.8) imply the existence of additional anti-BRST-like symmetries and anti-vector-like supersymmetries as in [4, 5].

4 Consequences of the Symmetries

4.1 Equations of motion

Let us now investigate in some detail the meaning of the relations of the last section, starting with the projection of the WI for the supersymmetry(3.9), along the gauge fixed direction. Without loss of generality we can choose the gauge vector n^μ to be $(0, 0, 1)$. We will denote this gauge fixed direction by u and the transverse coordinates by $x^{tr} = (x^i)$, $i = 1, 2$. Then,

$$\mathcal{V}_u Z_C = Tr \int d^3x \left(J_b \partial_u \frac{\delta Z_C}{\delta J_{\bar{c}}} + J_c \frac{\delta Z_C}{\delta J_A^u} + J_d \partial_u \frac{\delta Z_C}{\delta J_{\bar{\phi}}} + J_\phi \frac{\delta Z_C}{\delta J_B^u} \right) = 0. \quad (4.1)$$

Taking into account the gauge conditions (3.10), the latter can be written as

$$Tr \int d^3x \left(J^b \mathcal{X} + J^d \mathcal{Y} \right) = 0.$$

where \mathcal{X} and \mathcal{Y} are the most general forms compatible with (4.1)

$$\begin{aligned} \mathcal{X} &= \partial_u \frac{\delta Z_C}{\delta J_{\bar{c}}} + \lambda_1 \left[J^d, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right] + \lambda_2 \left[J^d, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right] + \lambda_3 \left[J^b, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right] + \lambda_4 \left[J^b, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right] - J_c = 0 \\ \mathcal{Y} &= \partial_u \frac{\delta Z_C}{\delta J_{\bar{\phi}}} + \xi_1 \left[J^d, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right] + \xi_2 \left[J^d, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right] + \xi_3 \left[J^b, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right] + \xi_4 \left[J^b, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right] - J_\phi = 0 \end{aligned}$$

At this level, one can use the consistency condition between the two equations we just found and the ghost equation (4.2) in order to fix the arbitrary parameters. Then, the result is

$$\begin{aligned} J_c - n^\mu \partial_\mu \frac{\delta Z_C}{\delta J_{\bar{c}}} + \left[J_b, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right] + \left[J_d, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right] &= 0, \\ J_\phi - n^\mu \partial_\mu \frac{\delta Z_C}{\delta J_{\bar{\phi}}} + \left[J_b, \frac{\delta Z_C}{\delta J_{\bar{\phi}}} \right] + \alpha \left[J_d, \frac{\delta Z_C}{\delta J_{\bar{c}}} \right] &= 0. \end{aligned} \quad (4.2)$$

which are nothing else than the ghost equations. Thus, the equations of motion for the ghost sector are a consequence of the gauge-fixed component of the supersymmetry WI, the gauge condition and the Slavnov identity.

For the gauge sector, let us consider the transverse component of (3.9)

$$\begin{aligned} \mathcal{V}_i Z_C &= Tr \int d^3x \left(J_b \partial_i \frac{\delta Z_C}{\delta J_{\bar{c}}} + \varepsilon_{ji} J_B^j \frac{\delta Z_C}{\delta J_{\bar{c}}} + J_c \frac{\delta Z_C}{\delta J_A^i} + \right. \\ &\quad \left. + J_d \partial_i \frac{\delta Z_C}{\delta J_{\bar{\phi}}} + \varepsilon_{ji} J_A^j \frac{\delta Z_C}{\delta J_{\bar{\phi}}} + J_\phi \frac{\delta Z_C}{\delta J_B^i} \right) = 0. \end{aligned} \quad (4.3)$$

together with the antighost equations (3.11) written as functional operators acting on Z_C

$$\begin{aligned} \left\{ \partial_u \frac{\delta}{\delta J_c} - \left[J_b, \frac{\delta}{\delta J_c} \right] - \alpha \left[J_d, \frac{\delta}{\delta J_\phi} \right] \right\} Z_C &= J_{\bar{c}}, \\ \left\{ \partial_u \frac{\delta}{\delta J_\phi} - \left[J_b, \frac{\delta}{\delta J_\phi} \right] - \left[J_d, \frac{\delta}{\delta J_c} \right] \right\} Z_C &= J_{\bar{\phi}}. \end{aligned} \quad (4.4)$$

Their consistency gives rise to the following identities

$$\begin{aligned} \left\{ \partial_u \frac{\delta}{\delta J_A^i} - \left[J_d, \frac{\delta}{\delta J_A^i} \right] - \alpha \left[J_b, \frac{\delta}{\delta J_B^i} \right] \right\} Z_C &= \varepsilon_{ji} J_B^j - \partial_i J_d, \\ \left\{ \partial_u \frac{\delta}{\delta J_B^i} - \left[J_d, \frac{\delta}{\delta J_B^i} \right] - \left[J_b, \frac{\delta}{\delta J_A^i} \right] \right\} Z_C &= \varepsilon_{ji} J_A^j - \partial_i J_b. \end{aligned} \quad (4.5)$$

which correspond to the equations of motion for the gauge fields. This concludes the analysis of the consequences of the supersymmetry for the equations of motion.

4.2 Calculation of the Propagators

Gauge conditions

We begin by looking at the gauge conditions (3.10) which imply the vanishing of the connected Green functions involving the components A_3 or B_3

$$\begin{aligned} \langle A_3^a(x) \prod_i \varphi_i(x_i) \rangle &= 0 \\ \langle B_3^a(x) \prod_i \varphi_i(x_i) \rangle &= 0 \end{aligned} \quad \forall \varphi_i \quad (4.6)$$

with two exceptions

$$\begin{aligned} \langle A_3^a(x) b^b(y) \rangle &= -\delta^{ab} \delta^{(3)}(x - y), \\ \langle B_3^a(x) d^b(y) \rangle &= -\delta^{ab} \delta^{(3)}(x - y). \end{aligned} \quad (4.7)$$

Antighost equations

The antighost equations (4.2) give the following differential equations for the connected Green functions involving one pair of ghost fields:

$$\partial_{x^3} \langle \bar{c}^a(x) c^b(y) \rangle = \delta^{ab} \delta^{(3)}(x - y), \quad (4.8)$$

$$\partial_{x^3} \langle \bar{\phi}^a(x) \phi^b(y) \rangle = \delta^{ab} \delta^{(3)}(x - y) \quad (4.9)$$

and

$$\begin{aligned} \partial_{x^3} \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_r}(y_r) c^b(y) \bar{c}^a(x) \rangle &= \\ \sum_{j=1}^s f^{ab_j c} \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots \widehat{b^{b_j}(y_j)} \dots b^{b_r}(y_r) c^b(y) \bar{c}^c(y_j) \rangle \delta(x - y_j) &+ \\ \sum_{i=1}^r f^{ac_i c} \langle d^{c_1}(z_1) \dots \widehat{d^{c_i}(z_i)} \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_r}(y_r) c^b(y) \bar{\phi}^c(z_i) \rangle \delta(x - z_i) \end{aligned} \quad (4.10)$$

and

$$\begin{aligned} \partial_{x^3} \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_r}(y_r) c^b(y) \bar{\phi}^a(x) \rangle = \\ \sum_{j=1}^s f^{ab_j c} \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots \widehat{b^{b_j}}(y_j) \dots b^{b_r}(y_r) c^b(y) \bar{\phi}^c(y_j) \rangle \delta(x - y_j) + \\ \alpha \sum_{i=1}^r f^{ac_i c} \langle d^{c_1}(z_1) \dots \widehat{d^{c_i}}(z_i) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_r}(y_r) c^b(y) \bar{c}^c(z_i) \rangle \delta(x - z_i). \end{aligned} \quad (4.11)$$

The solutions of the equations (4.8) and (4.9) are

$$\langle \bar{c}^a(x) c^b(y) \rangle = \delta^{ab} [\theta(x^3 - y^3) + \beta_1] \delta^{(2)}(x^{tr} - y^{tr}), \quad (4.12)$$

$$\langle \bar{\phi}^a(x) \phi^b(y) \rangle = \delta^{ab} [\theta(x^3 - y^3) + \beta_2] \delta^{(2)}(x^{tr} - y^{tr}) \quad (4.13)$$

The form of the terms proportional to β_1 , β_2 is dictated by transverse two dimensional Poincaré invariance and scale invariance. Indeed, the latter forbids solutions of the type $1/(x^{tr} - y^{tr})^2$ because this term is not a well defined distribution. To give it a meaning would need the introduction of UV subtraction point, i.e., of a dimensionful parameter which would break scale invariance. The integration constant can be fixed with the help of the discrete symmetry of the action (3.7) to be $\beta_1 = \beta_2 = -\frac{1}{2}$.

Integration of the equations (4.10) and (4.11) yields the following recursion relations for the Green functions with one pair of ghosts:

$$\begin{aligned} \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_s}(y_s) c^b(y) \bar{c}^a(x) \rangle = \\ = \sum_{j=1}^s f^{ab_j c} [\theta(x^3 - y_j^3) + \beta(r, s)] \delta^{(2)}(x^{tr} - y_j^{tr}) \times \\ \times \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots \widehat{b^{b_j}}(y_j) \dots b^{b_s}(y_s) c^b(y) \bar{c}^c(y_j) \rangle + \\ + \sum_{i=1}^r f^{ac_i c} [\theta(x^3 - z_i^3) + \beta(r, s)] \delta^{(2)}(x^{tr} - z_i^{tr}) \times \\ \times \langle d^{c_1}(z_1) \dots \widehat{d^{c_i}}(z_i) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_s}(y_s) c^b(y) \bar{\phi}^c(z_i) \rangle \end{aligned} \quad (4.14)$$

and

$$\begin{aligned} \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_s}(y_s) c^b(y) \bar{\phi}^a(x) \rangle = \\ = \sum_{j=1}^s f^{ab_j c} [\theta(x^3 - y_j^3) + \beta(r, s)] \delta^{(2)}(x^{tr} - y_j^{tr}) \times \\ \times \langle d^{c_1}(z_1) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots \widehat{b^{b_j}}(y_j) \dots b^{b_s}(y_s) c^b(y) \bar{\phi}^c(y_j) \rangle + \\ + \alpha \sum_{i=1}^r f^{ac_i c} [\theta(x^3 - z_i^3) + \beta(r, s)] \delta^{(2)}(x^{tr} - z_i^{tr}) \times \\ \times \langle d^{c_1}(z_1) \dots \widehat{d^{c_i}}(z_i) \dots d^{c_r}(z_r) b^{b_1}(y_1) \dots b^{b_s}(y_s) c^b(y) \bar{c}^c(z_i) \rangle. \end{aligned} \quad (4.15)$$

Using the discrete symmetry one could produce two additional recursion relations which

are not written explicitly since they are not needed for our calculations. The integration constants are all fixed to $\beta(r, s) = -\frac{1}{2}$ by the discrete symmetry (3.7) and Bose symmetry of the Lagrange multipliers b and d .

Now we will discuss the recursion relations for some special values of r and s .

The case $r = 0$

For this discussion we will use a more symbolic notion i.e. we will drop indices and variables because we only want to find the vanishing Green functions whereas the non vanishing Green functions can always be obtained from the explicit recursion relations (4.14) and (4.15). From (4.14) we get $\langle (b)^s c \bar{c} \rangle = \sum \langle (b)^{s-1} c \bar{c} \rangle$ which ends up after s steps with the $\langle c \bar{c} \rangle$ propagator defined in (4.8). On the contrary $\langle (b)^s \phi \bar{c} \rangle = 0$ since the recursion relation stops with the vanishing Green function $\langle \phi \bar{c} \rangle$. Using again the discrete symmetry we obtain $\langle (b)^s \bar{\phi} c \rangle = 0$ and $\langle (b)^s \phi \bar{\phi} \rangle = \sum \langle (b)^{s-1} \phi \bar{\phi} \rangle$.

The case $s = 0$

In this case we have to use (4.14) and (4.15) iteratively e.g. $\langle (d)^r c \bar{c} \rangle = \sum \langle (d)^{r-1} c \bar{\phi} \rangle = \langle (d)^{r-2} c \bar{c} \rangle = \dots$. The final result depends on whether the recursion relation ends up with $\langle c \bar{c} \rangle$ which gives a non vanishing result or with $\langle c \bar{\phi} \rangle$ which gives zero. Here we only want to compile the zero results:

$\langle (d)^r c \bar{c} \rangle = 0$ and $\langle (d)^r \phi \bar{\phi} \rangle = 0$ if r is an odd integer.

$\langle (d)^r \phi \bar{c} \rangle = 0$ and $\langle (d)^r c \bar{\phi} \rangle = 0$ if r is an even integer.

The Green functions with additional ghosts, gauge fields or Lagrange multipliers vanish in general as a consequence of the antighost equations (4.2):

$$\langle X c \bar{c} \rangle = \langle X c \bar{\phi} \rangle = \langle X \bar{c} \phi \rangle = \langle X \bar{\phi} c \rangle = 0, \text{ unless } X = (b)^m (d)^n. \quad (4.16)$$

Transverse supersymmetry

From equation (4.3) we get further relations along the same lines. For reasons of simplification we use the sloppy notation from the discussion above whenever possible.

The results for the two-point functions are:

$$\langle A_i^a(x) B_j^b(y) \rangle = \varepsilon_{ij} \delta^{ab} [\theta(x^3 - y^3) - \frac{1}{2}] \delta^{(2)}(x^{tr} - y^{tr}), \quad (4.17)$$

$$\langle b^a(x) A_i^b(y) \rangle = -\delta^{ab} [\theta(x^3 - y^3) - \frac{1}{2}] \partial_{x^i} \delta^{(2)}(x^{tr} - y^{tr}), \quad (4.18)$$

$$\langle d^a(x) B_i^b(y) \rangle = -\delta^{ab} [\theta(x^3 - y^3) - \frac{1}{2}] \partial_{x^i} \delta^{(2)}(x^{tr} - y^{tr}) \quad (4.19)$$

and $\langle AA \rangle = \langle BB \rangle = \langle bb \rangle = \langle dd \rangle = \langle dA \rangle = \langle bB \rangle = \langle c\bar{\phi} \rangle = \langle \bar{c}\phi \rangle = 0$. Furthermore we observe that all two-point functions with one ghost and one bosonic field vanish.

For the higher Green functions we obtain the recursion relations

$$\begin{aligned} \langle b^{d_1}(w_1) \dots b^{d_r}(w_r) d^{c_1}(z_1) \dots d^{c_s}(z_s) A_{l_1}^{b_1}(y_1) \dots A_{l_t}^{b_t}(y_t) B_{n_1}^{a_1}(x_1) \dots B_{n_u}^{a_u}(x_u) \left\{ \begin{matrix} A_i^b(y) \\ B_i^b(y) \end{matrix} \right\} \rangle = \\ = \sum_{k=1}^r \partial_{w_k^i} \langle b^{d_1}(w_1) \dots \widehat{b^{d_k}(w_k)} \dots b^{d_r}(w_r) (d)^s (A)^t (B)^u \left\{ \begin{matrix} c^b(y) \\ \phi^b(y) \end{matrix} \right\} \bar{c}^{d_k}(w_k) \rangle + \\ + \sum_{k=1}^s \partial_{z_k^i} \langle (b)^r d^{c_1}(z_1) \dots \widehat{d^{c_k}(z_k)} \dots d^{c_s}(z_s) (A)^t (B)^u \left\{ \begin{matrix} c^b(y) \\ \phi^b(y) \end{matrix} \right\} \bar{\phi}^{c_k}(z_k) \rangle + \\ + \sum_{k=1}^t \varepsilon_{il_k} \langle (b)^r (d)^s A_{l_1}^{b_1}(y_1) \dots \widehat{A_{l_k}^{b_k}(y_k)} \dots A_{l_t}^{b_t}(y_t) (B)^u \left\{ \begin{matrix} c^b(y) \\ \phi^b(y) \end{matrix} \right\} \bar{\phi}^{b_k}(y_k) \rangle + \\ + \sum_{k=1}^u \varepsilon_{in_k} \langle (b)^r (d)^s (A)^t B_{n_1}^{a_1}(x_1) \dots \widehat{B_{n_k}^{a_k}(x_k)} \dots B_{n_u}^{a_u}(x_u) \left\{ \begin{matrix} c^b(y) \\ \phi^b(y) \end{matrix} \right\} \bar{c}^{b_k}(y_k) \rangle. \end{aligned} \quad (4.20)$$

In the following we want to specify these recursion relations for special values of r, s, t and u to demonstrate that all Green functions can be obtained from our recursion relations. All Green functions with one pair of ghosts have been obtained in the previous subsection. Now we want to calculate the remaining Green functions with only bosonic fields and at least one A or B field.

The case $t = u = 0$

We obtain from (4.20):

$$\langle (b)^r (d)^s A \rangle = \sum \partial \langle (b)^{r-1} (d)^s c \bar{c} \rangle + \sum \partial \langle (b)^r (d)^{s-1} c \bar{\phi} \rangle$$

so the calculations breaks down to summing over already known Green functions. The same holds for $\langle (b)^r (d)^s B \rangle$.

The case $r = s = 0$

We obtain

$$\langle (A)^t (B)^u A \rangle = \varepsilon \langle (A)^t (B)^{u-1} c \bar{c} \rangle + \varepsilon \langle (A)^{t-1} (B)^u c \bar{\phi} \rangle$$

Using (4.16) we find $\langle (A)^a (B)^b \rangle = 0$ unless $a = 1$ and $b = 1$ which yields the two-point function $\langle AB \rangle$.

The case $s = u = 0$

In this case the relation (4.20) takes the form:

$$\langle (b)^r(A)^t \begin{pmatrix} A \\ B \end{pmatrix} \rangle = \sum \partial \langle (b)^{r-1}(A)^t \begin{pmatrix} c \\ \phi \end{pmatrix} \bar{c} \rangle + \sum \varepsilon \langle (b)^r(A)^{t-1} \begin{pmatrix} c \\ \phi \end{pmatrix} \bar{\phi} \rangle$$

For $t = 0$ we get the results:

$$\langle (b)^r A \rangle = \sum \partial \langle (b)^{r-1} c \bar{c} \rangle \text{ and } \langle (b)^r B \rangle = 0$$

For $t = 1$:

$$\langle (b)^r AB \rangle = \sum \varepsilon \langle (b)^r \phi \bar{\phi} \rangle \text{ and } \langle (b)^r AA \rangle = 0.$$

The case $r = u = 0$

In this case the relation (4.20) takes the form:

$$\langle (d)^s(A)^t \begin{pmatrix} A \\ B \end{pmatrix} \rangle = \sum \partial \langle (d)^{s-1}(A)^t \begin{pmatrix} c \\ \phi \end{pmatrix} \bar{\phi} \rangle + \sum \varepsilon \langle (d)^s(A)^{t-1} \begin{pmatrix} c \\ \phi \end{pmatrix} \bar{\phi} \rangle$$

For $t = 0$ we get the results:

$$\langle (d)^s A \rangle = 0 \text{ for } s \text{ odd and } \langle (d)^s B \rangle = 0 \text{ for } s \text{ even.}$$

For $t = 1$:

$$\langle (d)^s AA \rangle = 0 \text{ for } s \text{ even and } \langle (d)^s AB \rangle = 0 \text{ for } s \text{ odd.}$$

The case $r = t = 0$

In this case the relation (4.20) takes the form:

$$\langle (d)^s(B)^u \begin{pmatrix} A \\ B \end{pmatrix} \rangle = \sum \partial \langle (d)^{s-1}(B)^u \begin{pmatrix} c \\ \phi \end{pmatrix} \bar{\phi} \rangle + \sum \varepsilon \langle (d)^s(B)^{u-1} \begin{pmatrix} c \\ \phi \end{pmatrix} \bar{c} \rangle$$

For $t = 0$ we get the results:

$$\langle (d)^s A \rangle = 0 \text{ for } s \text{ odd and } \langle (d)^s B \rangle = 0 \text{ for } s \text{ even.}$$

For $t = 1$:

$$\langle (d)^s BA \rangle = 0 \text{ for } s \text{ odd and } \langle (d)^s BB \rangle = 0 \text{ for } s \text{ even.}$$

Gauge invariance

The two gauge symmetries (3.12) and (3.13) do not give a lot of new information besides consistency checks and the fact that all correlators consisting only of the Lagrange multipliers vanish:

$$\langle b^{a_1}(x_1) \dots b^{a_m}(x_m) d^{b_1}(y_1) \dots d^{b_n}(y_n) \rangle = 0 \quad \forall m, n$$

This is the unique solution obeying dimensional and scaling arguments.

5 Concluding Remarks

The main result of our study is that the Green functions of the model are the unique solutions of the Ward-identities defining the theory. Furthermore it turned out that the topological vector supersymmetry imposed a rather unexpected restriction on the *a priori* independent gauge vectors. It is also worth noticing that the Green functions correspond to tree graphs only. Note also that in principle there are loop graphs with external b and d fields only, however, as in Chern-Simons theory [15], the gauge-field and the ghost field contributions to these graphs cancel exactly due to the topological supersymmetry. Having investigated here the three-dimensional BF model it is now natural to apply the axial gauge also to higher dimensional BF models. It would be highly interesting to know whether the methods developed here and in [8] are also applicable to these cases.

6 Acknowledgements

The authors would like to thank O. Piguet for helpful discussions and comments.

References

- [1] D. Birmingham, M. Blau, M. Rakowski and G. Thompson,
Phys. Reports **209** (1991) 129.
- [2] A.S. Schwarz, Baku International Topological Conference, Abstracts, vol. 2, p.345 (1987);
- [3] E. Witten, *Comm. Math. Phys.* **121** (1989) 351.
- [4] F. Delduc, F. Gieres and S. P. Sorella *Phys. Lett.* **225** (1989) 367.
- [5] A. Brandhuber, M. Langer, O. Piguet, M. Schweda and S. P. Sorella,
Phys. Lett. **B300** (1993) 92.
- [6] T. Gajdosik and W. Kummer, *Phys. Rev.* **D48** (1993) 2897.
- [7] K. Landsteiner, M. Langer, M. Schweda and S. Sorella, *Phys. Lett.* **B337** (1994) 294.
- [8] A. Brandhuber, M. Langer, M. Schweda, S. Sorella, S. Emery and O. Piguet,
Helv. Phys. Acta **66** (1993) 551.;
- [9] M. Schweda, S.P. Sorella and M. Werneck de Oliveira, *Phys. Lett.* **B315** (1993) 93.
A. Boresch, M. Schweda and S. Sorella, *Phys. Lett.* **B328** (1994) 36.

- [10] M. Carvalho, L. C. Queiroz Vilar and S.P. Sorella *Algebraic characterization of anomalies in chiral $W(3)$ gravity* preprint CBPF-NF-042-94, hep-th:9408127
- [11] S. P. Sorella *Comm. Math. Phys.* **157** (1993) 231.
S. P. Sorella and M. Werneck de Oliveira, *Int. J. Mod. Phys. A***9** (1994) 2979.
O. Moritsch, S. P. Sorella and M. Schweda, *Class. Quantum Grav.* **11** (1994) 1225.
- [12] C. Lucchesi, O. Piguet, S. P. Sorella, *Nucl. Phys.* **B395** (1993) 325.
- [13] N. Maggiore, P. Provero, *Helv. Phys. Acta* **65** (1992) 993.
- [14] N. Maggiore and S.P. Sorella, *Nucl. Phys.* **B377** (1992) 236.
- [15] C. P. Martin, *Phys. Lett.* **B263** (1991) 69.