

Dissipative mechanics in granular superconductors

Autor(en): **Fazio, R. / Falci, G. / Giaquinta, G.**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **62 (1989)**

Heft 6-7

PDF erstellt am: **22.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-116088>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

DISSIPATIVE MECHANISMS IN GRANULAR SUPERCONDUCTORS

R.Fazio, Faculty of Applied Physics, Delft University, 2600 GA DELFT, The Netherlands, - perm. address : Istituto di Fisica, CATANIA.

G. Falci, Dottorato di Ricerca in Fisica, Università di Catania.

G. Giaquinta, Istituto di Fisica, Facoltà di Ingegneria, Università di Catania, Viale A.Doria 6, 95129 CATANIA, Italy

Abstract: The role played by dissipation in the coherent transition of granular superconductors is approached from a new viewpoint stressing the link between local and global properties of the system. A novel kind of free energy functional is presented;

1. Introduction

In accounting for the role of dissipation in the physics of a single Josephson Junction (SJJ), it is well known that starting from different dissipative mechanisms we are led to a non completely overlapping understandings of the superconductor-insulator transition, which rests, moreover, on limiting values of some constraints [1]. At present the extension to arrays does not look to be equally well developed and to this end the knowledge gained in the theory of granular superconductors (GS) [2] and related experiments [3] offers as a promising starting point. Indeed, in a GS, coherence results from the competition between the Josephson coupling, that determines the ordering of the phases of the single grain superconductive order parameters (singled out by a non vanishing value of an ordering field Ψ) and the electrostatic effects responsible for quantum fluctuations, dissipative tunneling suppressing this last leading to coherence.

It is our purpose to investigate the mutual interplay between the SJJ and the global transitions, stressing the role of dissipation in the former, by a novel kind of free energy accounting on the same footing for the respective ordering fields.

2. The Free Energy functional

The Euclidean effective phase action obtained in [2] describing a Josephson array exhibits the feature that, besides the spatial Josephson coupling between the single grain phases ϕ_i , dissipation couples on the imaginary time axis the differences of the phases ϕ_{ij} across every SJJ. The spatial decoupling is performed by introducing the Hubbard-Stratonovich field $\Psi_i(\tau)$ conjugated to $\exp(i\phi_i)$. We introduce a novel field $\Phi_{ij}(\tau)$, in order to perform "temporal" decoupling, conjugated with $\exp(i\phi_{ij})$. Integrating out the phase degrees of freedom, after a cumulant expansion in power of the two fields, we pervent to the following free energy functional form:

$$F|\Psi_i, \Phi_{ij}| = \sum_{ij} \sum_n |\Psi_i(\omega_n) K_{ij}^{-1} \Psi_j(\omega_n) - g(\omega_n) \delta_{ij} \Psi_i(\omega_n) \Psi_j(\omega_n)| + \\ \sum_{ij} \sum_n |\Phi_{ij}(\omega_n)|^2 - 2\alpha(\omega_n) g(\omega_n) |\Phi_{ij}(\omega_n)|^2 + \\ F_1(\Psi^4) + F_2(\Phi^4) + F_3(\Psi^2 \Phi^2)$$

where K_{ij} is the Josephson matrix and g is the phase-phase correlation function (details are omitted for brevity, they will appear in a forthcoming paper [4]). The fourth order coupling term $F_3(\Psi^2 \Phi^2)$ provides to the aimed link in the sense that it can be viewed as a second order term for the effective free energy describing the Ψ or the Φ transition as alone. The structure of the involved propagators obliges recursion to the RG methodology in accounting for the Φ transition [4] whereas a mean-field approximation is known to lead to reasonable findings as about the phase diagram of the Ψ transition.

3. References

- [1] S.A. Bulgadaev, JETP Lett. 45, 623 (1987).
- [2] U. Eckern, G. Schön and V. Ambegaokar, Phys. Rev. B30, 6914 (1984).
R. Fazio, G. Giaquinta and C. Di Mauro, Progr. Teor. Ph., 76, 592 (1986).
R. Fazio and G. Giaquinta, Phys. Rev. B34, 4909 (1986).
G. Falci, R. Fazio and G. Giaquinta, Phys. Rev. B, 1 May 1989.
- [3] B.G. Orr et al., Phys. Rev. Lett. 56, 378 (1986).
- [4] R. Fazio, G. Falci and G. Giaquinta, to be published.