

# Nonperturbatively renormalizable quantum field theories in 2+1 dimensions

Autor(en): **Kovner, A. / Rosenstein, B. / Gat, G.**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **65 (1992)**

Heft 2-3

PDF erstellt am: **19.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-116459>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## Nonperturbatively Renormalizable Quantum Field Theories in 2+1 Dimensions

A.Kovner, B. Rosenstein and G. Gat

Physics Department  
University of British Columbia  
Vancouver, B.C.  
Canada, V6T 2A6

**Abstract.** Recently several 2+1 dimensional models were found which are non-renormalizable in weak coupling expansion but renormalizable in  $1/N$ . We show that the IR fixed points of Yukawa theories and scalar  $QED_3$  (Landau-Ginzburg) are some of these theories. The critical 4-Fermi interaction model is the IR stable fixed point of Yukawa theories while the  $CP^{N-1}$  model is the IR fixed point of N-component scalar  $QED_3$ .

As is well known, there is a close relation between renormalizable relativistic quantum field theories (QFT) in 2+1 dimensions and second order phase transitions in 3D. For (rotationally invariant) statistical systems universal critical properties are determined by an infrared (IR) fixed point of the corresponding QFT. For example, critical exponents of Heisenberg (anti)ferromagnets are described by the  $O(3)$  symmetric  $\phi_3^4$  [1] theory

$$S_E[\phi] = \frac{1}{2}(\partial\phi_i)^2 - \frac{\mu^2}{2}\phi_i^2 + \frac{\lambda}{4!}\phi_i^4. \quad (1)$$

The IR fixed point of  $\phi^4$  is the conformally invariant critical  $\sigma$ -model. Note that although the nonlinear  $\sigma$ -model is non renormalizable in weak coupling perturbation theory it is renormalizable nonperturbatively.

Recently several new classes of nonperturbatively renormalizable QFT's in 2+1 dimensions were found using the  $1/N$  expansion. These include four-Fermi interaction models like the  $Z_2$  chiral invariant Gross-Neveu (GN) model [2]

$$\mathcal{L} = \bar{\psi}_i \not{\partial} \psi_i + \frac{g^2}{2N} (\bar{\psi}_i \psi_i)^2. \quad (2)$$

We show that the critical GN model is the IR fixed point of the superrenormalizable Yukawa theories

$$\mathcal{L} = \frac{1}{2}(\partial\sigma)^2 + \bar{\psi}_i \not{\partial} \psi_i + \lambda_0 \sigma \bar{\psi}_i \psi_i + \frac{m_0^2}{2} \sigma^2 \quad (3)$$

and therefore determines its universality class. At the phase transition point the chiral  $Z_2$  breaks down spontaneously. The renormalized coupling of the Yukawa model is determined by (we use notations of [1])

$$\sqrt{m}\lambda = \Gamma_{1,2}(p_1 = 0, p_2 = 0, p_3 = 0) \quad (4)$$

where  $m$  is the boson's mass,  $\Gamma_b^2(m) = 0$ . To fix the field's normalizations we choose to impose  $\frac{\partial \Gamma_b^2}{\partial p^2} \Big|_{p^2=m^2} = 1$ ,  $\frac{\partial \Gamma_f^2}{\partial p^2} \Big|_{p^2=0} = 1$ . To leading order in  $1/N$  the corresponding Callan-Symanzik  $\beta$  function is

$$\beta(\lambda) = -\frac{1}{2}\lambda + \frac{1}{16}\lambda^3 \quad (5)$$

It has a finite IR fixed point at  $\lambda^* = 2\sqrt{2}$ . The connection between the bare and renormalized couplings is determined by the RG equation

$$\left(1 + \beta(\lambda) \frac{\partial}{\partial \lambda}\right) \frac{\lambda_0^2}{m} = 0 \quad (6)$$

The solution is

$$\lambda_0(\lambda) = \sqrt{m}\lambda \exp \left\{ -\frac{1}{2} \int_0^\lambda \left[ \frac{1}{\beta(x)} + \frac{1}{x} \right] dx \right\} \quad (7)$$

As  $\lambda \rightarrow \lambda^*$ , the bare coupling  $\lambda_0$  diverges  $\lambda_0 \rightarrow \infty$ . In this limit the Yukawa action becomes the GN action eq.(2).

This is seen explicitly by calculating the dynamical dimension of the kinetic term for  $\sigma$ :  $[(\partial\sigma)^2] = 4$  [3]. It becomes irrelevant for  $d = 3$ . On the other hand some irrelevant operators at weak coupling become relevant or marginal near the fixed point. The dimension four  $(\bar{\psi}\psi)^2$  operator acquires scaling dimension two and thus becomes relevant, while the dimension six operator  $(\bar{\psi}\psi)^3$  has in the leading order in  $1/N$  scaling dimension three and thus becomes marginal. In this case more careful analysis is needed. The next to leading order in  $1/N$  shows that in fact it becomes relevant. Similar phenomenon occurs in abelian gauge theories. In scalar QED the IR fixed point is the nonperturbatively renormalizable  $CP^{N-1}$  model [4]. It is interesting to note that the vortex operator  $V$  that creates Nielsen-Olesen vortex [5] becomes relevant at  $N < 8$ .

To summarize we found that recently discovered nonperturbatively renormalizable 2+1 dimensional QFT contain IR fixed points describing new universality class. It is interesting to know whether these universality classes are realized in real materials.

The work of A.K and B.R was supported by NSERC of Canada.

### References

- [1] C. Itzykson and J.M. Drouffe, *Statistical Field Theory*, (Cambridge University Press, 1991) vol. I
- [2] B. Rosenstein, B.J. Warr and S.H. Park *Phys.Rev.Lett.* **62**, 1433 (1989) ; B. Rosenstein, B.J. Warr and S.H. Park *Phys.Rep.* **205**, 59 (1991) and references therein ; C. de Calan et al, *Phys.Rev.Lett* **66**, 3233 (1991)
- [3] Y. Kikukawa and K. Yamawaki, *Phys.Lett B* **234**, 497 (1990) ; G. Gat et al, *Phys.Lett. B* **240**,158 (1990)
- [4] A. Kovner and B. Rosenstein *Phys.Lett B* **261**,97 (1991)
- [5] A. Kovner and B. Rosenstein *Phys.Lett B* **261**,104 (1991)