

Chaotic dynamics of fluxons in large-area Josephson junctions

Autor(en): **Abdullaev, F.Kh. / Umarov, B.A.**

Objektyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **65 (1992)**

Heft 2-3

PDF erstellt am: **24.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-116498>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

CHAOTIC DYNAMICS OF FLUXONS IN LARGE-AREA JOSEPHSON JUNCTIONS

F.Kh.Abdullaev, B.A.Umarov

Thermophysics Department of the Uzbek Academy of Sciences, Katartal str.28, Tashkent 700135, USSR

Last years the investigation of the fluxon dynamics in the Josephson junctions attracts much interest of scientists. It is connected with a possibility to control the fluxon parameters with the help of external effects or the changes of junction parameters [1-3]. In this paper we consider the fluxon motion in large-area inhomogeneous Josephson junction at the presence of external variable current.

Let us consider a motion of the radial-symmetric vortex in inhomogeneous junction under the action of high frequency external force. In this case the equation for the phase difference between two superconducting films in standard dimensionless variables [1] takes the form

$$\Phi_{tt} - \Phi_{rr} + \sin\Phi = f \sin \Omega t + 1/r \Phi_r - \varepsilon \alpha(\Phi, r). \quad (1)$$

where α describes of the inhomogeneity influence.

Following [4-5] let us consider the dynamics of kink-shaped bubble satisfying (1) in which the energy is concentrated in a single well. We assume that the radius of such a bubble is large enough to be considered as a perturbation. For $\varepsilon=0, f=0$, and $\rho \gg 1$, one obtains the fluxon solution in the form

$$\Phi_0(z) = 4 \tanh^{-1} \exp\left(\frac{z}{\rho}\right), \quad z = \frac{r-\rho}{(1-v^2)^{1/2}}, \quad \rho = vt + \rho_0,$$

where v is fluxon velocity and ρ is the radius.

We apply perturbation theory to the fluxon dynamics by taking right hand side in (1) as a small perturbation. Then one obtains

$$\frac{dv}{dt} = \frac{1}{\rho} + \varepsilon \alpha(\rho) + \pi f / [8 \cdot (1-\Omega^2)] \cdot \sin \omega t, \quad (2)$$

$$\frac{d\rho}{dt} = v(t), \quad (3)$$

where

$$a(\rho) = \frac{1}{4} \int \alpha(z, \rho) \operatorname{sech}^4 z \, dz.$$

Hereby we came to nonrelativistic limit $v^2 \ll 1$ and take into account the renormalization of the external field following [2]. Let us investigate the case when $\varepsilon \gg f$, and $1/\rho \sim f$. In this case the system (2),(3) can be studied with the account of radially and external current as a perturbation. Let us consider the truncated system (2),(3)

$$dv/dt = \varepsilon a(\rho), \quad (2')$$

$$d\rho/dt = v. \quad (3')$$

Let us consider concrete models of the inhomogeneity

$$\alpha(r) = \delta(r - r_1) + \delta(r - r_2). \quad (4)$$

This is a model of the Josephson junction with micro short-circuit the regions of strong superconducting currents, the sizes of which are much less than Josephson depth of penetration λ_J . The separatrix solution of the system (2'-3') given in non explicit form, is found from the equation [6]

$$2 H_c (t - t_0) = \rho' - \frac{\alpha + y_0}{2(y_0^2 - 1)^{1/2}} \ln \left[\frac{y_0^{-1} + (y_0^2 - 1)^{1/2} \operatorname{tanh} \rho'}{y_0^{-1} - (y_0^2 - 1)^{1/2} \operatorname{tanh} \rho'} \right], \quad (5)$$

$$y_0 = \alpha - 2/\alpha, \quad \alpha = \cosh(r_2 - r_1), \quad \rho' = \rho - \frac{r_1 + r_2}{2},$$

$$H_c = \frac{\varepsilon}{4} \frac{\alpha^2}{(\alpha^2 - 1)}.$$

Taking into account the separatrix solution one can find the criteria of the chaotic motion near the separatrix with the help of the Melnikov method [7]. Let us calculate the Melnikov integral $D(t_0)$. If $\omega_1 \gg 1$, we obtain

$$D(t_0) \approx \varepsilon \cdot a \cdot \sin(k+L) \cdot \sin \omega t_0 / [2\omega_1 \cdot (1 - \Omega^2)] - \ln(r_1/r_2), \quad (6)$$

where $\omega_1 = \Omega / 2H_c$, $r_2 > r_1 \gg 1$, $L = r_2 - r_1$. For $1 \ll \rho_0$ we have

$$\ln(r_1/r_2) = 1/\rho_0.$$

It is known that if $D(t_0)$ changes a sign, then the separatrix is intersect and a motion becomes chaotic in this region. Having studied the result presented in formulae (6) we came to conclusion that in general case the fluxon dynamics in large-area Josephson junctions under the action of variable external field can have a stochastic character and the account of radial symmetry results in narrowing of the chaos region, i.e it acts as a dumping. This conclusion should be taken into account while designing the Josephson oscillators with variable frequency, in that case when we apply the alternating current to change the frequency.

References.

1. D.W. McLaughlin, A.C. Scott. In Solitons in action. Acad. Press, N.Y. 1978.
2. F.Kh. Abdullaev, A.A. Abdumalikov, B.A. Umarov. Doklady Uzbek Academy of Sciences. N.3, p.25, 1989. (In Russian)
3. P.L. Christiansen, P.S. Lomdahl, N.J. Zabusky. Appl. Phys. Lett. v.39, N.2, p.170, 1981.
4. E.M. Maslov. Preprint IZMIRAN SSSR, N.15(769), 1989.
5. E.M. Maslov. Physica D15, p.433, 1985.
6. F.Kh. Abdullaev, S.A. Darmanyan, B.A. Umarov. Phys. Rev. A, v.41, N.8, p.4498, 1990.
7. V.K. Melnikov. Trudy Moskovsk. Mat. Obshestva. v.12, p.3, 1963. (In Russian)