

Zeitschrift: Mitteilungen aus dem Gebiete der Lebensmitteluntersuchung und Hygiene = Travaux de chimie alimentaire et d'hygiène

Herausgeber: Bundesamt für Gesundheit

Band: 81 (1990)

Heft: 3

Artikel: Influence of menthol and thymol on *Aspergillus parasiticus* growth and production of aflatoxins in a synthetic medium

Autor: Karapinar, M.

DOI: <https://doi.org/10.5169/seals-982616>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Influence of Menthol and Thymol on *Aspergillus parasiticus* Growth and Production of Aflatoxins in a Synthetic Medium

M. Karapinar

Ege University, Faculty of Engineering, Food Engineering Department, Bornova-Izmir

Introduction

Aflatoxins are metabolites of the molds *Aspergillus flavus* and *A. parasiticus* which can grow on a wide variety of agricultural commodities and frequently contaminate foods and feeds inducing undesirable effects in both man and animals (1–5). There is no practical method to eliminate toxigenic molds from human and animal food resources, hence it is desirable to prevent aflatoxin formation or to inactivate aflatoxins in contaminated materials. Inhibition of the growth and the synthesis of aflatoxins by several means is the best approach to prevent contamination of foods with aflatoxins since their inactivation is very difficult (4, 6–10).

Recently, there has been considerable emphasis on the studies involving spices, spice oils and their constituents for inhibiting the growth and the synthesis of aflatoxins (11–22).

Menthol is the active principle of the essential oil of peppermint, and thymol is mainly present in the essential oils of thyme and origanum (17, 22–24).

This investigation was undertaken to determine growth and toxin production by *A. parasiticus* when menthol and thymol were introduced into a liquid medium.

Materials and Methods

Organisms

Strains of *Aspergillus parasiticus* NRRL 2999 and *Aspergillus parasiticus* CBS 26027 (ATCC 15517) were used. They were maintained on potato dextrose agar (PDA, Oxoid) slopes and stored in refrigerator.

Culture techniques

Yeast Extract Sucrose medium (YES) prepared according to *Davies et al.* (25) was used as basal medium throughout the study. The broth was dispensed in 10 ml quantities into test tubes and sterilized by autoclaving at 115 °C for 15 min. After sterilisation appropriate amounts of pure menthol, art. no 5995 and thymol, art. no. 8167 (Merck, Darmstadt, FRG) diluted in 1 ml of ethanol were added to separate tubes to give concentrations of 25, 50, 100, 150, 200, 300, 400 and 500 µg per ml. Ethanol containing no test material was added to culture media as a control. 0.1 ml portions of spore suspensions which were prepared according to the method described previously (21) were inoculated into 10 ml of culture medium. The tube cultures were slanted and incubated statically at 30 °C for 10 days. Experiments were carried out in duplicate.

Analyses

Mold growth was observed visually throughout the incubation period. After the incubation the cultures were autoclaved at 121 °C for 30 seconds to kill spores and mycelia. The mycelial weights were determined after filtering cultures through a sintered glass crucible No. 1 (Gallenkamp, Loughborough, U. K.) washing with distilled water and oven drying at 95 °C for 24 h.

Culture filtrates taken before washing were analysed for aflatoxin production according to the method of *Hitokoto et al.* (15), with use of Silica G-HR (Merck, Darmstadt, FGR) plates for thinlayer chromatography. The intensities of fluorescence of the separated aflatoxins were evaluated visually under UV at 366 nm.

Results and Discussion

Inhibition of growth by menthol and thymol was estimated qualitatively and quantitatively. Qualitative estimations were made visually using the scale explained at the bottom of table 1, whereas quantitative estimations were made on the basis of mycelial weight.

Although, as in the control tubes, growth of both strains initiated on the first day of incubation, growth rates were slowed down on media containing 25, 50, 100 µg/ml menthol (table 1). With level of menthol above 100 µg/ml, growth initiation retarded in both strains.

Growth inhibition increased with increasing concentrations of menthol and growth of both strains was completely inhibited on media containing 600 µg/ml. Similar results were observed when the dry weight of mycelial mats was used as a measure of growth (table 2).

At all concentrations, menthol caused slighter inhibition on the growth of *A. parasiticus* CBS 26027. Compared with the strain NRRL 2999 however, it will

also be seen from table 2 that up to the concentration of 300 $\mu\text{g/ml}$, menthol stimulated the aflatoxin production of *A. parasiticus* NRRL 2999, whereas its toxin-inhibiting activity against *A. parasiticus* CBS 26027 started at the level of 100 $\mu\text{g/ml}$ (57% inhibition).

Table 1. Growth of *A. parasiticus* in the presence of menthol (visual estimation)

Level of menthol ($\mu\text{g/ml}$)	Incubation time (days) at 30 °C									
	1	2	3	4	5	6	7	8	9	10
<i>A. parasiticus</i> NRRL 2999										
0	±	±	++							
25	±	±	+	++						
50	±	±	+	++						
100	±	±	±	+	++					
200	-	±	±	±	+	+	++			
300	-	-	-	±	±	±	±	±	+	++
400	-	-	-	-	-	-	-	±	±	±
500	-	-	-	-	-	-	-	-	±	±
600	-	-	-	-	-	-	-	-	-	-
<i>A. parasiticus</i> CBS 26027										
0	±	+	++							
25	±	+	++							
50	±	±	+	++						
100	±	±	±	+	++					
200	-	-	±	±	+	++				
300	-	-	-	-	±	±	±	+	+	++
400	-	-	-	-	-	±	±	±	±	+
500	-	-	-	-	-	-	±	±	±	±
600	-	-	-	-	-	-	-	-	-	--

- = no growth; ± = start of growth; + = scant growth; ++ = abundant growth

At concentrations of 200, 300, 400 and 500 $\mu\text{g/ml}$ menthol respectively caused ca. 67, 98, 99, 100% inhibition of the aflatoxin production of *A. parasiticus* CBS 26027. The percentage inhibition of the aflatoxin formation by *A. parasiticus* NRRL 2999 was ca. 31, 40 and 98 for menthol levels of 300, 400 and 500 $\mu\text{g/ml}$, respectively. Menthol, at a concentration of 500 $\mu\text{g/ml}$, seems to be critical for both strains and it dramatically suppressed growth and toxin production.

Table 2. Inhibitory effects of menthol on growth and toxin production of *A. parasiticus* after 10 days at 30 °C

Menthol ($\mu\text{g/ml}$)	Mycelium (mg/ml) ^a	A. parasiticus NRRL 2999				
		Aflatoxins ($\mu\text{g/ml}$)				
		B ₁	B ₂	G ₁	G ₂	Total
0	20 (0) ^b	273	21	196	20	510 (0)
25	17 (15)	581	51	458	34	1124 (0)
50	17 (15)	555	52	357	34	998 (0)
100	19 (5)	515	44	250	41	850 (0)
200	14 (30)	365	30	142	24	561 (0)
300	13 (35)	301	30	11	10	351 (31)
400	Tr	210	24	57	13	304 (40)
500	Tr	8	1	2	0	11 (98)
600	0 (100)	0	0	0	0	0 (100)
		A. parasiticus CBS 26027				
		Aflatoxins ($\mu\text{g/ml}$)				
		B ₁	B ₂	G ₁	G ₂	Total
0	21 (0) ^b	89	9	53	14	165 (0)
25	20 (5)	92	4	72	3	171 (0)
50	19 (10)	132	6	49	4	191 (0)
100	20 (5)	55	3	13	0	71 (57)
200	15 (29)	48	1	6	0	55 (67)
300	12 (43)	6	0	1	0	7 (96)
400	2 (90)	1	0	0	0	1 (99)
500	Tr	0	0	0	0	0 (100)
600	0 (100)	0	0	0	0	0 (100)

a = Dry weight; b = Numbers in parentheses indicate percentage inhibition;
Tr = Trace growth

However, initiation of growth at the end of a 10 days incubation period was observed as seen from tables 1 and 2 which indicate that the organisms may overcome the inhibitory effects of menthol at this level. Peppermint contains 0.5–3.0% volatile oil with 50–78% menthol (16, 26). *Mabrouk* and *El-Shayeb* (16) tested the antimicrobial activity of several spices including mint against growth and toxin production of *Aspergillus flavus*. They reported that, although mint stimulated mycelial growth, it inhibited aflatoxin formation at the levels of 5.0 and 10.0%. Results in table 3 and 4 indicate that *A. parasiticus* CBS 26027 is more sensitive to thymol than *A. parasiticus* NRRL 2999. As it can be seen from table 4, antimicrobial activity of thymol against growth and toxin production of *A. parasiticus* CBS 26027 started at the lowest level and the amount of mycelia and to-

xins decreased by increased concentration of thymol whereas lower concentrations of thymol, up to 100 $\mu\text{g/ml}$, enhanced the aflatoxin production of *A. parasiticus* NRRL 2999.

Table 3. Growth of *A. parasiticus* in the presence of thymol (visual estimation)

Level of thymol ($\mu\text{g/ml}$)	Incubation time (days) at 30 °C									
	1	2	3	4	5	6	7	8	9	10
<i>A. parasiticus</i> NRRL 2999										
0	+	+	++							
25	+	+	+	++						
50	-	+	+	+	+	++				
100	-	-	-	+	+	+	+	+	+	++
150	-	-	-	-	-	-	-	-	-	-
<i>A. parasiticus</i> CBS 26027										
0	+	+	++							
25	+	+	+	++						
50	-	+	+	+	+	++				
100	-	-	+	+	+	+	+	+	+	+
150	-	-	-	-	-	-	-	-	-	-

However, thymol at a concentration of 150 $\mu\text{g/ml}$ completely inhibited the growth of both strains. Thymol is a major constituent of the essential oil of thyme and origanum and the thyme contains 0.4–2.5% volatile oil with 20–60% thymol (15, 26).

Qualitatively the results are similar to those reported by Hitokoto et al. (15) Buchanan and Shepherd (17) and Farag et al. (22) who tested the antimicrobial activity of thymol against *A. flavus* and *A. parasiticus*. However, the *A. parasiticus* strains used in this study were substantially more sensitive to thymol than the *A. flavus* and *A. parasiticus* strains employed in earlier studies by other workers (15, 22). Hitokoto et al. (15) reported that a thymol concentration above 200 $\mu\text{g/ml}$ was necessary to inhibit the growth and toxin production of *A. flavus*. Similarly Farag et al. (22) who tested the inhibitory effects of thyme oil and thymol on the growth and toxin production of *A. parasiticus* ATCC 120920 have reported that a medium containing either 400 $\mu\text{g/ml}$ thyme oil or thymol completely inhibited the growth. It has been reported that a compound is considered a positive inhibitor if it reduces aflatoxin formation to 50% of that of control (16). Hence our results as well as other workers show that menthol and thymol if used in suf-

efficient quantities may be considered as effective inhibitors of aflatoxigenic fungi and consequently may offer some assistance in preventing of aflatoxin production.

Table 4. Inhibitory effects of thymol on the amount of growth and toxin production of *A. parasiticus* after 10 days at 30 °C

Thymol ($\mu\text{g/ml}$)	Mycelium (mg/ml) ^a	A. parasiticus NRRL 2999				
		Aflatoxins ($\mu\text{g/ml}$)				
		B ₁	B ₂	G ₁	G ₂	Total
0	20 (0) ^b	273	21	196	20	510 (0)
25	35 (0)	1124	53	253	30	1460 (0)
50	37 (0)	995	84	194	36	1309 (0)
100	14 (30)	124	9	14	19	166 (67)
150	0 (100)	0	0	0	0	0
	Mycelium (mg/ml) ^a	A. parasiticus CBS 26027				
		Aflatoxins ($\mu\text{g/ml}$)				
		B ₁	B ₂	G ₁	G ₂	Total
0	21 (0) ^b	89	9	53	14	165 (0)
25	20 (5)	88	2	11	1	102 (38)
50	18.7 (10)	28	2	5	1	36 (78)
100	3 (86)	2	0	0	0	2 (99)
150	0 (100)	0	0	0	0	0 (100)

a = Dry weight; b = Numbers in parentheses indicate percentage inhibition

Acknowledgement

The financial support of Ege University Research Foundation is gratefully acknowledged.

Summary

The antimicrobial and antitoxigenic activity of menthol and thymol which are active components of commonly used spices was investigated against two aflatoxigenic strains of *Aspergillus parasiticus*. Of the active components tested, thymol was the most effective against both strains leading to definite growth inhibition at the level of 150 $\mu\text{g/ml}$.

However, *A. parasiticus* CBS 26027 was more sensitive to thymol than the other strain. The growth and aflatoxin inhibition in this strain started at the lowest level of thymol (25 $\mu\text{g/ml}$); whereas low levels of thymol caused stimulation in growth and toxin accumulation by *A. parasiticus* NRRL 2999.

Both strains showed a similar inhibition pattern against menthol at the level of 600 µg/ml.

Zusammenfassung

In dieser Arbeit wurden die antimikrobiellen und antitoxigenen Einflüsse von Menthol und Thymol, die aktive Bestandteile von häufig verwendeten Gewürzen sind, bei zwei aflatoxigenen Stämmen von *Aspergillus parasiticus* untersucht. Thymol erwies sich bei beiden Stämmen als besonders wirksam.

Das Wachstum beider Stämme wurde bei einer Dosis von 150 µg/ml vollständig gehemmt, wobei *A. parasiticus* CBS 26027 empfindlicher reagierte. Bei *A. parasiticus* CBS 26027 begann das Schimmelpilzwachstum bei gleichzeitiger Hemmung der Toxinproduktion auf dem niedrigsten Niveau von Thymol (25 µg/ml). Paradoxaerweise stimulierte dieselbe Thymolkonzentration das Wachstum und die Toxinbildung von *A. parasiticus* NRRL 2999.

Beide Stämme zeigten eine ähnliche Empfindlichkeit gegenüber Menthol, wobei ihr Wachstum bei 600 µg/ml vollständig gehemmt wurde.

Résumé

Cette étude a pour but d'examiner l'effet antimicrobien et antitoxigénique des deux principes actifs que sont le menthol et le thymol, que l'on trouve dans certaines épices, à l'encontre de deux isolats aflatoxigènes de *Aspergillus parasiticus*.

Le thymol s'est révélé particulièrement efficace: une solution de 150 µg/ml inhibe complètement la croissance de ces deux souches; cependant c'est *A. parasiticus* CBS 26027 qui a été le plus sensible au thymol.

La croissance de la souche et l'inhibition de la production d'aflatoxines ont commencé vers une concentration minimale de 25 µg/ml de thymol. Tandis que, paradoxalement, cette même concentration a provoqué chez *A. parasiticus* NRRL 2999 une stimulation de sa croissance et de sa production de toxines. Les deux champignons ont réagi de manière similaire à l'encontre du menthol, une concentration de 600 µg/ml de menthol a inhibé leur croissance.

Literature

1. Denizel, T. and Kösker, Ö.: A mycological survey of various kinds of edible nuts commercially available in the U. K. with reference to mycotoxins. University of Ankara, Yearbook of the Faculty of Agriculture, 168–169 (1972).
2. Stolof, L.: Occurrence of mycotoxins in foods and feeds. Adv. Chem. Ser. **149**, 23–50 (1976).
3. Jewers, K.: Mycotoxins in food the application of survey and quality control. R. Soc. Health S. **102**, 114–118 (1982).
4. Bullerman, L. B., Schroeder, L. L. and Park, K. Y.: Formation and control of mycotoxins in food. J. Food Prot. **47**, 637–646 (1984).

5. Sanchis, V., Sala, N., Palones, A., Santamarima, A. and Burdaspal, P. A.: Occurrence of aflatoxin and aflatoxigenic molds in foods and feed in Spain. *J. Food Prot.* **49**, 445–448 (1986).
6. Jarvis, B.: Factors affecting the production of mycotoxins. *J. Appl. Bacteriol.* **34**, 199–213 (1971).
7. Hesseltine, C. W.: Conditions leading to mycotoxin contamination of foods and feeds. *Adv. Chem. Ser.* **49**, 1–22 (1976).
8. Goldblatt, L. A. and Dollear, F. G.: Review of prevention, elimination and detoxification of aflatoxins. *Pure Appl. Chem.* **49**, 1759–1764 (1977).
9. Northolth, M. D. and Bullerman, L. B.: Prevention of mold growth and toxin production through control of environmental conditions. *J. Food Prot.* **45**, 519–526 (1982).
10. Bullerman, L. B.: Interactive effects of temperature and pH on mycotoxin production. *Lebensm. Wiss. Technol.* **18**, 197–200 (1985).
11. Bullerman, L. B.: Inhibition of aflatoxin production by cinnamon. *J. Food Sci.* **39**, 1163–1165 (1974).
12. Alderman, G. G. and Marth, E. M.: Inhibition of growth and aflatoxin production of *Aspergillus parasiticus* by citrus oils. *Z. Lebensm. Unters.-Forsch.* **160**, 353–358 (1976).
13. Bullerman, L. B., Lieu, F. Y. and Seier, S. A.: Inhibition of growth and aflatoxin production by cinnamon and clove oils, cinnamic aldehyde and eugenol. *J. Food Sci.* **42**, 1107–1109 and 1116 (1977).
14. Sharma, A., Tewari, G. M., Shrihande, A. J., Padwal-Desai, S. R. and Bandyopadhyay, C.: Inhibition of aflatoxin-producing fungi by onion extracts. *J. Food Sci.* **44**, 1545–1547 (1979).
15. Hitokoto, H., Morozumi, S., Wauke, T., Sakai, S. and Kurata, H.: Inhibitory effects of spices on growth and toxin production of toxigenic fungi. *Appl. Environm. Microbiol.* **89**, 818–822 (1980).
16. Mabrouk, S. S. and El-Shayeb N. M. A.: Inhibition of aflatoxin formation by some spices. *Z. Lebensm. Unters. -Forsch.* **171**, 344–347 (1980).
17. Buchanan, R. L. and Shepherd, A. J.: Inhibition of *Aspergillus parasiticus* by thymol. *J. Food Sci.* **46**, 976–977 (1981).
18. Llewellyn, G. C., Burkett, M. L. and Eadie, T.: Potential mold growth, aflatoxin production and antimycotic activity of selected natural spices and herbs. *J. Assoc. Off. Anal. Chem.* **64**, 955–960 (1981).
19. Mabrouk, S. S. and El-Shayeb, N. M. A.: The effects of garlic on mycelial growth and aflatoxin formation by *Aspergillus flavus*. *Chem. Microbiol. Technol. Lebensm.* **7**, 37–41 (1981).
20. Azzous, M. A. and Bullerman, L. B.: Comparative antimycotic effects of selected herbs, spices, plant components and commercial antifungal agents. *J. Food Prot.* **45**, 1298–1301 (1982).
21. Karapinar, M.: The effect of citrus oils and some spices on growth and aflatoxin production by *Aspergillus parasiticus* NRRL 2999. *Inter. J. Food Microbiol.* **2**, 299–245 (1985).
22. Farag, R. S., Daw, Z. Y. and Abo-Raya, S. H.: Influence of some spice essential oils on *Aspergillus parasiticus* growth and production of aflatoxins in a synthetic medium. *J. Food Sci.* **54**, 74–76 (1989).
23. Parry, J. W.: Spices morphology, histology, chemistry. Vol. 2, p. 183. Chemical Publishing Comp. Inc, New York 1969.
24. Shelef, L. A.: Antimicrobial effects to spices. *J. Food Safety* **6**, 29–44 (1980).

25. *Davies, N. O., Diener, M. S. and Eldrige, W. B.*: Production of aflatoxin B₁ and G₁ by *Aspergillus flavus* in a semi-synthetic medium. *Appl. Microbiol.* **14**, 378–380 (1966).
26. *Egan, H., Kirk, R. and Sawyer, R.*: Pearson's chemical analysis of foods, p. 317. Churchill Livingstone, Edinburgh 1981.

Assoc. Prof. Dr. M. Karapinar
Ege University, Engineering Faculty
Food Engineering Department
Bornova-Izmir
Turkey TR-35100