

<b>Zeitschrift:</b>	Mitteilungen / Vereinigung Schweizerischer Versicherungsmathematiker = Bulletin / Association des Actuaires Suisses = Bulletin / Association of Swiss Actuaries
<b>Herausgeber:</b>	Vereinigung Schweizerischer Versicherungsmathematiker
<b>Band:</b>	63 (1963)
<b>Artikel:</b>	The variation problem in case of the linear decreasing rate of interest
<b>Autor:</b>	Sumitsuji, Osamu
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-966925">https://doi.org/10.5169/seals-966925</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# The Variation Problem in Case of the Linear Decreasing Rate of Interest

*By Osamu Sumitsuji, Osaka (Japan)*

## Summary

In this paper, the author gives approximation formulas for premiums and mathematical reserves of endowment assurances with linear decreasing rate of interest, in connection with the well-known variation problem. Also he shows the relation between the mathematical reserve of endowment assurance with decreasing rate of interest and that with increasing additive extra mortality.

## Zusammenfassung

In Anlehnung an frühere Arbeiten befasst sich der Autor mit der Herleitung von approximativen Formeln für Prämie und Deckungskapital der gemischten Versicherung im Falle eines linear sinkenden Zinsfusses. Er weist auch auf den Zusammenhang zwischen den Reserven einer solchen Versicherung hin, wenn einerseits von einem sinkenden Zinsfuss, anderseits von einer steigenden additiven Übersterblichkeit ausgegangen wird.

## Résumé

En s'appuyant sur des travaux antérieurs, l'auteur établit des formules approximatives en vue du calcul de la prime et de la réserve mathématique d'une assurance mixte fondée sur un taux d'intérêt linéairement décroissant. Certaines analogies sont également mises en évidence, pour cette forme d'assurance, entre les réserves mathématiques résultant, d'une part, de l'application d'un taux d'intérêt décroissant et, d'autre part, de la prise en considération d'une surmortalité croissante.

## Riassunto

Fondandosi su lavori precedenti, l'autore si preoccupa di dedurre formule approssimative per il calcolo del premio e della riserva matematica dell'assicurazione mista, nel caso di un tasso d'interesse decrescente in modo lineare. Egli accenna anche ai rapporti tra le riserve matematiche di una tale assicurazione, supposto, da una parte, un tasso d'interesse decrescente e, dall'altra, una maggior mortalità crescente.

§ 1. In my earlier paper<sup>1)</sup>, I showed the following relations:

$$a) \quad -\frac{d}{d\delta} P_{xn|} = \frac{\sum_{0}^{n-1} ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{N_x - N_{x+n}}. \quad (1)$$

$$\begin{aligned} b) \quad \frac{d}{d\delta} {}_t V_{xn|} &= \frac{d}{d\delta} P_{xn|} \frac{D_x}{D_{x+t}} \ddot{a}_{xt|} + \frac{\sum_{0}^{t-1} ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{D_{x+t}} \\ &= \frac{d}{d\delta} P_{xn|} \frac{D_x}{D_{x+t}} \ddot{a}_{xt|} \left( 1 - \frac{\ddot{a}_{xn|}}{\ddot{a}_{xt|}} \frac{\sum_{0}^{t-1} ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{\sum_{0}^{n-1} ({}_t V_{xn|} + P_{xn|}) D_{x+t}} \right) \\ &\stackrel{2)}{\sim} \frac{d}{d\delta} P_{xn|} \frac{D_x}{D_{x+t}} \ddot{a}_{xt|} \left( 1 - \frac{1 + \frac{I a_{xt-1|}}{\ddot{a}_{xt|}}}{1 + \frac{I a_{xn-1|}}{\ddot{a}_{xn|}}} \right) \\ &\sim \frac{d}{d\delta} P_{xn|} \frac{D_x}{D_{x+t}} \ddot{a}_{xt|} \left\{ 1 - \frac{t}{n} - \frac{t(i + q_{xn|})}{6} \left( 1 - \frac{t}{n} \right) \right. \\ &\quad \left. - \frac{1}{n} \left( 1 - \frac{t}{n} \right) - \frac{t}{12n} \left( 1 - \frac{t^2}{n^2} \right) (n q_{xn|})^2 - \frac{t}{24n} \left( 1 - \frac{t^3}{n^3} \right) (n q_{xn|})^3 \right\} \quad (2) \end{aligned}$$

$$\sim \frac{d}{d\delta} P_{xn|} \frac{D_x}{D_{x+t}} \ddot{a}_{xt|} \left( 1 - \frac{t}{n} \right) \quad (3)$$

(for the case where<sup>3)</sup>  $n$  and  $x$  are not large).

<sup>1)</sup> Mitteilungen der Vereinigung schweizerischer Versicherungsmathematiker, Bd. 59 and Bd. 60.

<sup>2)</sup>  $\frac{d}{d\delta} \ddot{a}_{xn|} = I a_{xn-1|}$

$$\frac{I a_{xn-1|}}{\ddot{a}_{xn|}} \sim \frac{n-1}{2} \left\{ 1 - \frac{(n+1)(i + q_{xn|})}{6} - \frac{(n q_{xn|})^2}{12} \left( 1 + \frac{n q_{xn|}}{2} \right) \right\},$$

where  $\frac{\sum_{0}^{n-1} l_{x+t}}{l_x} = n \left( 1 - \frac{n-1}{2} q_{xn|} \right).$

<sup>3)</sup> We assume  $10 \lesssim n \lesssim 30$  in this paper. Moreover in this case we take  $n(i + q_{xn|}) \lesssim 1.5$  and  $n q_{xn|} \lesssim 0.5$ .

Or

$$\sim \frac{d}{d\delta} P_{x\bar{n}} t \left( 1 - \frac{t}{n} \right) \quad (3')$$

(for the case where<sup>1)</sup>  $t$  is small, or  $x$  is large).

$$c) - \frac{\sum_0^{n-1} \left( \frac{d}{d\delta} {}_t V_{x\bar{n}} + \frac{d}{d\delta} P_{x\bar{n}} \right) D_{x+t}}{N_x - N_{x+n}} = \frac{1}{2} \left( \frac{d^2}{d\delta^2} P_{x\bar{n}} - \frac{d}{d\delta} P_{x\bar{n}} \right). \quad (4)$$

Using (2) we have for the case where  $n$  and  $x$  are not large

$$\begin{aligned} \frac{1}{2} \frac{d^2}{d\delta^2} P_{x\bar{n}} - \frac{1}{2} \frac{d}{d\delta} P_{x\bar{n}} &\sim - \frac{d}{d\delta} P_{x\bar{n}} \left\{ \frac{n+1}{2} - \left( 1 - \frac{1}{2n} \right) \frac{I a_{x\bar{n}-1}}{\ddot{a}_{x\bar{n}}} \right. \\ &\quad \left. - \frac{1}{6} + \frac{I^2 a_{x\bar{n}-1}}{2n \ddot{a}_{x\bar{n}}} - \frac{n^2 (i + q_{x\bar{n}})}{72} - \frac{2n}{15} \frac{(n q_{x\bar{n}})^2}{12} \left( 1 + \frac{n}{2} q_{x\bar{n}} \right) \right\}, \end{aligned} \quad (5)$$

$$\text{where } \frac{d^2}{d\delta^2} \ddot{a}_{x\bar{n}} = I^2 a_{x\bar{n}-1},$$

or using (3') we get for the case where  $x$  is large

$$\frac{1}{2} \frac{d^2}{d\delta^2} P_{x\bar{n}} - \frac{1}{2} \frac{d}{d\delta} P_{x\bar{n}} \sim - \frac{d}{d\delta} P_{x\bar{n}} \left\{ 1 + \frac{I a_{x\bar{n}-1}}{\ddot{a}_{x\bar{n}}} - \frac{I^2 a_{x\bar{n}-1}}{n \ddot{a}_{x\bar{n}}} \right\}. \quad (6)$$

d) In case of linear increasing extra mortality such as

$$q'_{x+t} = q_{x+t} + c \frac{t}{n} \quad (c \text{ is a constant}) \text{ we have}^2)$$

$$\begin{aligned} - \frac{{}_t V_{x\bar{n}}^{q_{x+t+c} \frac{t}{n}} - {}_t V_{x\bar{n}}}{c} &\sim 0.73 \left( 0.63 - \frac{t}{n} \right) \frac{{}_t V_{x\bar{n}}^{q_{x+t+c}} - {}_t V_{x\bar{n}}}{c} \\ &\sim 0.73 \left( 0.63 - \frac{t}{n} \right) \frac{d}{d\delta} {}_t V_{x\bar{n}}. \end{aligned} \quad (7)$$

<sup>1)</sup> We assume in this case  $n(i + q_{x\bar{n}}) \gtrsim 1.5$  and  $n q_{x\bar{n}} \gtrsim 0.5$ .

<sup>2)</sup> The better approximation is as follows:

$$\begin{aligned} - \frac{{}_t V_{x\bar{n}}^{q_{x+t+c} \frac{t}{n}} - {}_t V_{x\bar{n}}}{c} &\sim 0.46 \left( 1 - \frac{8}{5} \frac{t}{n} + \frac{3 \left( 1 - \frac{t}{2n} \right)}{n} \right) \frac{d}{d\delta} {}_t V_{x\bar{n}} \quad \text{for small } n, \text{ say } n \sim 10. \\ &\sim 0.46 \left( 1 - \frac{8}{5} \frac{t}{n} \right) \frac{d}{d\delta} {}_t V_{x\bar{n}} \quad \text{for not small } n. \end{aligned}$$

§ 2. Now we assume the rates of interest such as

$$\delta + \delta', \quad \delta + \delta' \frac{n-1}{n}, \quad \dots, \quad \delta + \delta' \frac{n-t}{n}, \quad \dots, \quad \delta + \frac{\delta'}{n}, \quad \delta.$$

The following relation is fundamental.

$$\begin{aligned} \frac{P_{xn|}^{\delta+\delta' \frac{n-t}{n}} - P_{xn|}}{\delta'} &= -\frac{\sum_0^{n-1} (n-t) ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{n(N_x - N_{x+n})} \\ &= \frac{d}{d\delta} P_{xn|} + \frac{\sum_0^{n-1} t ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{n(N_x - N_{x+n})}. \end{aligned}$$

To evaluate  $\frac{\sum_0^{n-1} t ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{n(N_x - N_{x+n})}$ , differentiate the equation (1) with respect to  $\delta$ , and we have

$$\begin{aligned} \frac{d^2}{d\delta^2} P_{xn|} + \frac{d}{d\delta} P_{xn|} \frac{\frac{d}{d\delta} \ddot{a}_{xn|}}{\ddot{a}_{xn|}} &= -\frac{\sum_0^{n-1} \left( \frac{d}{d\delta} {}_t V_{xn|} + \frac{d}{d\delta} P_{xn|} \right) D_{x+t}}{N_x - N_{x+n}} \\ &\quad + \frac{\sum_0^{n-1} t ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{N_x - N_{x+n}}. \end{aligned}$$

Carrying (4) to the right hand side of this equation, we have

$$\frac{\sum_0^{n-1} t ({}_t V_{xn|} + P_{xn|}) D_{x+t}}{n(N_x - N_{x+n})} = \frac{1}{2n} \frac{d^2}{d\delta^2} P_{xn|} + \frac{1}{2n} \frac{d}{d\delta} P_{xn|} - \frac{1}{n} \frac{I a_{xn-1|}}{\ddot{a}_{xn|}} \frac{d}{d\delta} P_{xn|}.$$

Therefore we have

$$\frac{P_{xn|}^{\delta+\delta' \frac{n-t}{n}} - P_{xn|}}{\delta'} = \frac{d}{d\delta} P_{xn|} \left( 1 + \frac{1}{2n} - \frac{I a_{xn-1|}}{n \ddot{a}_{xn|}} \right) + \frac{1}{2n} \frac{d^2}{d\delta^2} P_{xn|} \quad (8)$$

$$= D \frac{d}{d\delta} P_{xn|}, \quad (8')$$

$$\text{where } D = 1 + \frac{1}{2n} - \frac{I a_{xn-1|}}{n \ddot{a}_{xn|}} + \frac{1}{2n} \frac{\frac{d^2}{d\delta^2} P_{xn|}}{\frac{d}{d\delta} P_{xn|}}.$$

Consequently  $\frac{P_{x\bar{n}}^{\delta+\delta'\frac{t}{n}} - P_{x\bar{n}}}{\delta'} = (1-D) \frac{d}{d\delta} P_{x\bar{n}}$ . (9)

The approximation to  $D$  is obtained as follows:

- (i) For the case where  $x$  and  $n$  are not large. From (8), using the approximation (5), we have

$$D \sim \frac{1}{2} + \frac{2}{3n} - \frac{Ia_{x\bar{n-1}}}{2n^2 \ddot{a}_{x\bar{n}}} - \frac{I^2 a_{x\bar{n-1}}}{2n^2 \ddot{a}_{x\bar{n}}} + \frac{n(i+q_{x\bar{n}})}{72} + \frac{(nq_{x\bar{n}})^2}{90} \left(1 + \frac{n}{2} q_{x\bar{n}}\right).$$

On the other hand, we may write

$$\begin{aligned} \frac{Ia_{x\bar{n-1}}}{\ddot{a}_{x\bar{n}}} &\sim \frac{n-1}{2} \left\{ 1 - \frac{(n+1)(i+q_{x\bar{n}})}{6} - \frac{(nq_{x\bar{n}})^2}{12} \left(1 + \frac{nq_{x\bar{n}}}{2}\right) \right\}, \\ \frac{I^2 a_{x\bar{n-1}}}{\ddot{a}_{x\bar{n}}} &\sim \frac{(n-1)(2n-1)}{6} \left\{ 1 - \frac{(n-1)(i+q_{x\bar{n}})}{4} - \frac{(nq_{x\bar{n}})^2}{8} \left(1 + \frac{nq_{x\bar{n}}}{2}\right) \right\}. \end{aligned}$$

Using these approximations, we may have

$$D \sim \frac{1}{3} + \frac{2}{3n} + \frac{(n-1)(i+q_{x\bar{n}})}{18} + \frac{(nq_{x\bar{n}})^2}{31} \left(1 + \frac{n}{2} q_{x\bar{n}}\right). \quad (10)$$

From (10) we have, for the case where  $ni \lesssim 1$  and  $nq_{x\bar{n}} \lesssim \frac{1}{3}$ ,  $D \sim 0.4$ .

Examples:

$n = 30$	$x = 30$	
C.S.O. (41) Table, $2\frac{1}{2}\%$		JPM(3) Table, $3\frac{1}{2}\%$
$q_{30\bar{30}} = 0.0063$		$q_{30\bar{30}} = 0.010$
the true value of $D = 0.401$		$D = 0.417$
by (10) $= 0.407$	by (10) $= 0.431$	

- (ii) For the case where  $x$  is large. Carrying (6) into the formula (8), we obtain

$$D \sim 1 - 2 \frac{Ia_{x\bar{n-1}}}{n \ddot{a}_{x\bar{n}}} + \frac{I^2 a_{x\bar{n-1}}}{n^2 \ddot{a}_{x\bar{n}}}.$$

Hence we may have

$$D \sim \frac{1}{3} + \frac{1}{2n} + \frac{(n-1)(i+q_{x\bar{n}})}{12} + \frac{(nq_{x\bar{n}})^2}{24} \left(1 + \frac{n}{2} q_{x\bar{n}}\right). \quad (11)$$

Examples:

$n = 30$	$x = 50$	$n = 20$	$x = 50$
C.S.O. Table, $2\frac{1}{2}\%$	$J^{PM(3)}$ , $3\frac{1}{2}\%$	$J^{PM(3)}$ , $3\frac{1}{2}\%$	
$q_{50 \bar{30}} = 0.0215$	$q_{50 \bar{30}} = 0.029$	$q_{50 \bar{20}} = 0.026$	
the true value of $D = 0.491$		$0.557$	$0.456$
by (11) $= 0.482$		$0.549$	$0.469$
by (10) $= 0.448$		$0.494$	$0.440$

§ 3. As to the premium reserve we have

$$\begin{aligned} \frac{\delta' \rightarrow 0}{\delta'} \frac{{}_t V_{x \bar{n}}^{\delta+\delta' t/n} - {}_t V_{x \bar{n}}}{\delta'} &= \frac{P_{x \bar{n}}^{\delta+\delta' t/n} - P_{x \bar{n}}}{\delta' \rightarrow 0} \frac{D_x}{D_{x+t}} \ddot{a}_{x \bar{t}} \left( 1 - \frac{\ddot{a}_{x \bar{n}}}{\ddot{a}_{x \bar{t}}} \frac{\sum_{0}^{t-1} t({}_t V_{x \bar{n}} + P_{x \bar{n}}) D_{x+t}}{\sum_{0}^{n-1} t({}_t V_{x \bar{n}} + P_{x \bar{n}}) D_{x+t}} \right) \\ &\sim \frac{P_{x \bar{n}}^{\delta+\delta' t/n} - P_{x \bar{n}}}{\delta' \rightarrow 0} \frac{D_x}{D_{x+t}} \ddot{a}_{x \bar{t}} \left( 1 - \frac{\frac{I^2 a_{x \bar{t}-1}}{\ddot{a}_{x \bar{n}}}}{\frac{I^2 a_{x \bar{n}-1}}{\ddot{a}_{x \bar{n}}}} \right), \end{aligned}$$

using for  $\frac{I^2 a_{x \bar{n}-1}}{\ddot{a}_{x \bar{n}}}$  the same approximation as on p. 37 as well as (3) and (9), we may have<sup>1)</sup>

$$\sim (1-D) \left( 1 + \frac{t}{n} \right) \frac{d}{d\delta} {}_t V_{x \bar{n}}, \quad (12)$$

when  $n$  and  $x$  are not large.

<sup>1)</sup> The better approximation is as follows:

$$\begin{aligned} \frac{\delta' \rightarrow 0}{\delta'} \frac{{}_t V_{x \bar{n}}^{\delta+\delta' t/n} - {}_t V_{x \bar{n}}}{\delta'} &\sim (1-D) \left( 1 + \frac{t}{n} \right) \left\{ 1 + \frac{1}{n} - \frac{t(i+q_{x \bar{n}})}{6} \left( 1 - \frac{3}{2 \left( 1 + \frac{n}{t} \right)} \right) \right\} \frac{d}{d\delta} {}_t V_{x \bar{n}} \\ &\sim (1-D) \left( 1 + \frac{t}{n} \right) \left( 1 + \frac{1}{n} \right) \frac{d}{d\delta} {}_t V_{x \bar{n}} \quad \text{for small } n, \text{ say } n \sim 10 \\ &\sim (1-D) \left( 1 + \frac{t}{n} \right) \frac{d}{d\delta} {}_t V_{x \bar{n}} \quad \text{for not small } n. \end{aligned}$$

Consequently, from the fundamental relation of

$$\frac{\underset{\delta' \rightarrow 0}{t} V_{xn}^{\delta+\delta' \frac{n-t}{n}} - t V_{xn}}{\delta'} + \frac{\underset{\delta' \rightarrow 0}{t} V_{xn}^{\delta+\delta' \frac{t}{n}} - t V_{xn}}{\delta'} = \frac{d}{d\delta} t V_{xn}$$

we obtain<sup>1)</sup>, for the case where  $n$  and  $x$  are not large,

$$\frac{\underset{\delta' \rightarrow 0}{t} V_{xn}^{\delta+\delta' \frac{n-t}{n}} - t V_{xn}}{\delta'} \sim D \left( 1 - \frac{1-D}{D} \frac{t}{n} \right) \frac{d}{d\delta} t V_{xn}. \quad (13)$$

Putting  $D \sim 0.4$ , we have<sup>1)</sup> for the case where  $ni \lesssim 1$  and  $nq_{xn} \lesssim \frac{1}{3}$

$$\frac{\underset{\delta' \rightarrow 0}{t} V_{xn}^{\delta+\delta' \frac{n-t}{n}} - t V_{xn}}{\delta'} \sim 0.4 \left( 1 - \frac{3}{2} \frac{t}{n} \right) \frac{d}{d\delta} t V_{xn}. \quad (13')$$

It is worth while to notice that from (7) and (13) we have, when

$$ni \lesssim 1 \quad \text{and} \quad nq_{xn} \lesssim \frac{1}{3},$$

$$\underset{c \rightarrow 0}{-} \frac{\underset{c \rightarrow 0}{t} V_{xn}^{q_x+t+c \frac{t}{n}} - t V_{xn}}{c} \sim \frac{\underset{\delta' \rightarrow 0}{t} V_{xn}^{\delta+\delta' \frac{n-t}{n}} - t V_{xn}}{\delta'}. \quad (14)$$

This interesting relation is also obtained as follows: Clearly we have

$$\underset{c \rightarrow 0}{-} \frac{\underset{c \rightarrow 0}{P}_{xn}^{q_x+t+c \frac{t}{n}} - P_{xn}}{c} - \frac{\underset{\beta \rightarrow 0}{P}_{xn}^{q_x+t \left(1 + \frac{t}{n}\beta\right)} - P_{xn}}{\beta} = v \frac{\underset{n \ddot{a}_{xn}}{Ia_{xn-1}} - \underset{\delta' \rightarrow 0}{P}_{xn}^{\delta+\delta' \frac{t}{n}} - P_{xn}}{\delta'},$$

$$\underset{c \rightarrow 0}{-} \frac{\underset{c \rightarrow 0}{P}_{xn}^{q_x+t+c \frac{n-t}{n}} - P_{xn}}{c} - \frac{\underset{\beta \rightarrow 0}{P}_{xn}^{q_x+t \left(1 + \frac{n-t}{n}\beta\right)} - P_{xn}}{\beta} = v \left( 1 - \frac{\underset{n \ddot{a}_{xn}}{Ia_{xn-1}}}{\underset{n \ddot{a}_{xn}}{Ia_{xn}}} \right) - \frac{\underset{\delta' \rightarrow 0}{P}_{xn}^{\delta+\delta' \frac{n-t}{n}} - P_{xn}}{\delta'}.$$

Therefore

$$\underset{c \rightarrow 0}{-} \frac{\underset{c \rightarrow 0}{t} V_{xn}^{q_x+t+c \frac{t}{n}} - t V_{xn}}{c} - \frac{\underset{\delta' \rightarrow 0}{t} V_{xn}^{\delta+\delta' \frac{n-t}{n}} - t V_{xn}}{\delta'}$$

$$= \frac{d}{d\delta} t V_{xn} + \frac{v}{n} \frac{D_x}{D_{x+t}} \ddot{a}_{xt} \left( \frac{\underset{n \ddot{a}_{xn}}{Ia_{xn-1}}}{\ddot{a}_{xn}} - \frac{\underset{n \ddot{a}_{xn}}{Ia_{xt-1}}}{\ddot{a}_{xt}} \right) + \frac{\underset{\beta \rightarrow 0}{t} V_{xn}^{q_x+t \left(1 + \frac{t}{n}\beta\right)} - t V_{xn}}{\beta}.$$

<sup>1)</sup> When  $n \sim 10$ , the formulas (13) and (13') may be slightly modified.

And when  $ni \lesssim 1$  and  $nq_{xn} \lesssim \frac{1}{3}$  we see easily that

$$\begin{aligned} -\frac{d}{d\delta} {}_t V_{xn} &= \frac{\ddot{a}_{x+t-n}}{\ddot{a}_{xn}} \left( \frac{Ia_{xn-1}}{\ddot{a}_{xn}} - \frac{Ia_{x+t-n-1}}{\ddot{a}_{x+t-n}} \right) \\ &\sim \frac{v}{n} \frac{D_x}{D_{x+t}} \ddot{a}_{xt} \left( \frac{Ia_{xn-1}}{\ddot{a}_{xn}} - \frac{Ia_{xt-1}}{\ddot{a}_{xt}} \right), \end{aligned}$$

and  $\frac{{}_t V_{xn}^{q_{x+t}(1+\frac{t}{n}\beta)}}{\beta \rightarrow 0} - {}_t V_{xn}$  is small. Therefore we get the above relation.

§ 4. The formula (8) is suitable for the case where  $\delta'$  is small. When  $\delta'$  is not small the following modified formula is better.

$$\begin{aligned} \frac{P_{xn}^{i+i'\frac{n-t}{n}} - P_{xn}}{i'} &= v D \frac{d}{d\delta} P_{xn} + \frac{v \sum_{0}^{n-1} (|\Delta {}_t V_{xn}| + |\Delta P_{xn}|) D_{x+t}}{N_x - N_{x+n}} \\ &\quad - \frac{v \sum_{0}^{n-1} t (|\Delta {}_t V_{xn}| + |\Delta P_{xn}|) D_{x+t}}{n (N_x - N_{x+n})}, \end{aligned}$$

$$\text{where } \Delta P_{xn} = P_{xn}^{i+i'\frac{n-t}{n}} - P_{xn}, \quad \Delta {}_t V_{xn} = {}_t V_{xn}^{i+i'\frac{n-t}{n}} - {}_t V_{xn}.$$

Using the following approximations

$$\begin{aligned} |\Delta P_{xn}| &\sim i' v D \left| \frac{d}{d\delta} P_{xn} \right|, \\ |\Delta {}_t V_{xn}| &\sim i' v D \left| \frac{d}{d\delta} {}_t V_{xn} \right| \left( 1 - \frac{1-D}{D} \frac{t}{n} \right), \quad \text{we have} \end{aligned}$$

$$\frac{\sum_{0}^{n-1} (|\Delta {}_t V_{xn}| + |\Delta P_{xn}|) D_{x+t}}{N_x - N_{x+n}} \sim i' v D \left| \frac{d}{d\delta} P_{xn} \right| \left\{ 1 + \frac{n}{12} \left( 3 - \frac{1}{D} \right) \right\} \quad (15)$$

$$\text{and } \frac{\sum_{0}^{n-1} t (|\Delta {}_t V_{xn}| + |\Delta P_{xn}|) D_{x+t}}{n (N_x - N_{x+n})} \sim i' v D \left| \frac{d}{d\delta} P_{xn} \right| \left\{ \frac{1}{2} + \frac{n}{5} \left( \frac{2}{3} - \frac{1}{4D} \right) \right\}. \quad (16)$$

Therefore we may write when  $x$  and  $n$  are not large

$$P_{xn}^{i+i'\frac{n-t}{n}} - P_{xn} \sim i' v D \frac{d}{d\delta} P_{xn} \left\{ 1 - i' v \left[ \frac{1}{2} + \frac{n}{60} \left( 7 - \frac{2}{D} \right) \right] \right\}. \quad (17)$$

When  $t \equiv 0$ , we get  $D = 1$ , (16) = 0, and we may write from (17)

$$P_{xn}^{i+i'} - P_{xn} \sim i' v \frac{d}{d\delta} P_{xn} \left\{ 1 - i' v \left( 1 + \frac{n}{6} \right) \right\} \quad (\text{for the case where } x \text{ and } n \text{ are not large}).$$