

<b>Zeitschrift:</b>	Mitteilungen / Vereinigung Schweizerischer Versicherungsmathematiker = Bulletin / Association des Actuaires Suisses = Bulletin / Association of Swiss Actuaries
<b>Herausgeber:</b>	Vereinigung Schweizerischer Versicherungsmathematiker
<b>Band:</b>	64 (1964)
<b>Artikel:</b>	A theorem in operational calculus
<b>Autor:</b>	Dahiya, R.S.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-966968">https://doi.org/10.5169/seals-966968</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 01.09.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## A Theorem in Operational Calculus

By R. S. Dahiya, Pilani (Rajasthan), India

### Summary

The author proves a theorem in the field of Laplace transformations. Two examples show the possible applications.

#### 1. Introduction. The integral equation

$$\Phi(p) = p \int_0^\infty e^{-pt} f(t) dt, \quad R(p) > 0. \quad (1.1)$$

represents the classical Laplace transform and the functions  $\Phi(p)$  and  $f(t)$  related by (1.1), are said to be operationally related to each other.  $\Phi(p)$  is called the image and  $f(t)$  the original. Symbolically we can write

$$\Phi(p) \doteq f(t) \quad \text{or} \quad f(t) \doteq \Phi(p).$$

The transforms, we have dealt with, are either the Hankel transform or another transform in which the kernel is designated by  $\tilde{\omega}_{\mu,\nu}(x)$ , where

$$\tilde{\omega}_{\mu,\nu}(x) = \sqrt{x} \int_0^\infty J_\nu\left(\frac{x}{t}\right) J_\mu(t) \frac{dt}{t}. \quad (1.2)$$

The integral on the right was first evaluated by C.V.H. Rao [1] and that it plays the role of a transform was conjectured by Watson [2]. Later on Bhatnagar has proved in detail that it plays the role of a transform. In this paper, we make use of Goldstein's theorem [3] as follows:

Let

- (i)  $\Phi_1(p) \doteq f_1(t),$
- (ii)  $\Phi_2(p) \doteq f_2(t),$

then after applying Goldstein's theorem, we get

$$\int_0^\infty f_1(t) \Phi_2(t) \frac{dt}{t} = \int_0^\infty \Phi_1(t) f_2(t) \frac{dt}{t},$$

where  $\Phi(p)$  is the image and  $f(t)$  is the original.

2. Theorem. Let

- (i)  $\psi(p) \doteq f(x),$
- (ii)  $t^{\mu-\lambda-\frac{1}{2}} f(t) \quad \text{be } R_{\mu,\nu},$

then

$$x^{2\mu-\lambda} f(x) \doteq - \frac{2^{2\mu-\lambda+1} \Gamma\left(\mu + \frac{3}{2}\right) \Gamma\left(\frac{\mu+\nu+3}{2}\right)}{\Gamma\left(\frac{\lambda+3}{2}\right) \Gamma\left(\frac{\lambda}{2} + 2\right) \Gamma\left(\frac{\nu-\mu-1}{2}\right)} \\ \cdot p^2 \int_0^\infty t^{\lambda+1} \psi(t) {}_3F_2\left(\begin{matrix} \mu + \frac{3}{2}, \frac{\mu+\nu+3}{2}, \frac{\nu-\mu+3}{2} \\ \frac{\lambda+3}{2}, \frac{\lambda}{2} + 2 \end{matrix}; t^2 p^2\right) dt, \quad (2.1)$$

provided  $f(x)$  is absolutely integrable in  $(0, \infty)$ ,  $t^{\mu-\lambda-\frac{1}{2}} f(t)$  and  $x^{2\mu-\lambda} f(x)$  are integrable in  $(0, \infty)$ , and  $R(\mu) \geqq -\frac{1}{2}$ ,  $R(\nu) \geqq -\frac{1}{2}$ ,  $R(\lambda+2) \leqq 0$ .

Proof: Let

$$p^m \tilde{\omega}_{\mu,\nu}(p) \doteq F(x)$$

then

$$x^\lambda F\left(\frac{1}{x}\right) \doteq p^{\frac{1-\lambda}{2}} \int_0^\infty J_{\lambda+1}(2\sqrt{xp}) x^{\frac{\lambda-1}{2}+m} \tilde{\omega}_{\mu,\nu}(x) dx, \quad (2.2)$$

$R(\lambda+2) > 0$ ,  $R(\lambda+m+\mu) \geqq -\frac{3}{2}$ ,  $R(\lambda+m+\nu) \geqq -\frac{3}{2}$ ,  $R(\lambda+2m) < 0$

$$\text{or, } x^\lambda F\left(\frac{1}{x}\right) \doteq p^{\frac{1-\lambda}{2}} \int_0^\infty J_{\lambda+1}(2\sqrt{pt}) t^{\frac{\lambda-1}{2}+m} \tilde{\omega}_{\mu,\nu}(t) dt, \quad (2.3)$$

$$\text{or, } x^\lambda F\left(\frac{1}{x}\right) \doteq \frac{p^{1-m-\lambda}}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\Gamma(\lambda+m+s)}{\Gamma(-m-s+2)} 2^{1-2s} \frac{\Gamma\left(\frac{\mu-s}{2} + \frac{3}{4}\right) \Gamma\left(\frac{\nu-s}{2} + \frac{3}{4}\right)}{\Gamma\left(\frac{\mu+s}{2} + \frac{1}{4}\right) \Gamma\left(\frac{\nu+s}{2} + \frac{1}{4}\right)} p^{-s} ds, \quad (2.4)$$

$$\text{or, } x^\lambda F\left(\frac{1}{x}\right) \doteq \frac{p^{1-\lambda-m}}{2\pi i} \int_{c-i\infty}^{c+i\infty} 2^{\lambda+2m-1} \frac{\Gamma\left(\frac{\lambda+m+s}{2}\right) \Gamma\left(\frac{\lambda+m+s+1}{2}\right) \Gamma\left(\frac{\mu-s}{2} + \frac{3}{4}\right)}{\Gamma\left(\frac{-m-s+2}{2}\right) \Gamma\left(\frac{-m-s+3}{2}\right) \Gamma\left(\frac{\mu+s}{2} + \frac{1}{4}\right)} \cdot \frac{\Gamma\left(\frac{\nu-s}{2} + \frac{3}{4}\right)}{\Gamma\left(\frac{\nu+s}{2} + \frac{1}{4}\right)} p^{-s} ds. \quad (2.5)$$

By putting  $m = \mu - \lambda + \frac{1}{2}$  and then evaluating the integral, we get

$$x^\lambda F\left(\frac{1}{x}\right) \doteq 2^{2\mu-\lambda+1} p^2 \sum_{n=0}^{\infty} \frac{(-1)^n}{(n)!} \frac{\Gamma\left(\mu+n+\frac{3}{2}\right) \Gamma\left(\frac{\mu+\nu+3}{2}+n\right)}{\Gamma\left(\frac{\lambda}{2}+n+\frac{3}{2}\right) \Gamma\left(\frac{\lambda}{2}+n+2\right) \Gamma\left(\frac{\nu-\mu}{2}-n-\frac{1}{2}\right)} p^{2n} \quad (2.6)$$

$$\text{or, } x^\lambda F\left(\frac{1}{x}\right) \doteq 2^{2\mu-\lambda+1} p^2 \frac{\Gamma\left(\mu+\frac{3}{2}\right) \Gamma\left(\frac{\mu+\nu+3}{2}\right)}{\Gamma\left(\frac{\lambda+3}{2}\right) \Gamma\left(\frac{\lambda}{2}+2\right) \Gamma\left(\frac{\nu-\mu-1}{2}\right)} \cdot {}_3F_2\left(\begin{matrix} \mu+\frac{3}{2}, \frac{\mu+\nu+3}{2}, \frac{\mu-\nu+3}{2}; \\ \frac{\lambda+3}{2}, \frac{\lambda}{2}+2; \end{matrix} p^2\right). \quad (2.7)$$

On writing  $\frac{x}{t}$  for  $x$  we have

$$\left(\frac{x}{t}\right)^{\lambda} F\left(\frac{t}{x}\right) \doteq 2^{2\mu-\lambda+1} \frac{\Gamma\left(\mu+\frac{3}{2}\right) \Gamma\left(\frac{\mu+\nu+3}{2}\right) p^2 t^2}{\Gamma\left(\frac{\lambda+3}{2}\right) \Gamma\left(\frac{\lambda}{2}+2\right) \Gamma\left(\frac{\nu-\mu-1}{2}\right)} \\ \cdot {}_3F_2\left(\begin{matrix} \mu+\frac{3}{2}, \frac{\mu+\nu+3}{2}, \frac{\mu-\nu+3}{2} \\ \frac{\lambda+3}{2}, \frac{\lambda}{2}+2; \end{matrix} t^2 p^2\right). \quad (\text{A})$$

Let

$$(i) \quad p^{\mu-\lambda+\frac{1}{2}} \tilde{\omega}_{\mu,\nu}(p) \doteq F(x), \quad (\text{B})$$

$$(ii) \quad \psi(p) \doteq f(x), \quad (\text{C})$$

$$\text{Also from (B), } (ap)^{\mu-\lambda+\frac{1}{2}} \tilde{\omega}_{\mu,\nu}(ap) \doteq F\left(\frac{x}{a}\right). \quad (\text{D})$$

Applying Goldstein's theorem to (C) and (D), we get

$$\int_0^\infty f(t) (at)^{\mu-\lambda+\frac{1}{2}} \tilde{\omega}_{\mu,\nu}(at) \frac{dt}{t} = \int_0^\infty \psi(t) F\left(\frac{t}{a}\right) \frac{dt}{t}. \quad (2.8)$$

On writing  $x$  for  $a$  and multiplying by  $x^\lambda$  to both sides, it follows

$$x^\lambda \int_0^\infty f(t) (xt)^{\mu-\lambda+\frac{1}{2}} \tilde{\omega}_{\mu,\nu}(xt) \frac{dt}{t} = \int_0^\infty \psi(t) F\left(\frac{t}{x}\right) \left(\frac{x}{t}\right)^\lambda t^{\lambda-1} dt. \quad (2.9)$$

Interpreting with the help of (A), we get

$$x^{\mu+\frac{1}{2}} \int_0^\infty f(t) t^{\mu-\lambda-\frac{1}{2}} \tilde{\omega}_{\mu,\nu}(xt) dt \doteq \frac{2^{2\mu-\lambda+1} \Gamma\left(\mu+\frac{3}{2}\right) \Gamma\left(\frac{\mu+\nu+3}{2}\right)}{\Gamma\left(\frac{\lambda+3}{2}\right) \Gamma\left(\frac{\lambda}{2}+2\right) \Gamma\left(\frac{\nu-\mu-1}{2}\right)} \\ \cdot p^2 \int_0^\infty t^{\lambda+1} \psi(t) {}_3F_2\left(\begin{matrix} \mu+\frac{3}{2}, \frac{\mu+\nu+3}{2}, \frac{\mu-\nu+3}{2} \\ \frac{\lambda+3}{2}, \frac{\lambda}{2}+2; \end{matrix} t^2 p^2\right) dt. \quad (2.10)$$

If  $t^{\mu-\lambda-\frac{1}{2}} f(t)$  is  $R_{\mu,\nu}$ , we obtain the required result.

3. Example 1. Let  $t^{\frac{\mu-\nu+1}{2}} J_{\frac{\mu+\nu}{2}}(t)$  be  $R_{\mu,\nu}$ ,

then  $f(t) = t^{\frac{\lambda-\nu+\mu+1}{2}} J_{\frac{\mu+\nu}{2}}(t)$

$$\stackrel{.}{=} \frac{\Gamma(\lambda+2) p^{-\lambda-1}}{2^{\frac{\mu+\nu}{2}} \Gamma\left(\frac{\mu+\nu+2}{2}\right)} {}_2F_1\left(\begin{array}{c} \frac{\lambda}{2}+1, \frac{\lambda+3}{2}; \\ \frac{\mu+\nu+2}{2}; \end{array} -\frac{1}{p^2}\right) \equiv \psi(p)$$

and  $x^{2\mu-\lambda} f(x) = x^{\frac{3\mu-\nu}{2}+1} J_{\frac{\mu+\nu}{2}}(x)$

$$\stackrel{.}{=} \frac{\Gamma(2\mu+2) p^{-2\mu-1}}{2^{\frac{\mu+\nu}{2}} \Gamma\left(\frac{\mu+\nu+2}{2}\right)} {}_2F_1\left(\begin{array}{c} \mu+1, \mu+\frac{3}{2}; \\ \frac{\mu+\nu+2}{2}; \end{array} -\frac{1}{p^2}\right). \quad (3.1)$$

Hence from (2.1) we get

$$\int_0^\infty {}_2F_1\left(\begin{array}{c} \frac{\lambda}{2}+1, \frac{\lambda+3}{2}; \\ \frac{\mu+\nu+2}{2}; \end{array} -\frac{1}{t^2}\right) {}_3F_2\left(\begin{array}{c} \mu+\frac{3}{2}, \frac{\mu+\nu+3}{2}, \frac{\mu-\nu+3}{2}; \\ \frac{\lambda+3}{2}, \frac{\lambda}{2}+2; \end{array} t^2 p^2\right) dt$$

$$= \frac{\Gamma\left(\frac{\lambda+3}{2}\right) \Gamma\left(\frac{\lambda}{2}+2\right) \Gamma\left(\frac{\nu-\mu-1}{2}\right) \Gamma(2\mu+2) p^{-2\mu-3}}{2^{2\mu-\lambda+1} \Gamma\left(\mu+\frac{3}{2}\right) \Gamma\left(\frac{\mu+\nu+3}{2}\right) \Gamma(\lambda+2)} {}_2F_1\left(\begin{array}{c} \mu+1, \mu+\frac{3}{2}; \\ \frac{\mu+\nu+2}{2}; \end{array} -\frac{1}{p^2}\right).$$

In particular, if  $\mu = \nu$ , then (3.2)

$$\int_0^\infty {}_2F_1\left(\begin{array}{c} \frac{\lambda}{2}+1, \frac{\lambda+3}{2}; \\ \nu+1; \end{array} -\frac{1}{t^2}\right) {}_3F_2\left(\begin{array}{c} \nu+\frac{3}{2}, \nu+\frac{3}{2}, \frac{3}{2}; \\ \frac{\lambda+3}{2}, \frac{\lambda}{2}+2; \end{array} t^2 p^2\right) dt$$

$$= -\frac{\sqrt{\pi} \Gamma\left(\frac{\lambda+3}{2}\right) \Gamma\left(\frac{\lambda}{2}+2\right) \Gamma(2\nu+2) p^{-2\nu-3}}{2^{2\nu-\lambda} \Gamma\left(\nu+\frac{3}{2}\right) \Gamma\left(\frac{2\nu+3}{2}\right) \Gamma(\lambda+2)} {}_1F_0\left(\nu+\frac{3}{2}; -\frac{1}{p^2}\right).$$

4. Example 2. Let

$$t^{n+\mu+\frac{1}{2}} K_n(t) \quad \text{be} \quad R_{\mu,\nu}, \quad \text{where} \quad \nu = \mu + 2n$$

then  $f(t) = t^{n+\lambda+1} K_n(t)$

$$\doteqdot \frac{\sin [(n+\lambda+1)\pi] \Gamma(\lambda+2)}{\sin [(2n+\lambda+1)\pi] (p^2-1)^{\frac{n+\lambda+2}{2}}} Q_{n+\lambda+1}^n \left( \frac{p}{\sqrt{p^2-1}} \right) \equiv \varphi(p)$$

and

$$\begin{aligned} x^{2\mu-\lambda} f(x) &= x^{2\mu+n+1} K_n(x) \\ &\doteqdot \frac{\sin [(2\mu+n+1)\pi] \Gamma(2\mu+2)}{\sin [(2\mu+2n+1)\pi] (p^2-1)^{\frac{2\mu+n+2}{2}}} Q_{2\mu+n+1}^n \left( \frac{p}{\sqrt{p^2-1}} \right). \end{aligned}$$

Hence from (2.1) we get

$$\begin{aligned} &\int_0^\infty \frac{t^{\lambda+1}}{(t^2-1)^{\frac{2\mu+n+2}{2}}} Q_{2\mu+n+1}^n \left( \frac{t}{\sqrt{t^2-1}} \right) {}_3F_2 \left( \begin{matrix} \mu + \frac{3}{2}, \frac{2\mu+2n+3}{2}, \frac{3-2n}{2} \\ \frac{\lambda+3}{2}, \frac{\lambda}{2} + 2; \end{matrix} t^2 p^2 \right) dt \\ &= \frac{\Gamma(2\mu+2) \Gamma\left(\frac{\lambda+3}{2}\right) \Gamma\left(\frac{\lambda}{2}+2\right) \Gamma\left(\frac{2n-1}{2}\right) \sin [(2\mu+n+1)\pi] \sin [(2n+\lambda+1)\pi]}{2^{2\mu-\lambda+1} \Gamma\left(\mu + \frac{3}{2}\right) \Gamma\left(\frac{2\mu+2n+3}{2}\right) \Gamma(\lambda+2) \sin [(2\mu+n+1)\pi] \sin [(n+\lambda+1)\pi]} \\ &\quad \cdot \frac{1}{p^2 (p^2-1)^{\frac{2\mu+n+2}{2}}} Q_{2\mu+n+1}^n \left( \frac{p}{\sqrt{p^2-1}} \right), \quad R(2\mu+2n-\lambda) > 0. \quad (4.1) \end{aligned}$$

I am thankful to Prof. S. C. Mitra and Dr. B. Singh for their kind help and encouragement in the preparation of this paper.

## References

- [1] Rao, C. V. H.: Messenger of Mathematics, Vol. 47 (1918), 134–137.
- [2] Watson, G. N.: Quart. Jour. of Math., Vol. 2 (1931), 298–309.
- [3] Goldstein: Proc. Lond. Math. Soc., Vol. 2 (1932), 34, P (103).
- [4] Erdélyi, A., and others: Tables of Integral Transforms, *i* and *ii*, Bateman Project.

## Zusammenfassung

Der Autor beweist einen Satz aus dem Gebiet der Laplace-Transformationen und illustriert ihn an zwei Beispielen.

## Résumé

Dans la présente note, l'auteur établit un théorème appartenant à la transformation de Laplace et en donne deux applications.

## Riassunto

L'autore prova un teorema riguardante la trasformazione di Laplace e ne da due esempi.