

Zeitschrift: Mitteilungen / Schweizerische Vereinigung der
Versicherungsmathematiker = Bulletin / Association Suisse des
Actuaires = Bulletin / Swiss Association of Actuaries

Herausgeber: Schweizerische Vereinigung der Versicherungsmathematiker

Band: - (1989)

Heft: 2

Artikel: From the convolution of uniform distributions to the probability of ruin

Autor: Gerber, Hans U.

DOI: <https://doi.org/10.5169/seals-967222>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

HANS U. GERBER, Lausanne

From the Convolution of Uniform Distributions to the Probability of Ruin

1. Introduction

Let y_1, y_2, \dots, y_n be positive numbers. We consider the uniform distribution over $(0, y_i)$; the corresponding cumulative distribution function is

$$F_i(x) = \begin{cases} 0 & \text{for } x < 0 \\ \frac{x}{y_i} & \text{for } 0 \leq x < y_i \\ 1 & \text{for } x \geq y_i \end{cases} \quad (1)$$

$i = 1, 2, \dots, n$. The cumulative distribution function of their convolution is

$$H_n(x) = F_1 * F_2 * \dots * F_n(x). \quad (2)$$

Explicit formulas for $H_n(x) = H_n(x; y_1, \dots, y_n)$ and its derivative are well known and have been derived in different ways, see for example *Seal* (1950) and *Shiu* (1987). In this note, which is mostly of a pedagogical nature, we shall first show by geometric reasoning how two dual expressions for H_n can be obtained. In section 4 the extension to the convolution of distributions with decreasing probability density functions is discussed. Thanks to these more general formulas certain series expressions for the probability of ruin can be derived easily.

2. Geometric reasoning: the rocket principle

We start with $n = 2$ and prefer to consider $y_1 y_2 H_2(x)$ (instead of $H_2(x)$), since this expression has a geometric interpretation: It is the area of that part of the rectangle

$$\{(x_1, x_2) \mid 0 \leq x_1 \leq y_1, \quad 0 \leq x_2 \leq y_2\}$$

Figure 1a Geometric derivation of formula (3 a)





Figure 1b Geometric derivation of formula (3 b)



that is below the line $x_1 + x_2 = x$. There are two formulas for $y_1 y_2 H_2(x)$:

$$\begin{aligned}
 y_1 y_2 H_2(x) &= y_1 y_2 - \frac{1}{2}(y_1 + y_2 - x)_+^2 \\
 &\quad + \frac{1}{2}(y_1 - x)_+^2 + \frac{1}{2}(y_2 - x)_+^2 \\
 &\quad - \frac{1}{2}(-x)_+^2,
 \end{aligned} \tag{3a}$$

and

$$y_1 y_2 H_2(x) = \frac{1}{2}(x)_+^2 - \frac{1}{2}(x - y_1)_+^2 - \frac{1}{2}(x - y_2)_+^2 + \frac{1}{2}(x - y_1 - y_2)_+^2. \tag{3b}$$

These inclusion-exclusion type formulas are best derived geometrically. The geometric derivation of (3a), term by term, is illustrated in Figure 1a. The idea is to start with a line above the rectangle (i.e., $x > y_1 + y_2$), to lower the line successively and thereby to make the necessary corrections by adding and subtracting the areas of certain isosceles right triangles. In Figure 1b we illustrate the geometric derivation of (3b). Here, the idea is to start with a line that is below the rectangle and to raise it successively.

For $n = 3$ we consider $y_1 y_2 y_3 H_3(x)$, which is the volume of that part of the three dimensional rectangle

$$\{(x_1, x_2, x_3) \mid 0 \leq x_i \leq y_i, \quad i = 1, 2, 3\}$$

that is below the plane $x_1 + x_2 + x_3 = x$. If we start with a plane that is above the three-dimensional rectangle ($x > y_1 + y_2 + y_3$) and lower the plane successively, we have to add and subtract the volumes of certain isosceles right pyramids. This way we get

$$\begin{aligned} y_1 y_2 y_3 H_3(x) = & y_1 y_2 y_3 \\ & - \frac{1}{6}(y_1 + y_2 + y_3 - x)_+^3 \\ & + \frac{1}{6}(y_1 + y_2 - x)_+^3 + \frac{1}{6}(y_1 + y_3 - x)_+^3 + \frac{1}{6}(y_2 + y_3 - x)_+^3 \\ & - \frac{1}{6}(y_1 - x)_+^3 - \frac{1}{6}(y_2 - x)_+^3 - \frac{1}{6}(y_3 - x)_+^3 \\ & + \frac{1}{6}(-x)_+^3. \end{aligned} \quad (4a)$$

If we start with a plane that is below the three-dimensional rectangle ($x < 0$) and raise it successively, we get

$$\begin{aligned} y_1 y_2 y_3 H_3(x) = & \frac{1}{6}(x)_+^3 \\ & - \frac{1}{6}(x - y_3)_+^3 - \frac{1}{6}(x - y_2)_+^3 - \frac{1}{6}(x - y_1)_+^3 \\ & + \frac{1}{6}(x - y_2 - y_3)_+^3 + \frac{1}{6}(x - y_1 - y_3)_+^3 + \frac{1}{6}(x - y_1 - y_2)_+^3 \\ & - \frac{1}{6}(x - y_1 - y_2 - y_3)_+^3. \end{aligned} \quad (4b)$$

The artistic reader is invited to draw diagrams similar to those of Figure 1!

3. The general formulas

For arbitrary n , the formulas are now the following natural extensions:

$$\begin{aligned}
 \left(\prod_{i=1}^n y_i \right) H_n(x) &= \prod_{i=1}^n y_i \\
 &\quad - \frac{1}{n!} (y_1 + \cdots + y_n - x)_+^n \\
 &\quad + \frac{1}{n!} \sum (y_{i_1} + \cdots + y_{i_{n-1}} - x)_+^n \\
 &\quad - \frac{1}{n!} \sum (y_{i_1} + \cdots + y_{i_{n-2}} - x)_+^n \\
 &\quad \dots \\
 &\quad \pm \frac{1}{n!} (-x)_+^n
 \end{aligned} \tag{5a}$$

and

$$\begin{aligned}
 \left(\prod_{i=1}^n y_i \right) H_n(x) &= \frac{1}{n!} (x)_+^n \\
 &\quad - \frac{1}{n!} \sum_{i=1}^n (x - y_i)_+^n \\
 &\quad + \frac{1}{n!} \sum (x - y_{i_1} - y_{i_2})_+^n \\
 &\quad \dots \\
 &\quad \pm \frac{1}{n!} (x - y_1 - \cdots - y_n)_+^n.
 \end{aligned} \tag{5b}$$

These formulas can be verified by induction with respect to n . For (5b) this can be done, for example, by using the recursive relation

$$\left(\prod_{i=1}^n y_i \right) H_n(x) = \prod_{i=1}^{n-1} y_i \int_0^{y_n} H_{n-1}(x - y) dy \tag{6}$$

and observing that

$$\frac{1}{(n-1)!} \int_0^{y_n} (x - a - y_{n-1})_+^{n-1} dy = \frac{1}{n!} (x - a - y_n)_+^n - \frac{1}{n!} (x - a)_+^n \tag{7}$$

for any $a \geq 0$.

4. Application: The convolution of distributions with decreasing probability density functions

Let Y_1, Y_2, \dots, Y_n be independent, positive random variables. We denote the cumulative distribution function of Y_i by $P_i(x)$ and suppose that its mean, μ_i is finite. To $P_i(x)$ we associate another cumulative distribution function, given by the formula

$$Q_i(x) = \frac{1}{\mu_i} \int_0^x [1 - P_i(z)] dz. \quad (8)$$

Note that there is a one-to-one correspondence between Q_i and P_i : as $Q_i(x)$ is a distribution function with decreasing probability density function, the underlying P_i can be retrieved by the formula

$$P_i(x) = 1 - \frac{Q_i'(x)}{Q_i'(0)}. \quad (9)$$

Furthermore, the Laplace transform of Q_i can be expressed in terms of the Laplace transform of P_i :

$$\int_0^{\infty} e^{-tx} Q_i(dx) = \frac{1 - \int_0^{\infty} e^{-tx} P_i(dx)}{t\mu_i}. \quad (10)$$

The proof of this well known formula is a simple exercise of integration by parts.

Let us now consider random variables X_1, \dots, X_n which (given Y_1, \dots, Y_n) are conditionally independent, such that X_i is uniformly distributed over $(0, Y_i)$; then $H_n(x; Y_1, \dots, Y_n)$ is the conditional cumulative distribution function of $X_1 + \dots + X_n$. The following Lemma will allow us to apply the results of section 3.

Lemma:

$$E \left[\left(\prod_{i=1}^n Y_i \right) H_n(x; Y_1, \dots, Y_n) \right] = \left(\prod_{i=1}^n \mu_i \right) Q_1 * Q_2 * \dots * Q_n(x).$$

Proof: We verify this formula by calculating the Laplace transforms. Since

$$\int_0^{\infty} e^{-tx} H_n(dx; y_1, \dots, y_n) = \prod_{i=1}^n \left(\frac{1 - e^{-ty_i}}{ty_i} \right), \quad (11)$$

we get

$$\begin{aligned} & \int_0^{\infty} e^{-tx} E \left[\left(\prod_{i=1}^n Y_i \right) H_n(dx; Y_1, \dots, Y_n) \right] \\ &= E \left[\int_0^{\infty} e^{-tx} \left(\prod_{i=1}^n Y_i \right) H_n(dx; Y_1, \dots, Y_n) \right] \\ &= E \left[\prod_{i=1}^n \left(\frac{1 - e^{-tY_i}}{t} \right) \right] \\ &= \prod_{i=1}^n \left(\frac{1 - E[e^{-tY_i}]}{t} \right) \\ &= \left(\prod_{i=1}^n \mu_i \right) \prod_{i=1}^n \left(\frac{1 - E[e^{-tY_i}]}{\mu_i t} \right). \end{aligned} \quad (12)$$

Q.E.D.

We consider now the special case where the Y_i 's are identically distributed. We write P , μ , and Q instead of P_i , μ_i , and Q_i , and set

$$S_k = Y_1 + \dots + Y_k \quad (13)$$

for $k = 1, \dots, n$ ($S_0 = 0$). Let us look at the expression

$$\begin{aligned} & \left(\prod_{i=1}^n Y_i \right) Pr(X_1 + \dots + X_n \leq x \mid Y_1, \dots, Y_n) \\ &= \left(\prod_{i=1}^n Y_i \right) H_n(x; Y_1, \dots, Y_n). \end{aligned} \quad (14)$$

Now we substitute (5a) or (5b), with y_i replaced by Y_i . Finally we take expectations and use the Lemma to obtain

$$\mu^n Q^{*n}(x) = \mu^n - \frac{1}{n!} \sum_{j=0}^n (-1)^j \binom{n}{j} E \left[(S_{n-j} - x)_+^n \right] \quad (15a)$$

and

$$\mu^n Q^{*n}(x) = \frac{1}{n!} \sum_{j=0}^n (-1)^j \binom{n}{j} E \left[(x - S_j)_+^n \right]. \quad (15b)$$

Formula (15a) can also be found in *Gerber* (1988) where it has been derived by probabilistic methods.

5. Application: The probability of ruin

We consider the classical model of the collective theory of risk, where

$$U(t) = u + ct - S(t), \quad t \geq 0, \quad (16)$$

is the surplus of an insurance company at time t . Here u is the initial surplus, c the rate at which the premiums are received, and

$$S(t) = Y_1 + \cdots + Y_{N(t)} \quad (17)$$

the aggregate claims, a compound Poisson process; the claim amounts Y_1, Y_2, \dots are i.i.d. random variables with common distribution function $P(x)$, mean μ , and $\{N(t)\}$ is a Poisson process with parameter λ . It is assumed $c > \lambda\mu$.

The probability of ruin, $\psi(u)$, can be expressed by the convolution formula,

$$\psi(u) = (1 - a\mu) \sum_{n=1}^{\infty} (a\mu)^n [1 - Q^{*n}(u)], \quad (18)$$

where $a = \lambda/c$ and

$$Q(x) = \frac{1}{\mu} \int_0^x [1 - P(t)] dt \quad (19)$$

in agreement with the notation of section 4.

Thanks to formulas (15) we can now derive two alternative formulas for the probability of ruin. First we replace $1 - Q^{*n}(u)$ in (18) by using (15a). After simplification of the resulting double summation we get

$$\psi(u) = (1 - a\mu) \sum_{k=1}^{\infty} \frac{a^k}{k!} E \left[(S_k - u)_+^k e^{-a(S_k - u)} \right]. \quad (20a)$$

Alternatively we may substitute (15b) into (18). After simplification we obtain the formula

$$1 - \psi(u) = (1 - a\mu) \sum_{k=0}^{\infty} \frac{(-a)^k}{k!} E \left[(u - S_k)_+^k e^{a(u-S_k)} \right]. \quad (20b)$$

In the special case where all claims are of size μ , $S_k = k\mu$, this formula is well known, sometimes by the name of Khintchine-Pollaczek, see *Feller* (1966, formula (2.11) of XIV.3). The general expression (20 b) has been derived by *Shiu* (1988). Formula (20 a) has been derived by *Prabhu* (1965, formula (5.55)). If we add (20 a) and (20 b) we obtain the identity

$$1 = (1 - a\mu) \sum_{k=0}^{\infty} \frac{a^k}{k!} E \left[(S_k - u)^k e^{-a(S_k - u)} \right], \quad (21)$$

which is the starting point of *Gerber* (1988).

H. U. Gerber
Ecole des H.E.C
Université de Lausanne
1015 Lausanne

References

- Feller, W.* (1966): An Introduction to Probability Theory and its application, vol. II. Wiley, New York.
- Gerber, H. U.* (1988): Mathematical fun with ruin theory. *Insurance: Mathematics & Economics* 7, 15–23.
- Prabhu, N. U.* (1965): Queues and Inventories: A Study of Their Basic Stochastic Processes. Wiley, New York.
- Seal, H. L.* (1951): Spot the prior reference. *J. Inst. Act. Students' Society* 10, 255–258.
- Shiu, E. S. W.* (1987): Convolution of Uniform Distributions and Ruin Probability. *Scandinavian Actuarial Journal*, 191–197.
- Shiu, E. S. W.* (1988): Calculation of the probability of eventual ruin by Beekman's convolution series. *Insurance: Mathematics & Economics* 7, 41–47.

Summary

The dual formulas for the convolution of uniform distributions can be obtained by a geometric reasoning of the inclusion-exclusion type. The two formulas can be generalized to the convolution of distributions with decreasing probability density functions. Finally, this result can be used to get two series expressions for the probability of ruin.

Zusammenfassung

Zwei klassische Formeln für die Faltung von Gleichverteilungen können leicht auf geometrische Weise (gemäß dem "Raketenprinzip", d.h. mit sukzessiven Korrekturen) hergeleitet werden. Die beiden Formeln werden verallgemeinert für den Fall der Faltung von Verteilungen mit abnehmender Dichtefunktion. Dieses Resultat kann man wiederum benutzen, um zwei duale Reihenentwicklungen der Ruinwahrscheinlichkeit herzuleiten.

Résumé

On peut obtenir des formules duales pour la convolution de distributions uniformes en utilisant une approche géométrique de type inclusion-exclusion. On généralise les deux formules au cas de convolution de distributions dont la fonction de densité de probabilité est une fonction décroissante. Finalement, on utilise ce dernier résultat pour développer des expressions duales (sous forme de séries) pour la probabilité de ruine.