Zeitschrift: Mitteilungen / Schweizerische Vereinigung der

Versicherungsmathematiker = Bulletin / Association Suisse des

Actuaires = Bulletin / Swiss Association of Actuaries

Herausgeber: Schweizerische Vereinigung der Versicherungsmathematiker

Band: - (1994)

Heft: 2

Artikel: Dependent contracts in Bühlmann's credibility model

Autor: De Vijlder, F. / Cossette, H.

DOI: https://doi.org/10.5169/seals-967202

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

B. Wissenschaftliche Mitteilungen

F. DE VIJLDER, H. COSSETTE*, Louvain-la-Neuve

Dependent contracts in Bühlmann's credibility model

1. Introduction: credibility on a roulette, dependent contracts

We consider a roulette with holes numbered 1, 2, ..., k. We denote by θ_j (j = 1, ..., k) the probability that the ball falls in the hole j, in any play. In a perfect roulette,

$$\theta_1 = \theta_2 = \dots = \theta_k = \frac{1}{k}$$
.

We do not assume the roulette to be perfect here. We have

$$\theta_1 + \theta_2 + \dots + \theta_k = 1.$$

Let us observe the roulette during t plays, numbered $1, 2, \ldots, t$. We define X_{js} $(j=1,\ldots,k;\ s=1,\ldots,t)$ to be equal to 1 if the ball falls in the hole j at the play s, and to be equal to 0 if not. We want to find credibility estimators for θ_1,\ldots,θ_k . Bühlmann's model does not apply directly, because it treats portfolio's with independent contracts. The independence assumption is not satisfied on the roulette, because

$$X_{1s} + X_{2s} + \dots + X_{ks} = 1$$
 $(s = 1, \dots, t).$

But Bühlmann's estimator is constructed on a fixed contract, and the portfolio with several contracts is used only for the estimation of the involved parameters. Starting with this idea, we were convinced that only a small adaptation of Bühlmann's model would solve our roulette problem.

We like to believe that the contracts of insurance portfolios can be considered as being independent, mostly because this is very convenient for the construction of a theory. But chain car crashes are not exceptional, and on misty days all cars of a region have higher probabilities to be involved in an accident. During dry hot summers, all wooden cottages are more exposed to fire. In such situations, the stochastic dependence can hardly be neglected. Next credibility model shows that it is not so difficult to take it into account.

^{*}This research is supported by l'Université Laval and Fonds pour la Formation de Chercheurs et l'Aide à la Recherche.

The considerations of this note also apply, mutatis mutandis, to the Bühlmann-Straub model with weighted observations.

2. Bühlmann columns

A Bühlmann column with characteristics $\mu(.), m, a, s^2$ is a column

$$(\Theta, X_1, \dots, X_t)' \tag{1}$$

of random variables such that

$$E(X_s/\Theta) = \mu(\Theta)$$
 $(s = 1, ..., t),$
 $E\mu(\Theta) = m,$
 $Var \mu(\Theta) = a,$ (2)

$$E\operatorname{Cov}(X_r, X_s/\Theta) = \delta_{rs}s^2 \qquad (r, s = 1, \dots, t). \tag{3}$$

(No confusion is possible between the time subscript s and the s occurring in the parameter s^2). Using the indicated covariances, we implicitly assume that

$$EX_s^2 < \infty$$
 $(s = 1, \dots, t).$

Any column (1), such that for fixed Θ , the variables X_1, \ldots, X_t are conditionally i.i.d. is a Bühlmann column with characteristics defined by (2) and (3).

In the paper Bühlmann (1967), at the origin of all the developments of modern credibility theory, the author argued on a single column (1), and he only used the relations (2) and (3). He derived the *credibility estimator* $\hat{\mu}$ for $\mu(\Theta)$:

$$\widehat{\mu} = z \frac{1}{t} (X_1 + \dots + X_t) + (1 - z)m, \quad \text{where} \quad z = \frac{at}{s^2 + at}.$$

The classical Bühlmann model, with several contracts, can be defined as being a finite number of independent Bühlmann columns with the same characteristics. The parameters m, a, s^2 can be estimated from observations in this model.

As we shall see in next sections, the independence assumption can be dropped if some bias is allowed in the estimation of a. In fact, already in the classical Bühlmann and Bühlmann-Straub models, the usual estimators for a are biased. Indeed, pseudo-estimators are biased because they do not use the exact z's. If the classical estimator is used, say \widehat{a} , it may take negative values and it is replaced by $\max(\widehat{a}, 0)$, introducing a bias again.

3. Bühlmann's generalized model

We consider the model with random variables

$$\Theta_1 \quad \dots \quad \Theta_i \quad \dots \quad \Theta_k \\
X_{11} \quad \dots \quad X_{i1} \quad \dots \quad X_{k1} \\
\dots \quad \dots \quad \dots \quad \dots \\
X_{1s} \quad \dots \quad X_{is} \quad \dots \quad X_{ks} \\
\dots \quad \dots \quad \dots \quad \dots \\
X_{1t} \quad \dots \quad X_{it} \quad \dots \quad X_{kt}$$

We only assume that the columns of this tableau are Bühlmann columns with the same characteristics $\mu(.)$, m, a, s^2 :

$$E(X_{is}/\Theta_i) = \mu(\Theta_i),$$

$$E\mu(\Theta_i) = m,$$

$$\operatorname{Var} \mu(\Theta_i) = a \qquad (i = 1, \dots, k; \ s = 1, \dots, t)$$

$$E\operatorname{Cov}(X_{ir}, X_{is}/\Theta_i) = \delta_{rs}s^2 \qquad (i = 1, \dots, k; \ r, s = 1, \dots, t).$$

The credibility estimator for $\mu(\Theta_i)$ is

$$\widehat{\mu}_i = z X_{iM} + (1-z)m,$$
 where $z = \frac{at}{s^2 + at}$ and $X_{iM} = \frac{1}{t} \big(X_{i1} + \dots + X_{it} \big).$

Our problem is to estimate m, s^2, a .

Whereas the columns of the tableau are independent in the classical model, we later (section 6) make a conditional independence assumption on the lines of the tableau.

4. Parameter estimation

Estimation of m.

The obvious estimator of m is

$$\widehat{m} = \frac{1}{kt} \sum_{is} X_{is} .$$

But m might also be estimated in a larger portfolio than the one under consideration. Hereafter, m' denotes any estimate (a constant, not a random variable),

of m. In the roulette problem, m is known exactly, and then we take of course m' = m.

Estimation of s^2 .

As in the classical Bühlmann model, the right estimator of s^2 is

$$\hat{s}^2 = \frac{1}{k(t-1)} \sum_{i} \sum_{s} (X_{is} - X_{iM})^2 \tag{4}$$

It is unbiased. Indeed, the unbiasedness of the estimator

$$\widehat{s}_{i}^{2} = \frac{1}{t-1} \sum_{s} (X_{is} - X_{iM})^{2},$$

based on the observations of the *i*-th contract only, results from the fact that the *i*-th column of the tableau of section 3 is a Bühlmann column. Moreover,

$$\widehat{s}^2 = \frac{1}{k} \sum_i \widehat{s}_i^2 \,.$$

Estimation of a.

We suggest the estimator

$$\widehat{a} = \frac{1}{kt(t-1)} \sum_{i} \sum_{r \neq s} (X_{ir} - m') (X_{is} - m').$$
 (5)

It satisfies

$$E\widehat{a} = a + (m - m')^2, \tag{6}$$

i.e. it has the bias $(m - m')^2$.

Indeed, we have

$$Cov(X_{ir}, X_{is}) = a + \delta_{rs}s^2$$
 $(i = 1, ..., k; r, s = 1, ..., t)$

because the *i*-th contract is a Bühlmann column. Also,

$$(X_{ir} - m')(X_{is} - m')$$

$$= ((X_{ir} - m) + (m - m'))((X_{is} - m) + (m - m'))$$

$$= (X_{ir} - m)(X_{is} - m) + (m - m')(X_{is} - m)$$

$$+ (X_{ir} - m)(m - m') + (m - m')^{2}.$$

Hence, if $r \neq s$,

$$E((X_{ir} - m')(X_{is} - m'))$$

$$= E((X_{ir} - m)(X_{is} - m)) + (m - m')^{2}$$

$$= Cov(X_{ir}, X_{is}) + (m - m')^{2} = a + (m - m')^{2}.$$

This implies (6) because the double sum in the last member of (5) has kt(t-1) terms with the same expectation

$$a + (m - m')^2.$$

We notice that

$$\widehat{a} = \frac{1}{k} \sum_{i} \widehat{a}_{i} ,$$

where

$$\widehat{a}_i = \frac{1}{t(t-1)} \sum_{r \neq s} (X_{ir} - m') (X_{is} - m').$$

Because $a \ge 0$, we cannot accept negative estimates and we have to replace \widehat{a} by $\max(0, \widehat{a})$, or, perhaps even better, by

$$\frac{1}{k} \sum_{i} \max(0, \widehat{a}_i).$$

5. Application to the roulette problem

In the roulette problem of the introduction, we interpret the unknown vector $(\theta_1, \dots, \theta_k)$ as realization of some random vector $(\Theta_1, \dots, \Theta_k)$. To other realizations of that random vector correspond other hypothetical roulettes.

At each play, the ball falls in exactly one hole. Hence

$$X_{1s} + \dots + X_{ks} = 1.$$

Taking expectations, we have km = 1. Hence m = 1/k and we can use the estimator (5) with m' = 1/k. Then (6) shows its unbiasedness.

We use the notations

$$X_{i\Sigma} = X_{i1} + \dots + X_{it} \qquad (i = 1, \dots, k),$$

$$X_{\Sigma\Sigma} = X_{1\Sigma} + \dots + X_{k\Sigma} = t,$$

$$Q = X_{1\Sigma}^2 + \dots + X_{k\Sigma}^2.$$

We have

$$t < Q < t^2$$
.

The case Q=t occurs if the ball falls in different holes in the plays $s=1,\ldots,t$. This is only possible if $t \leq k$. The case $Q=t^2$ occurs if the ball falls in the same hole in each of the plays $s=1,\ldots,t$.

Here we have

$$\mu(\Theta_i) = E(X_{is}/\Theta_i)$$

$$= P(X_{is} = 1/\Theta_i)$$

$$= \Theta_i$$

and $\widehat{\mu}_i$ is the credibility estimator $\widehat{\theta}_i$ of θ_i :

$$\hat{\theta}_i = zX_{iM} + (1-z)\frac{1}{k}$$
 $(i=1,\ldots,k).$ (7)

Estimation of s^2 .

From (4) results

$$k(t-1)\hat{s}^2 = \sum_{i} \sum_{s} \left(X_{is} - \frac{1}{t} X_{i\Sigma} \right)^2$$

$$= \sum_{i} \sum_{s} \left(X_{is} - \frac{2}{t} X_{is} X_{i\Sigma} + \frac{1}{t^2} X_{i\Sigma}^2 \right)$$

$$= X_{\Sigma\Sigma} - \frac{2}{t} \sum_{i} X_{i\Sigma}^2 + \frac{1}{t^2} t \sum_{i} X_i^2$$

$$= t - \frac{Q}{t}.$$

Hence

$$\hat{s}^2 = \frac{t^2 - Q}{kt(t-1)} \,, \tag{8}$$

$$0 \le \hat{s}^2 \le \frac{1}{k} \,. \tag{9}$$

Estimation of a.

$$\sum_{i} \sum_{r \neq s} (X_{ir} - m) (X_{is} - m)$$

$$= \sum_{irs} (X_{ir} - m) (X_{is} - m) - \sum_{is} (X_{is} - m)^{2}$$

$$= \sum_{irs} (X_{ir} X_{is} - m X_{is} - m X_{ir} + m^{2})$$

$$- \sum_{is} (X_{is} - 2m X_{is} + m^{2})$$

$$= Q - mt^{2} - mt^{2} + m^{2}kt^{2} - t + 2mt - ktm^{2}$$

$$= Q - \frac{1}{k}t^{2} - \frac{1}{k}t^{2} + \frac{1}{k^{2}}kt^{2} - t + \frac{2}{k}t - kt\frac{1}{k^{2}}$$

$$= (Q - t) + \frac{t}{k} - \frac{t^{2}}{k} = t(t - 1) - \frac{1}{k}t(t - 1) - (t^{2} - Q).$$

Hence, by (5) and (8),

$$\widehat{a} = \left(\frac{1}{k} - \widehat{s}^2\right) - \frac{1}{k^2} \tag{10}$$

This is a relation between the unbiased estimators \hat{a} , \hat{s}^2 . Taking expectations, we obtain that the corresponding relation holds between the exact values a, s^2 . In fact, the latter relation is obvious because

$$a + s^2 = \text{Var } X_{is} = E(X_{is}^2) - E^2(X_{is}) = E(X_{is}) - m^2$$

= $m - m^2 = \frac{1}{k} - \frac{1}{k^2}$.

Discussion of the solution.

Case 1: The last member of (10) is negative.

Then we can only adopt the estimate a=0. Then z=0 and from (7) results

$$\widehat{\theta}_i = \frac{1}{k}$$
 $(i = 1, \dots, k).$

This means that we cannot contradict the perfectness of the roulette. Case 1 certainly occurs if the balls fall in different holes in the considered plays. Indeed, then Q = t, $\hat{s}^2 = 1/k$ and the last member of (10) equals $-1/k^2$.

Case 2: The last member of (10) is positive.

Then we adopt (8) and (10) as estimators for s^2 , a and the estimate of θ_i results from (7). Of course, all involved random variables X_{is} are supposed to be replaced by their realizations x_{is} .

We tried different values of k, t, Q. In each case there was complete agreement with the intuition that we have of the roulette problem, or at least, there was no contradiction with this intuition. As extreme example, let us assume that in all the considered plays the ball falls in the hole 1. Then

$$Q = t^2, \qquad \widehat{s}^2 = 0, \qquad z = 1,$$

$$\widehat{\theta}_1 = 1, \qquad \widehat{\theta}_i = 0 \qquad (i = 2, \dots, k).$$

Conclusion.

This study implies the following advice to the gamblers: play the numbers that come out most frequently. Most strangely, a lot of gamblers do just the opposite. They commit a double error. First they believe that the roulette is perfect. This is certainly not the case in obscure gamblers clubs, and even not in big casinos, as one of us was told by a manager. Further, they know just enough of the law of large numbers to apply it incorrectly. If a number has not come out for some time, they believe that the law of large numbers sanctions the roulette by forcing its outcome in the next plays. Of course they ignore a statement by Borel, we believe, "La roulette n'a ni yeux, ni mémoire".

6. A restriction on Bühlmann's generalized model

Our next problem is the detection of the stochastic dependence of the contracts (i.e. the columns in the tableau of 3.) in Bühlmann's generalized model, restricted conveniently.

We shall assume that, in any fixed portfolio, corresponding to a realization

$$\theta = (\theta_1, \dots, \theta_k)$$

of the vector

$$\Theta = (\Theta_1, \dots, \Theta_k),$$

the observations in different years $s=1,\ldots,t$, are independent. In fact we assume less. Precisely, we suppose that

$$E(X_{is}/\Theta) = E(X_{is}/\Theta_i)$$
 $(i = 1, \dots, k; s = 1, \dots, t)$

and that constants b, c exist, satisfying the relations

$$Cov(\mu(\Theta_i), \mu(\Theta_j)) = b \qquad (i, j = 1, \dots, k; i \neq j),$$

$$E Cov(X_{ir}, X_{js}/\Theta) = \delta_{rs}c \quad (i, j = 1, \dots, k; i \neq j; r, s = 1, \dots, t).$$

If the contracts are independent, then b = c = 0.

The values of b and c give an idea of the stochastic dependence, more precisely of the correlation, of the contracts. In next section, we indicate how they can be estimated.

7. Parameter estimation in the restricted model

The estimation of b and c is based on the relation

$$\operatorname{Cov}(X_{ir}, X_{js}) = b + \delta_{rs}c \qquad (i, j = 1, \dots, k; \ i \neq j;$$

$$r, s = 1, \dots, t)$$

$$(11)$$

proved in the usual way from the assumptions on the restricted model. Indeed, for $i \neq j$,

$$\operatorname{Cov}(X_{ir}, X_{js}) = \operatorname{Cov}(E(X_{ir}/\Theta), E(X_{js}/\Theta)) + E \operatorname{Cov}(X_{ir}, X_{js}/\Theta)$$

$$= \operatorname{Cov}(E(X_{ir}/\Theta_i), E(X_{js}/\Theta_j)) + \delta_{rs}c$$

$$= \operatorname{Cov}(\mu(\Theta_i), \mu(\Theta_j)) + \delta_{rs}c$$

$$= b + \delta_{rs}c.$$

Estimation of b.

For b we suggest the estimator

$$\widehat{b} = \frac{1}{k(k-1)t(t-1)} \sum_{i \neq j, r \neq s} (X_{ir} - m') (X_{js} - m'), \tag{12}$$

where m' is an approximation of m.

By a proof similar to that of (6), we obtain

$$E\hat{b} = b + (m - m')^2. \tag{13}$$

In the roulette case, we can take m' = m and then \hat{b} is unbiased.

Estimation of c.

For c we suggest the estimator

$$\widehat{c} = \frac{1}{tk(k-1)} \sum_{s} \sum_{i \neq j} (X_{is} - m') (X_{js} - m') - \widehat{b}.$$
(14)

It is unbiased: $E\hat{c} = c$.

Indeed, from (14) results that

$$E(\widehat{b} + \widehat{c}) = b + c + (m - m')^2,$$

and the difference with (13) proves the unbiasedness of \hat{c} , whatever be m'.

General expressions for the estimators.

Let us abbreviate

$$\sum_{ijrs} = \sum_{ijrs} (X_{ir} - m')(X_{js} - m'),$$

$$\sum_{ijs} = \sum_{ijs} (X_{is} - m')(X_{js} - m'),$$

$$\sum_{irs} = \sum_{irs} (X_{ir} - m')(X_{is} - m'),$$

$$\sum_{is} = \sum_{is} (X_{is} - m')(X_{is} - m'),$$

where $i, j = 1, \dots, k$ and $r, s = 1, \dots, t$.

Then

$$\widehat{b} = \frac{1}{k(k-1)t(t-1)} \left(\sum_{ijrs} -\sum_{ijs} -\sum_{irs} + \sum_{is} \right),$$
 (15)

$$\widehat{c} = \frac{1}{tk(k-1)} \left(\sum_{ijs} - \sum_{is} \right) - \widehat{b}, \tag{16}$$

$$\widehat{a} = \frac{1}{kt(t-1)} \left(\sum_{irs} - \sum_{is} \right) \tag{17}$$

From (15) and (17) results

$$\hat{b} = \frac{1}{k(k-1)t(t-1)} \left(\sum_{ijrs} - \sum_{ijs} \right) - \frac{1}{k-1} \hat{a}$$
 (18)

8. Estimation of b and c in the roulette case

In the roulette case we take $m' = m = \frac{1}{k}$.

Estimation of b.

$$\sum_{ijrs} = \sum_{ijrs} (X_{ir}X_{js} - mX_{js} - mX_{ir} + m^2)$$

$$= t^2 - mkt^2 - mkt^2 + m^2k^2t^2$$

$$= t^2 - t^2 - t^2 + t^2$$

$$= 0$$

$$\sum_{ijs} = \sum_{ijs} (X_{is}X_{js} - mX_{is} - mX_{js} + m^2)$$

$$= t - mkt - mkt + m^2k^2t$$

$$= t - t - t + t$$

$$= 0.$$

By (18),

$$\hat{b} = -\frac{1}{k-1}\hat{a}.\tag{19}$$

Taking expectations, we have

$$b = -\frac{1}{k-1}a\tag{20}$$

For a direct proof of (20), we first observe the obvious result:

$$\sum_{i} \Theta_{i} = \sum_{i} E(X_{is}/\Theta_{i}) = E\left(\sum_{i} X_{is}/\Theta\right) = E(1/\Theta) = 1.$$

Then

$$0 = \operatorname{Cov}\left(\sum_{i} \Theta_{i}, \sum_{j} \Theta_{j}\right) = \sum_{ij} \operatorname{Cov}(\Theta_{i}, \Theta_{j})$$
$$= \sum_{i \neq j} \operatorname{Cov}(\Theta_{i}, \Theta_{j}) + \sum_{i} \operatorname{Var}(\Theta_{i})$$
$$= k(k-1)b + ka.$$

Estimation of c.

$$\sum_{is} = \sum_{is} (X_{is}^2 - 2mX_{is} + m^2)$$

$$= \sum_{is} (X_{is} - 2mX_{is} + m^2)$$

$$= t - 2mt + m^2kt$$

$$= t - \frac{t}{k}$$

$$= \frac{1}{k}(k-1)t.$$

Then, by (10) and (19)

$$\widehat{c} = -\frac{1}{k^2} - \widehat{b} = -\frac{1}{k^2} + \frac{1}{k-1} \left(\frac{1}{k} - \widehat{s}^2 - \frac{1}{k^2} \right) = -\frac{1}{k-1} \widehat{s}^2,$$

$$\widehat{c} = -\frac{1}{k-1} \widehat{s}^2.$$
(21)

Taking expectations, we obtain

$$c = -\frac{1}{k-1}s^2. (22)$$

For a direct proof of (22):

$$0 = E \operatorname{Cov}(1, 1/\Theta)$$

$$= E \operatorname{Cov}\left(\sum_{i} X_{is}, \sum_{j} X_{js}/\Theta\right)$$

$$= \sum_{i \neq j} E \operatorname{Cov}\left(X_{is}, X_{js}/\Theta\right) + \sum_{i} E \operatorname{Var}\left(X_{i}/\Theta\right)$$

$$= k(k-1)c + ks^{2}.$$

Remarks.

In the roulette case, the dependence between the "contracts" is negative: $b \le 0$, $c \le 0$. In automobile insurance, fire insurance, ... positive dependencies must be expected.

In the roulette case, \widehat{a} , \widehat{b} , \widehat{c} depend on k and \widehat{s}^2 only, and \widehat{s}^2 only depends on k and Q.

The quality of the estimator \hat{s}^2 increases with t and with k. For small values of t and k, say 2 or 3, the number of observations X_{is} is too small, and \hat{s}^2 can only be a poor estimator of s^2 .

9. Improved credibility estimators in the restricted model

The credibility estimator $\widehat{\mu}_i$ is the linear combination of the random variables $1, X_{i1}, \dots, X_{it}$ closest to $\mu(\Theta_i)$ in the least squares sense. Let μ_i^* be the linear combination of 1 and all the observable random variables X_{js} $(j=1,\dots,k;$ $s=1,\dots,t)$ closest to $\mu(\Theta_i)$ in the least squares sense. In the classical model $\widehat{\mu}_i = \mu_i^*$ because the contracts are independent. Here μ_i^* is a strictly better estimator than $\widehat{\mu}_i$ (except in special cases, where the two estimators may be equal). For symmetry reasons, μ_i^* can be displayed as

$$\mu_i^* = z_1 X_{iM} + z_2 X_{MM} + z_3 m, \tag{23}$$

where X_{iM} is defined in section 3 and

$$X_{MM} = \frac{1}{k} \sum_{i} X_{iM} = \frac{1}{kt} \sum_{is} X_{is}$$

Then the unknown coefficients z_1, z_2, z_3 result from the relations

$$E\mu_i^* = E\mu(\Theta_i) \tag{24}$$

$$\operatorname{Cov}(\mu_i^*, X_{js}) = \operatorname{Cov}(\mu(\Theta_i), X_{js}) \qquad (j = 1, \dots, k; s = 1, \dots, t).$$
(25)

(For instance, see Theorem 5 of De Vylder e.a. (1992)). In order to explicit (25), we need more

Covariance relations.

From the relation following (6) for $Cov(X_{ir}, X_{is})$ and the relation (11) for $Cov(X_{ir}, X_{js})$ $(i \neq j)$, results

$$\operatorname{Cov}(X_{ir}, X_{js}) = \operatorname{Cov}(X_{ir}, X_{is})\delta_{ij} + \operatorname{Cov}(X_{ir}, X_{js})(1 - \delta_{ij})$$

$$= b + c\delta_{rs} + (a - b)\delta_{ij} + (s^2 - c)\delta_{ij}\delta_{rs}$$

$$(i, j = 1, \dots, k; r, s = 1, \dots, t).$$
(26)

From (26) we obtain, for all i, j = 1, ..., k and r, s = 1, ..., t:

$$\operatorname{Cov}\left(X_{ir}, X_{jM}\right) = \operatorname{Cov}\left(X_{iM}, X_{jM}\right)$$
$$= b + (a - b)\delta_{ij} + \frac{1}{t}\left(c + (s^{2} - c)\delta_{ij}\right) \tag{27}$$

$$\operatorname{Cov}\left(X_{ir}, X_{iM}\right) = \operatorname{Cov}\left(X_{iM}, X_{iM}\right)$$
$$= a + \frac{1}{t}s^{2} \tag{28}$$

$$\operatorname{Cov}\left(X_{ir}, X_{MM}\right) = \operatorname{Cov}\left(X_{iM}, X_{MM}\right)$$
$$= b + \frac{1}{k}(a - b) + \frac{1}{t}c + \frac{1}{kt}(b - c). \tag{29}$$

$$\operatorname{Cov}(\mu(\Theta_{i}), X_{js}) = E \operatorname{Cov}(\mu(\Theta_{i}), X_{js}/\Theta) + \operatorname{Cov}(E(\mu(\Theta_{i})/\Theta), E(X_{js}/\Theta)) = 0 + \operatorname{Cov}(\mu(\Theta_{i}), \mu(\Theta_{j})) = a\delta_{ij} + b(1 - \delta_{ij}).$$
(30)

Determination of μ_i^* .

From (23), (24):

$$z_1 + z_2 + z_3 = 1. (31)$$

From (23), (27), (29):

$$Cov (\mu_i^*, X_{js}) = z_1 Cov (X_{iM}, X_{js}) + z_2 Cov (X_{MM}, X_{js})$$
$$= z_1 \left(b + (a - b)\delta_{ij} + \frac{1}{t} (c + (s^2 - c)\delta_{ij}) \right) + z_2 d, \quad (32)$$

where d is the last member of (29).

Hence, from (25) for i = j and for $i \neq j$, using (30):

$$z_1\left(a + \frac{1}{t}s^2\right) + z_2d = a\tag{33}$$

$$z_1\left(b + \frac{1}{t}c\right) + z_2d = b. ag{34}$$

Then z_1, z_2, z_3 result from (31), (33), (34).

If m, in (23), is estimated and replaced by X_{MM} , we obtain

$$\mu_i^* = z_1 X_{iM} + (1 - z_1) X_{MM} \qquad (i = 1, \dots, k). \tag{35}$$

Then we only need z_1 . From (33) and (34) results, taking the difference of these relations:

$$z_1 = \frac{(a-b)t}{(s^2 - c) + (a-b)t} \tag{36}$$

In the roulette case, (20) and (22) imply $z = z_1$, $\hat{\mu}_i = \mu_i^*$.

Best unbiased homogeneous linear estimator.

The estimator μ_i^* defined by (35) is the best (least squares sense) unbiased approximation of $\mu(\Theta_i)$, linear homogeneous in the observable random variables X_{js} $(j=1,\ldots,k; s=1,\ldots,t)$. This results from Theorem 6 of De Vylder e.a. (1992). By that theorem it is enough to verify that

$$E(\mu(\Theta_i) - \mu_i^*) X_{js} = E((\mu(\Theta_i) - m) - (\mu_i^* - m)) (X_{js} - m)$$

= $Cov (\mu(\Theta_i), X_{js}) - Cov (\mu_i^*, X_{js})$

does not depend on the subscripts j, s. By (27), (29), (30), it is easily verified that this is so.

F. De Vijlder, H. Cossette, U.C.L. Place des Doyens 1 B-1348 Louvain-la-Neuve

References

- Bühlmann H. (1967). Experience rating and credibility. The ASTIN Bulletin, vol IV, part II, 199–207.
- De Vylder F., Goovaerts M. (1992). Optimal parameter estimation under zero excess assumptions in a classical model. IME 11, 1–6.

Abstract

We show how Bühlmann's model in credibility theory can be adapted in order to cope with contracts that are not necessarily stochasticly independent. As illustration, outside the insurance world, we estimate the probability that the ball falls in a fixed hole of an imperfect casino roulette.

We also indicate how the dependence between the contracts could be detected in a distribution-free set-up.

Zusammenfassung

Das Modell von Bühlmann in der Credibility-Theorie kann so angepasst werden, dass es auf nicht notwendig unabhängige Verträge angewendet werden kann. Zur Illustrierung ausserhalb des Versicherungswesens schätzen wir die Wahrscheinlichkeit, dass die Kugel bei einem unvollkommenen Roulette in ein festes Loch fällt.

Wir weisen ebenfalls darauf hin, wie die Abhängigkeit zwischen Verträgen ohne die Kenntnis von Verteilungen gefunden werden kann.

Résumé

Nous montrons comment le modèle de Bühlmann de la théorie de la crédibilité peut être modifié afin qu'il puisse également être appliqué lorsque l'indépendance des contrats d'assurance n'est pas garantie. Comme illustration hors du domaine de l'assurance, nous estimons la probabilité que la boule d'une roulette de casino non parfaite s'arrête dans une case donnée.

Nous montrons également comment la dépendance des contrats peut être détectée lorsque les fonctions de distributions ne sont pas connues.