

Zeitschrift:	Mémoires de la Société Vaudoise des Sciences Naturelles
Herausgeber:	Société Vaudoise des Sciences Naturelles
Band:	12 (1958-1961)
Heft:	5
Artikel:	Les viroses du tabac en Suisse
Autor:	Aubert, Olivier
Kapitel:	Summary
DOI:	https://doi.org/10.5169/seals-257914

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

mosaïque du tabac. Les symptômes produits par ces deux types de souches sur divers hôtes sont décrits. En outre, une souche à mosaïque blanche, très virulente, a été isolée.

Le *virus de la mosaïque de la luzerne* est assez répandu, mais peu redoutable. Les symptômes provoqués sur le tabac par diverses souches de ce virus sont décrits.

Le *virus X de la pomme de terre* est très rare sur le tabac.

Au point de vue économique, le virus Y de la pomme de terre est le plus redoutable des virus qui attaquent le tabac en Suisse. Cependant, l'importance des virus de la mosaïque du concombre et de la mosaïque de la luzerne n'est pas négligeable non plus. En revanche, le virus de la mosaïque du tabac et le virus X de la pomme de terre n'ont aucune influence sur l'état sanitaire des plantations de tabac.

SUMMARY

The following viruses were isolated from Swiss tobacco fields : Potato virus Y, Tobacco mosaic virus, Cucumber mosaic virus, Alfalfa mosaic virus and Potato virus X.

Potato virus Y occurred abundantly as the *typical* strain which was harmless to tobacco, and as a *necrotic* strain which caused a disease called *tobacco veinal necrosis*. Until 1957, the necrotic strain had only attacked tobacco of the Burley type. Then in 1958, a new variant appeared which infected the previously resistant Mont-Calme Brun and Mont-Calme Jaune varieties as well as the Burley tobacco.

In greenhouse tests various solanaceous plants were inoculated with the typical and the necrotic strains. The symptoms were compared.

The physical properties of necrotic virus Y were established. The dilution end point was as high as 1/100 000. The temperature of inactivation was at 60° C. The longevity *in vitro* varied from 30 to 70 days depending upon the method of conservation.

Typical and necrotic strains were serologically related, but the presence of the typical strain in the host plant failed to prevent subsequent infection and multiplication of the necrotic strain, and vice-versa.

The necrotic virus Y was transmissible mechanically and by aphids, but it was not seed-borne.

Isolates of lesser virulence could readily be obtained from necrotic virus Y. One such isolate caused little necrosis. Another none. Both were not readily transmitted by aphids. Cross protection studies

showed these isolates to be related more closely to the necrotic rather than to the typical virus Y. They were therefore considered as attenuated variants of the necrotic strain.

Tobacco mosaic virus rarely occurred in tobacco fields. It was never found in plants of the variety Burley R. This variety showed a hypersensitive reaction with the formation of necrotic local lesions.

Cucumber mosaic virus was extremely wide-spread, but did not give very apparent symptoms. Two strains were identified. The first strain occurred frequently and caused a fine mosaic. The second one was rare and caused a mosaic similar to that caused by Tobacco mosaic virus. Various host plants were inoculated with two strains, and the symptoms were described. Another strain of Cucumber mosaic virus was isolated. It was very virulent and caused a white mosaic.

Alfalfa mosaic virus was fairly wide-spread but caused little damage. Tobacco plants reacted differently to the various strains tested.

Potato virus X very rarely occurred in tobacco fields.

From an economical stand point, Potato virus Y was the most damaging virus on tobacco in Switzerland. Less damaging, but still to be reckoned with, were Cucumber mosaic virus and Alfalfa mosaic virus. On the other hand, Tobacco mosaic virus and Potato virus X were of no economical importance.
