

Zeitschrift: Mémoires de la Société Vaudoise des Sciences Naturelles
Herausgeber: Société Vaudoise des Sciences Naturelles
Band: 19 (1991-1999)
Heft: 1: The cytogenetics of the *Sorex araneus* group and related topics

Artikel: Whole-arm rearrangements in the common shrew in a wider context
Autor: Garagna, S. / Searle, J.B. / Redi, C.A.
DOI: <https://doi.org/10.5169/seals-260076>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Whole-arm rearrangements in the common shrew in a wider context¹

BY

S. GARAGNA², J.B. SEARLE³ and C.A. REDI²

Summary.—GARAGNA S., SEARLE J.B. and REDI C.A., 1991. Whole-arm rearrangements in the common shrew in a wider context. In: J. HAUSSER, ed. The cytogenetics of the *Sorex araneus* group and related topics. Proceedings of the ISACC's Second International Meeting. *Mém. Soc. vaud. Sc. nat.* 19.1: 45-49.

The common shrew has one of the most variable karyotypes of any mammal. Recent karyotypic evolution in this species has predominantly involved rearrangements which result in changes in the disposition of chromosome arms without changes within the arms. A chromosome arm may be present as an acrocentric chromosome or as a part of a variety of different metacentric chromosomes. While Robertsonian (centric) fusions are apparently the predominant form of rearrangement, other mechanisms like fissions or whole-arm reciprocal translocations should not be excluded.

The analogous chromosomal variation encountered in the house mouse can be explained by the presence of very homogeneous highly repetitive (satellite) DNA in the centromeric region. Recent data indicate the presence of heterochromatin near the centromeres of the common shrew, except for the evolutionary oldest metacentrics. Preliminary results of *in situ* hybridisation with the highly conserved (TTAGGG)_n telomeric sequence suggests a loss of telomeric sequences during or after fusions.

Résumé.—GARAGNA S., SEARLE J.B. et REDI C.A., 1991. Réarrangements des bras chromosomiques de la musaraigne carrelet considérés dans un contexte plus général. In: J. HAUSSER, dir. The cytogenetics of the *Sorex araneus* group and related topics. Proceedings of the ISACC's Second International Meeting. *Mém. Soc. vaud. Sc. nat.* 19.1: 45-49.

¹This work was supported by the Royal Society of London and a bilateral Project Grant from the Italian C.N.R. (89.02932.04).

²Dipartimento di Biologia Animale and Centro di Studio per l'Istochimica del CNR, University of Pavia, Piazza Botta 10, 27100 Pavia, Italy

³Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, U.K.

La musaraigne carrelet présente le caryotype le plus variable de tous les mammifères. L'évolution récente du caryotype de cette espèce implique surtout des réarrangements qui changent les relations entre les bras chromosomiques, sans changer la structure des bras eux-mêmes. Un bras chromosomique peut être présent sous forme de chromosome acrocentrique ou participer à différents chromosomes métacentriques. Si les fusions Robertsoniennes (ou centriques) constituent vraisemblablement le mécanisme prédominant dans ces réarrangements, d'autres modifications comme les fissions ou les translocations réciproques ne peuvent pas être exclues.

La situation analogue rencontrée chez la souris domestique peut être expliquée par la présence d'ADN hautement répétitif (ADN satellite) très homogène à proximité des centromères. Des données récentes indiquent également la présence d'hétérochromatine près des centromères chez la musaraigne carrelet, sauf pour les métacentriques évolutivement les plus anciens. Les résultats préliminaires d'hybridations *in situ* avec une séquence télomérique très conservée (TTAGGG)_n suggèrent une perte du DNA télomérique pendant ou après le processus de fusion.

In the common shrew (*Sorex araneus*), an extremely large number of chromosomal rearrangements have been described due primarily to centric (Robertsonian: Rb) fusion of the ancestral acrocentric chromosomes giving rise to metacentrics. As a consequence of these frequent rearrangements, there are many karyotypic races of the common shrew all over Europe, making it one of the most interesting mammals for studies of karyotypic evolution.

The role of structural chromosome rearrangements in the speciation process has long been debated (WHITE 1968, 1978, BUSH *et al.* 1977, BUSH 1981, CAPANNA 1982, BAKER and BICKHAM 1986) and a number of hypotheses have been proposed based mainly on the general assumption that karyotypic heterozygosity can perturb the meiotic process, with gene flow between populations prevented by the consequent subfertility or sterility of the hybrids. However, BENGTSSON and FRYKMAN (1990) have argued that simple chromosomal rearrangements do not play a special role in speciation in the common shrew. They cite evidence that gene flow occurs between different shrew populations in contact zones (FRYKMAN 1984, FRYKMAN and BENGTSSON 1984), despite likely heterozygote disadvantage (SEARLE 1988). In addition, an increase of acrocentric chromosomes (a sort of "return" to the ancestral karyotype) has been found in the karyotypes of individuals caught in several shrew hybrid zones (FEDYK and LENIEC 1987, FREDGA 1982, SEARLE 1986) which may further ease flow of genetic information across the zones. The situation found in Britain where new karyotypic races are apparently being formed as a result of modification of hybrid zones, reinforces the idea that speciation, through chromosomal rearrangements, is unlikely to occur in the common shrew (review in SEARLE *et al.* 1990).

Thus, we have some appreciation of the relevance of chromosomal rearrangements to the present-day genetic architecture of the common shrew. However, it remains difficult to understand how such rearrangements become established in this species and, crucially in this regard, what are the molecular processes involved in the structural change of the karyotype. At this point, it must be underlined that although Rb fusions are likely to have been the predominant form of chromosomal rearrangement in the recent karyotypic

evolution of *Sorex araneus* (VOLOBOUEV 1989), we may also have to envisage molecular mechanisms for the following additional categories of rearrangement: Rb fissions (OLERT 1973, HALKKA *et al.* 1987, SEARLE *et al.* 1990), whole-arm reciprocal translocations (WÓJCIK 1986, HALKKA *et al.* 1987) and centromeric shifts (VOLOBOUEV 1989).

To help predict the molecular processes important in the karyotypic evolution of the common shrew, we will digress to a much better studied system: the house mouse, *Mus musculus*. As is well known, a very high frequency of Rb fusions have occurred in *M. musculus domesticus* (CAPANNA *et al.* 1976, GROPP *et al.* 1982, WINKING *et al.* 1988, review in REDI and CAPANNA 1988). The study of the molecular constitution of the pericentromeric region in this species is crucial to our understanding of these chromosomal rearrangements (REDI *et al.* 1990c). We have found that the highly repetitive (satellite) DNA, which is located near the centromeres of all chromosomes except the Y (PARDUE and GALL 1970), is particularly homogeneous and arranged in very long tandem repeats in *M. musculus domesticus* (REDI *et al.* 1990b). This pericentromeric constitution could favour a high frequency of Rb rearrangement through mispairing of homologous sequences of satellite DNA of nonhomologous acrocentrics (REDI *et al.* 1990a). Interestingly, in *M. musculus musculus*, in which the satellite DNA is less homogeneous, only one Rb chromosome has been detected (ZIMA and MACHOLAN 1989).

Even though a wider array of chromosomal rearrangements may occur in the common shrew than in the house mouse, a molecular analysis of the pericentromeric region of autosomes is again of pre-eminent importance for our understanding of the mechanism of chromosomal mutation in this species. Until now, efforts to isolate the satellite DNA from the shrew genome have failed (WORSMAN 1981). However, some preliminary data on the pericentromeric constitution of shrew chromosomes are available. Heterochromatin has been found located around the centromeres of all autosomes except the three pairs of evolutionary-old metacentrics (SEARLE 1983), indicating the presence of highly repetitive DNA. Highly repetitive DNA has also been detected after DNA digestion with restriction enzymes (WORSMAN 1981). Recently, we analysed the pericentromeric regions of Rb chromosomes in the common shrew by *in situ* hybridization with synthetic oligonucleotides that have the highly conserved vertebrate telomeric sequence (TTAGGG)_n (MEYNE *et al.* 1989). Our preliminary data suggest that there is loss of these telomeric sequences during or after fusion.

In addition to satellite DNA, it is probable that other factors, such as transposons (HALKKA *et al.* 1987) or elements able to increase the recombination frequency like minisat-DNAs (WAHLS *et al.* 1990), could be active in the karyotypic evolution of the common shrew at the molecular level.

REFERENCES

- BAKER R.J. and BICKHAM J.W., 1986. Speciation by monobrachial centric fusions. *Proc. Natl. Acad. Sci. USA* 83: 8245-8248.
- BENGSSON B.O. and FRYKMAN I., 1990. Karyotype evolution: Evidence from the common shrew (*Sorex araneus* L.). *J. evol. Biol.* 3: 85-101.
- BUSH G.L., 1981. Stasipatric speciation and rapid evolution in animals. In: W.R. ATCHLEY and D.S. WOODRUFF, eds. Evolution and Speciation: 201-218. Cambridge University Press.
- BUSH G.L., CASE S.M., WILSON A.C. and PATTON J.L., 1977. Rapid speciation and chromosomal evolution in mammals. *Proc. Natl. Acad. Sci. USA* 74: 3942-3946.
- CAPANNA E., 1982. Robertsonian numerical variation in animal speciation: *Mus musculus*, an emblematic model. In: C. BARIGOZZI, ed. Mechanisms of Speciation: 155-177. A. Liss, New York.
- CAPANNA E., GROPP A., WINKING H., NOACK G. and CIVITELLI M.V., 1976. Robertsonian metacentrics in the mouse. *Chromosoma* 58: 341-353.
- FEDYK S. and LENIEC H., 1987. Genetic differentiation of Polish population of *Sorex araneus* L. I. Variability of autosome arm combinations. *Folia Biol. (Kraków)* 35: 57-68.
- FREDGA K., 1982. Distribution of and chromosome polymorphism in chromosome races of the common shrew in Sweden. *Hereditas* 97: 317.
- FRYKMAN I., 1984. An electrophoretic study of genetic differentiation in *Sorex*. Thesis, Department of Genetics, University of Lund.
- FRYKMAN I. and BENGSSON B.O., 1984. Genetic differentiation in *Sorex*. III. Electrophoretic analysis of a hybrid zone between two karyotypic races in *Sorex araneus*. *Hereditas* 100: 259-270.
- GROPP A., WINKING H., REDI C.A., CAPANNA E., BRITTON-DAVIDIAN J. and NOACK G., 1982. Robertsonian karyotype variation in wild house mice from Rhaeto-Lombardia. *Cytogenet. Cell Genet.* 34: 67-77.
- HALKKA L., SÖDERLUND V., SKARÉN U. and HEIKKILÄ J., 1987. Chromosomal polymorphism and racial evolution of *Sorex araneus* L. in Finland. *Hereditas* 106: 257-275.
- MNEYNE J., RATLIFF R.L. and MOYZIS R.K., 1989. Conservation of the human telomere sequence (TTAGGG)_n among vertebrates. *Proc. Natl. Acad. Sci. USA* 89: 7049-7053.
- OLERT J., 1973. A case of chromosome dissociation in a shrew. *Genet. Res.* 22: 323-324.
- PARDUE M.L. and GALL J.G., 1970. Chromosomal localization of mouse satellite DNA. *Science* 168: 1356-1358.
- REDI C.A. and CAPANNA E., 1988. Robertsonian Heterozygotes in the House Mouse and the Fate of their Germ Cells. In: A. DANIEL, ed. The Cytogenetics of Mammalian Autosomal Rearrangements: 315-359. A. Liss, New York.
- REDI C.A., GARAGNA S. and CAPANNA E., 1990a. Nature's experiment with *in situ* hybridization? A hypothesis for the mechanism of Rb fusion. *J. evol. Biol.* 3: 133-137.
- REDI C.A., GARAGNA S., DELLA VALLE G., BOTTIROLI G., DELL'ORTO P., VIALE G., PEVERALI F.A., RAIMONDI E. and FOREJT J., 1990b. Differences in the organization and chromosomal allocation of satellite DNA between the European long tailed house mice *Mus domesticus* and *Mus musculus*. *Chromosoma* 99: 11-17.
- REDI C.A., GARAGNA S. and ZUCCOTTI M., 1990c. Robertsonian chromosome formation and fixation: the genomic scenario. *Biol. J. Linn. Soc. Lond.* 41: 235-255.

- SEARLE J.B., 1983. Robertsonian chromosomal variation in the common shrew *Sorex araneus* L. *Thesis, University of Aberdeen.*
- SEARLE J.B., 1986. Factors responsible for a karyotypic polymorphism in the common shrew, *Sorex araneus*. *Proc. R. Soc. Lond. (B)* 229: 277-298.
- SEARLE J.B., 1988. Selection and Robertsonian Variation in Nature: The Case of the Common Shrew. In: A. DANIEL, ed. *The Cytogenetics of Mammalian Autosomal Rearrangements*: 507-531. A. Liss, New York.
- SEARLE J.B., HÜBNER R., WALLACE B.M.N. and GARAGNA S., 1990. Robertsonian variation in wild mice and shrews. *Chromosomes Today* 10: 253-263.
- VOLOBOUEV V.T., 1989. Phylogenetic relationships of the *Sorex araneus-arcticus* species complex (Insectivora, Soricidae) based on high-resolution chromosome analysis. *J. Hered.* 80: 284-290.
- WAHLS W.P., WALLACE L.J. and MOORE P.D., 1990. Hypervariable Minisatellite DNA is a hotspot for homologous recombination in human cells. *Cell* 60 : 95-103.
- WHITE M.J.D., 1968. Models of speciation. *Science* 159: 1065-1071.
- WHITE M.J.D., 1978. Modes of Speciation. Freeman, San Francisco.
- WINKING H., DULIĆ B. and BULFIELD G., 1988. Robertsonian karyotype variation in the European house mouse, *Mus musculus*: survey of present knowledge and new observations. *Z. Säugetierkunde* 53: 148-161.
- WÓJCIK J.M., 1986. Karyotypic races of the common shrew (*Sorex araneus* L.) from northern Poland. *Experientia* 42: 960-962.
- WORSMAN R.K., 1981. A comparative study of restriction site periodicities in highly repetitive DNA of *Sorex araneus*, *Sorex minutus* and *Neomys fodiens*. *Thesis, University of Aberdeen.*
- ZIMA J. and MACHOLÁN M., 1989. Robertsonian fusion 5.12 in a population of *Mus musculus musculus*. *Folia Zoologica* 38: 233-238.

