

Zeitschrift: Mycologia Helvetica
Herausgeber: Swiss Mycological Society
Band: 1 (1983-1986)
Heft: 1

Artikel: A new species of Lyophyllum from Japan
Autor: Hongo, T. / Clémenton, H.
DOI: <https://doi.org/10.5169/seals-1036450>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 03.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

MYCOLOGIA HELVETICA

Vol. I No 1

1983

A NEW SPECIES OF LYOPHYLLUM FROM JAPAN

T. Hongo

Laboratory of Biology, Faculty of Education
Shiga University, Otsu 520, Japan

and

H. Cléménçon

Institut de Botanique Systématique
Université de Lausanne, Bâtiment de Biologie
CH 1015 Lausanne - Dorigny, Suisse

On close inspection the Japanese fungus described as Lyophyllum transforme (Britz.) Sing. by Imazeki & Hongo (1965) proved to be different from the European L. transforme. Both have been described with "triangular" spores. However, "triangular" is a rough approximation of the projection image of a three-dimensional body into two dimensions. In both species the spore shape can be described using the image of a short banana bearing an enormous dorsal thickening. The spores of the European species bear a high, flattened, dorsal cone centered around the middle of the abaxial side, whereas the Japanese species bears an incomplete, apical ring thickest on the abaxial side and tapering towards the adaxial side where it remains open. It becomes to resemble a fig, and so the Japanese species is called sykosporum, meaning "fig-spored".

LYOPHYLLUM SYKOSPORUM nov. spec.

Pileus e convexo repandus, carnosus, siccus, brunneo-fuliginosus vel fusco-brunneus, subsericeus, tactu nigrescens, 4-9 cm latus. Lamellae subconfertae, postice adnato-rotundatae vel subdecurrentes, pallide

griseae, tactu nigro-maculatae. Stipes fibrilloso-striatus, pallide griseus, fuscescens, apice pruinoso-tomentoso, basi clavatus vel bulbosus, 7-10 cm longus, 10-18 mm crassus, deorsum 20-30 mm crassus, solidus vel farctus. Caro alba, subinodora, mitis, tactu nigro-maculata. Sporae 5,5-8,5 x 4,5-6,5 μ m, ficiformes, laeves, inamyloideae, cyanophilae, uninucleatae. Basidia 28-35 x 8-9 μ m, clavata, fibulata, tetrasporigera, cum granulis siderophilis. Cystidia et cheilocystidia nullae. Pileipellis subgelatinosa, 30-40 μ m crassa, hyphis intertextis vel subregularis, 2-5 μ m crassis, fibulatis. Trama lamellarum regularis.

Hab.: Caespitibus, ad terram silvis abietis.

Holotypus: Hongo 2752 (LAU).

Pileus 4-9 cm broad, convex then expanded, fleshy; surface dry, slightly innately fibrillose, brownish-grey to umbrinous; margin incurved when young; flesh thick, soft, white to greyish white, gradually blackening when injured; taste mild, odour slight. Gills sinuate to adnate-sinuate or adnate-subdecurrent, finally separating; moderately crowded; about 5 mm broad; pale grey, staining black when touched. Stipe 7-10 cm long, 1-1.8 cm thick at apex, 2-3 cm thick at the clavate to bulbous base; fibrillose striate, pruinose at apex; pale grey, becoming darker in age; solid or stuffed.

Spores asymmetrically fig-shaped by an enormous apical thickening tapering towards the abaxial side so that the cross section near the apex looks like a segment roughly a quarter of a circle, therefore irregularly triangular to quadrangular in projection; 5.5-8.5 x 4.5 - 6.5 μ m; uniculate; wall smooth, cyanophilous and siderophilous, inamyloid. Basidia 28-35 x 8-9 / 3-4 μ m; four spored; clamped; with siderophilous granules. Cystidia and marginal cells absent. Pileipellis subgelatinized, the hyphae being separated by a space about 1-2 times their diameter; hyphae cylindrical, 2-5 μ m broad, interwoven on the disk, more radially arranged near the margin of the pileus, clamped; pigment intrusting and interhyphal. Hymenophoral trama regular; hyphae clamped, 8-16 μ m broad in the mediostratum. No gloeopleurous hyphae seen.

Cepitose, gregarious or in small clusters, on the ground in woods, especially of conifers. Summer and autumn. Japan.

Material studied: Hongo 3164 and 2752 (type, LAU).




Figure 1: *Lyophyllum sykosporum* Hongo & Clç. Bar = 1 cm.




Figure 2: Lyophyllum sykosporum Hongo & Clç., spores and basidia.
Bar = 10 μ m.