

**Zeitschrift:** Mycologia Helvetica  
**Herausgeber:** Swiss Mycological Society  
**Band:** 1 (1983-1986)  
**Heft:** 3

**Artikel:** Flaviporus semisupiniformis (Polyporaceae) in Italy  
**Autor:** Ginns, J. / Bernicchia, Annarosa  
**DOI:** <https://doi.org/10.5169/seals-1036462>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# MYCOLOGIA HELVETICA

Vol. I No 3

1984

Manuscript received on 25th February 1984)

## FLAVIPORUS SEMISUPINIFORMIS (POLYPORACEAE) IN ITALY

J. Ginns

Biosystematics Research Institute, Agriculture Canada  
Ottawa, Ont., Canada K1A 0C6

and

Annarosa Bernicchia  
Instituto de Patologia Vegetale, Universita Degli Studi,  
40126 Bologna, Italia

Several interesting or rare fungi were collected in the Mesola forest of Italy. One polypore, identified as *Flaviporus semisupiniformis* is of interest because this species was known only from Mexico (Ginns 1980). A description of the Italian specimen is given and it is compared with the type specimen.

### FLAVIPORUS SEMISUPINIFORMIS (Murr.) Ginns

Basidiome when dry pileate, lobed, overall 3 cm broad and deep, and 1.3 mm thick, broadly attached; the margin white, densely cottony, even to slightly lobed, up to 0.5 mm wide; pileus surface of matted, radiating fibrils or with a glabrous, thin crust, pallid to pale yellowish brown; context pallid, woody but soft, becoming fibrous when wetted, up to 0.2 mm thick; pore layer 1.1 mm thick the pores somewhat collapsed and angular, thin-walled, (5-)6 per mm, the mouths even, granulose (i.e., appearing frosted).

Hyphal system dimitic. Hyphae inamyloid, non-dextrinoid and acyanophilous. Context generative hyphae branched, hyaline, thin to moderately thick-walled, 2.5–5 $\mu\text{m}$  diam., bearing large, prominent clamp connections; skeletal hyphae rarely branched, rather straight, aseptate, 3–5 $\mu\text{m}$  diam., the walls pale yellow, smooth, 0.8–2 $\mu\text{m}$  thick. Tramal generative hyphae as in the context but the skeletal hyphae straight to slightly wavy, 2.5–3.5(–4) $\mu\text{m}$  diam., the walls 0.5–1 $\mu\text{m}$  thick with some hyphae more contorted, infrequently branched and slightly narrower. Cystidia (Fig. 1) common, scattered throughout the trama, some projecting through the hymenium, cylindrical with the apex clavate to lanceolate, thick-walled, about 40–100 X 7–9 $\mu\text{m}$ , approx. 40 $\mu\text{m}$  of the apical part incrusted; cystidioles lacking; basidia (Fig. 1) clavate, 12–16 X 5–6 $\mu\text{m}$ , narrowed to 3 $\mu\text{m}$  diam. at




Figure 1: *Flaviporus semisupiniformis*. Basidiospores, basidium and crystalline incrusted cystidial apex. Bar = 5 $\mu\text{m}$ .

the base, with 4 sterigmata, 3 $\mu\text{m}$  long; spores (Fig. 1) broadly ellipsoid, some subglobose, 3.6-4.2 X 2.5-3.2 $\mu\text{m}$ , the wall hyaline, thin, smooth, acyanophilous, inamyloid, non-dextrinoid, with a minute apiculus.

Specimen examined: Italia, Mesola Forest (Ferrara), 22 Oct. 1981, A. Bernicchia 341 (Bologna and DAOM 185538).

Substratum: On *Fraxinus oxycarpa* trunk lying on the ground.

The excellent condition of the specimen when collected has allowed a more detailed description of the basidia. The type specimen, re-described by Ginns (1980), differed in having smaller pores (10 per mm), cystidia confined to the lower half of the tube-layer, slightly smaller spores (2.5-3 $\mu\text{m}$  in diam.), and scattered cystidioles. The cystidia and gross morphology of the Italian collection were identical with the illustrations of the type (Ginns 1980). Additional collections from Mexico and Italy will be needed before the significance of these differences can be evaluated.

This fungus was collected in the Mesola forest (Ferrara, Italy) which extends along the Adriatic coast, about half-way between Ravenna and Venice. It belongs to the *Quercion-ilicis* vegetation type and was probably established during the tenth century on lands which had emerged from the sea. This forest, one of the few remaining parts of the vast forest that once covered the northern part of Italy, is now protected as a nature reserve.

#### BIBLIOGRAPHY

- Ginns, J. 1980. The genus *Flaviporus* Murrill (Polyporaceae). Canadian J. Bot. 58: 1578-1590.

*Zusammenfassung*

In den Wäldern von Mesola (Ferrara, Italien) wurde *Flaviporus semisupiniformis* (Murr.) Ginns gefunden, ein Porling, der bis anhin lediglich aus Mexiko bekannt war. Der italienische Fund wird beschrieben und mit dem Typus verglichen.

*Résumé*

On a trouvé dans les forêts de Mesola (Ferrare, Italie) *Flaviporus semisupiniformis* (Murr.) Ginns, un Polypore qui n'était connu jusqu'ici qu'au Mexique. Les carpophores trouvés en Italie sont décrits et comparés à l'espèce type.

*Riassunto*

Nel bosco della Mesola (Ferrara, Italia) è stato trovato *Flaviporus semisupiniformis* (Murr.) Ginns, un poliporo che finora era conosciuto soltanto dal Messico. Viene descritto il ritrovamento italiano e si fa pure un confronto con il tipo.