

Zeitschrift:	Mycologia Helvetica
Herausgeber:	Swiss Mycological Society
Band:	3 (1988-1990)
Heft:	3
Artikel:	Bambusicolous fungi collected in southwestern France : 1982-1989
Autor:	Petrini, Orlando / Candoussau, Françoise / Petrini, Liliane E.
DOI:	https://doi.org/10.5169/seals-1036536

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

MYCOLOGIA HELVETICA

Vol. 3 No 3

pp. 263—279

1989

(Manuscript received on 14th July 1989)

BAMBUSICOLOUS FUNGI COLLECTED IN SOUTHWESTERN FRANCE 1982 - 1989

ORLANDO PETRINI¹⁾, FRANÇOISE CANDOUSSAU²⁾ AND LILIANE E. PETRINI¹⁾

¹⁾ Mikrobiologisches Institut, ETH-Zentrum/LFV,
8092 Zürich, Switzerland

²⁾ 53, rue Aristide Briand, 64000 Pau, France

Abstract: More than 60 taxa in the Ascomycotina and Deuteromycotina collected between 1982 and 1989 on bamboo culms and leaves in the Southwest of France are listed. Two new species, *Aulographum emiliae* and *Hadrotrichum bambusicola* are described. Some interesting records are briefly discussed and illustrated.

Zusammenfassung: Mehr als 60 Taxa der Ascomyceten und Deuteromyceten, zwischen 1982 und 1989 auf Stängeln und Blättern von Bambus in Südwestfrankreich gesammelt, werden aufgelistet. Zwei neue Arten, *Aulographum emiliae* und *Hadrotrichum bambusicola* werden beschrieben. Einige interessante Funde werden kurz besprochen.

In the last few years one of us (F.C.) has been extensively collecting ascomycetes, basidiomycetes and deuteromycetes growing on several species of bamboos. CANDOUSSAU et al. (1985), BOIDIN et al. (1986), and SPOONER & CANDOUSSAU (1988) have described new or interesting species of fungi growing on this substrate. We now present a comprehensive list of all ascomycetes and deuteromycetes collected so far on bamboo culms in the Southwest of France.

All collections are from dead culms of bamboos growing in well-sheltered, humid sites protected from wind exposure. More than 40 taxa in the Ascomycotina and about 20 Deuteromycotina have been collected. While some of them are cosmopolitan species able to grow on many Poaceae, a number of taxa (e.g. *Eutypella leprosa*, *Hypoxyylon bipapillatum*, *Herpotrichia herpotrichioides*, *Lophiostoma macrostomum*) are unusual records for these hosts. The common occurrence of some fungal species so far reported only from tropical regions of the world can be explained at least partly by their host-specificity, as it can be assumed that they were introduced in France

with their hosts, and by the apparently very favourable climatic conditions typical of the Southwest of France (CANDOUSSAU et al., 1985).

Collecting sites

Saprobic ascomycetes and deuteromycetes growing on dead culms and sheaths of *Arundinaria japonica* SIEB. et ZUCC. ex STEUDEL, *Phyllostachys bambusoides* SIEB. et ZUCC., *P. mitis* RIVIERE, *P. nigra* (LODD) MUNRO, and *P. viridiglaucescens* (CARRIERE) A. et C. RIVIERE have been collected in following Départements (Official French designation no. of the département in brackets) and sites:

Gard (30):

Anduze (Alès, Bambouseraie de Prafrance)

Gers (32):

Mirande, Loubersan (Mirande)

Gironde (33):

Bernos - Beaulac (N 134, Pau - Bordeaux).

Landes (40):

Peyrehorade, Capbreton, Momuy (Orthez) Habas, Tartas, Seignosse.

Pyrénées Atlantiques (64):

Pau, Gan, Assat, Lavie (Salies de Béarn), Sauveterre de Béarn, Osserain, Uhart-Mixe, Viven, Mont (Orthez), Biarrotte, Biarritz.

as well as in the Pyrénées National Park of Torla (Spain).

Unless otherwise stated, the fungi were found on the dead culms of their hosts. The collections are kept in the private Herbarium of F.C. or in ZT.

ASCOMYCOTINA

1. PEZIZALES

Scutellinia hirtella (REHM) KUNTZE

Description: LE GAL (1966)

Collection examined: 64 Pau, on *Phyllostachys bambusoides*, FC 24.11.1984 (FC 3839).

2. OSTROPALES

Stictis stellata WALLR.

Description and illustrations: SHERWOOD (1977)

Collections examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 3.5.1983; 40 Capbreton, Bord du Boudigau, on *Phyllostachys* sp., FC 9.6.1984

(FC 4686); 64 Assat, Maison Diriart, on *P. bambusoides*, FC 29.11.1984 (FC 4686A); 64 Osserain, on *P. bambusoides*, FC 15.01.1989 (FC 4686B).

FC 4686A has a peculiar pinkish colour and its periphysoids are unbranched; the hymenium has little or no iodine reaction. Similar collections have also been recorded from the Southern USA by M. SHERWOOD (in litt. to FC, 29.11.1984).

3. SPHAERIALES

Amphisphaeria striatispora F. CANDOUSSAU et H. KATUMOTO

Description and illustrations: CANDOUSSAU et al. (1985)

Collections examined: 40 Capbreton, Tombstone, on dead sheaths of *Arundinaria japonica*, FC 11.6.1984 (FC 5604); same site and host, FC 16.12.1984 (5604A) and FC 9.6.1986 (5604B).

Anthostomella nitidissima (DURIEU et MONT.) SACC.

Description: SACCARDO (1882)

Collection examined: 40 Capbreton, Tombstone, on *Arundinaria japonica*, FC 16.12.1984 (FC 5205-4).

This species is near *A. punctulata* (FRANCIS, 1975) but the ascospores are slightly larger and more compressed. It also occurs on culms whereas *A. punctulata* is found on leaves (S.M. FRANCIS, pers. comm.).

Apiospora montagnei SACC.

Description and illustrations: MÜLLER & VON ARX (1962); ELLIS (1971).

Collections examined: This fungus is present everywhere on bamboos throughout the collecting sites. Its anamorph, belonging to the genus *Arthrinium* Kunze: Fr. is almost always associated with it (e.g. FC 5386A, FC 5386B, FC 5386C).

Chaetosphaeria vermicularioides (SACC. et ROUM.) W. GAMS et V. HOL.-JECH.

Description and illustrations: GAMS & HOLUBOVA -JECHOVA. (1976)

Collections examined: 64 Viven, Château de Viven, on *Phyllostachys mitis*, FC 25.11.1984 (FC 6232); 40 Capbreton, Bord du Boudigau, on *P. mitis*, FC 30.11.1986 (FC 6232A).

Eutypella leprosa (FR.) BERLESE

Description and illustrations: RAPPAZ (1987)

Collections examined: 40 Peyrehorade, Landes, 40 km from Pau, on *Phyllostachys cf. bambusoides*, FC 29.11.87 (FC 5427); Sauveterre de Béarn, on *Bambusa* sp., FC 13.5.1988 (FC 180).

Eutypella scoparia (SCHWEIN.: FR.) ELLIS et EVERH.

Description and illustrations: RAPPAZ (1987).

Collection examined: 64 Osserain, on *Phyllostachys bambusoides*, FC 15.1.1989 (FC 200).

Gibberella zeae (SCHWEIN.) PETCH

Description and illustrations: BOOTH (1971)

Anamorph: *Fusarium graminearum* SCHWÄBE

Collections examined: 40 Tartas, Bord de la Midouze, on *Arundinaria japonica*, FC 2.12.1984 (FC 4878A); 40 Capbreton, Monument, on *Arundinaria japonica*, FC 16.12.1984 (FC 4878).

Hypocrea rufa (PERS.: FR.) FR.

Description and illustrations: DENNIS (1978)

Collections examined: 64 Pau, Hippodrome, on *Arundinaria japonica*, FC 4.2.1984 (FC 4805A); 64 Biarrotte, on *Phyllostachys bambusoides*, FC 10.3.1985 (FC 4805B); 64 Assat, Maison Diriart, on *P. bambusoides*, FC 15.6.1986 (FC 4805); 64 Uhart-Mixe, on *P. bambusoides*, FC 28.9.1986 (FC 4805C).

Hypoxyton bipapillatum BERK. et CURT.

Description and illustrations: PETRINI & MÜLLER (1986, sub *H. serpens* var II); PETRINI & ROGERS (1986)

Collection examined: 64 Assat, Maison Diriart, on *Phyllostachys bambusoides*, FC 15.6.1986 (FC 5237).

Hypoxyton serpens (PERS.: FR.) KICKX

Description and illustrations: PETRINI & MÜLLER (1986); PETRINI & ROGERS (1986).

Collection examined: 30 Anduze, Bambouseraie de Prafrance, Generargues, on *Phyllostachys viridis glaucescens*, FC 21.11.87 (FC 174).

Hypoxyton serpens (PERS.: FR.) KICKX var. *macrosporum* J.H. MILLER

Description and illustrations: PETRINI & ROGERS (1986).

Collection examined: 64 Uhart-Mixe, on *Phyllostachys bambusoides*, FC 28.9.1986 (FC 5286).

A taxon widely distributed in the USA, *H. serpens* var. *macrosporum* has so far only once been collected in Europe on broad-leaved trees (POUZAR, 1985; PETRINI & ROGERS, 1986).

Nectriopsis squamulosa (ELLIS) SAMUELS

Description and illustrations: SAMUELS (1988)

Collections examined: 64 Tartas, Landes, on *Phyllostachys*, G. Gilles 23.11. 1984 (FC 4862); 64 Pau, Hippodrome, on *P. mitis*, FC 6.3.1985 (FC 4862A); 64 Pau, La Ventourlère on *P. mitis*, FC 10.3.1983 (FC 4862B); 64 Assat, maison Diriart, on *P. mitis*, FC 11.3.1984 (FC 4862C); 64 Biarotte, on *P. bambusoides*, FC 10.3.1985 (FC 4862D); 40 Capbreton, Bord du Boudigau, on *P. mitis*, FC 9.6.1986 (FC 4862E).

Peroneutypa heteracantha (SACC.) BERL.

Description and illustrations: DENNIS (1978)

Collections examined: 64 Pau, Hippodrome, on *Phyllostachys bambusoides*, FC 6.3.1983 (FC 5460A); 64 Mont, on *Arundinaria japonica*, FC 5.2.1984 (FC 5460B); 64 Assat, Maison Diriart, on *Phyllostachys mitis*, FC 12.2.1984 (FC 5460).

This species, almost always associated with *Graphium fasciculatum* SACC., is very frequent in all collecting sites visited.

Rosellinia muroiana HINO et KATUMOTO

Description and illustrations: HINO & KATUMOTO (1958).

Collections examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 20.3.1983 (FC 5379); 64 Mont, on *Arundinaria japonica*, FC 5.2.1984 (FC 5379A); 64 Osserain, on *Phyllostachys bambusoides*, FC 25.3.1984 (FC 5379B); 64 Assat, Maison Diriart, on *P. bambusoides*, FC 5.6.1986 (FC 5379C); 64 Osserain, on *P. bambusoides*, FC 15.1.1989 (FC 5379D).

R. muroiana is probably a synonym of *Astrocystis sublimbata* (DUR. et MONT.) S. HUGHES [= *R. sublimbata* (DUR. et MONT.) PASS.] and it shows striking similarities to *R. amblystoma* BERL. et SACC. Unfortunately, it was not possible to study the type material of *R. muroiana*: it is very likely, however, that the three taxa are conspecific, in which case *R. amblystoma* and *R. muroiana* would have to be removed from the genus *Rosellinia* and relegated into *Astrocystis*.

Rosellinia sp.

Fig. 1c

Stromata nearly globose, 0.7-1.1 mm diam., brown, dark brown to black in age, gregarious and sometimes confluent. **Subiculum** in young stage covering the stromata, light brown, felted, gradually disappearing and eventually absent in old material.- **Perithecia** 550-720 µm diam.- **Asci**: spore bearing part 180-205 µm, stipe 10-40 µm, total 215 - 230 x 10 µm.- **Ascal tip**

7-11 x 4-5 µm wide, staining blue in Melzer's reagent.- Ascospores 25-39 x 6-8 µm, narrowly ellipsoidal with one pinched end, light brown, with short, sigmoid germslit. Appendages or slimy sheaths not observed.

Collections examined: 64 Sauveterre de Béarn, on *Phyllostachys bambusoides*, FC 20.2.1983 (FC 52, ZT 3.118); same site, on *Phyllostachys* sp., FC 22.3.1987 (FC 52A, ZT 3.152).

This collection cannot be assigned to any known species of *Rosellinia* on bamboos. It may be a new taxon but we prefer to restrain from describing it as new until the study of additional material will be possible.

Roussöëlla verruculosa F. CANDOUSSAU et K. KATUMOTO

Description and illustrations: CANDOUSSAU et al. (1985)

Collections examined: 33 Bernos Beaulac, Hôtel Theas, on *Phyllostachys mitis*, FC 20.3.1984 (FC 5396); 64 Sauveterre de Béarn, on *P. bambusoides*, J. Vivant & FC 31.12.1982 and a second collection in June 1986, same site (FC 5396A); 64 Osserain, on *P. bambusoides*, FC 15.1.1989 (FC 5396B).

Trichosphaeria pilosa (PERS.) FUCK.

Description and illustration: VON ARX & MÜLLER (1954)

Collection examined: 40 Capbreton, Monument, on *Arundinaria japonica*, FC 24.6.1984 (FC 5654).

Zignoëlla pulviscula (CURR.) SACC. sensu MUNK 1957

Description and illustrations: MUNK (1957); BOOTH (1957, sub *Chaetosphaeria*)

Collections examined: 64 Lavie, on *Phyllostachys nigra*, FC 25.3.1984 (FC 5662); 64 Assat, Maison Diriart, on *P. mitis*, FC 15.6.1986 (FC 5662A); 64 Uhart Mixe, on *P. bambusoides*, FC 7.12.1986 (FC 5662B).

4. LEOTIALES

Arachnopeziza aurelia (PERS.) FUCK.

Description and illustrations: KORF (1981)

Collection examined: 64 Osserain, near Sauveterre, on *Phyllostachys* sp., FC 3.7.1988 (FC 4375).

Bisporella scolochloae (DE NOT.) SPOONER

Description: KIRK & SPOONER (1984).

Collections examined: 64 Assat, Maison Diriart, on *Phyllostachys mitis*, FC 12.2.1984; same site and host, FC 11.6.1986 (FC 4296); 64 Osserain, on *P.*

bambusoides, FC 15.1.1989 (FC 4296A); 40 Tartas, on *Phyllostachys* sp., FC 23.1.1989 (FC 4296B).

B. scolochloae occurs on stromatic pyrenomycetes and on culms of various larger grasses. The bright colour and the 3-septate ascospores are good diagnostic characters.

Crocicreas bambusicola CARPENTER

Description & illustrations: CARPENTER (1981)

Collection examined: 40 Capbreton, Bord du Boudigau, on sheaths of *Phyllostachys mitis*, FC 9.6.1986 (FC 4242).

Dasyscyphus barbatus (KUNZE) MASSEE

Description and illustrations: DENNIS (1978)

Collection examined: Spain, Torla, on *Arundinaria japonica*, FC 18.6.1973 (FC 4355).

Durella brunnea HINO et KATUMOTO

Description and illustrations: HINO & KATUMOTO (1955)

Collection examined: 64 Osserain, near Sauveterre, on *Phyllostachys* sp., FC 3.7.1988 (FC 4494).

Encoelia mollisioides B. SPOONER

Description and illustrations: SPOONER & CANDOUSSAU (1988)

Collection examined: 40 Capbreton, Le Boudigau, on sheaths of *Phyllostachys mitis*, FC 9.6.1986 (FC 152).

Lasiostictella bambusae M. SHERWOOD-PIKE

Description and illustrations: CANDOUSSAU et al. (1985)

Collections examined: 40 Capbreton, Tombstone, on *Arundinaria japonica*, FC 15.5.1984 (FC 4667); same site and host, FC 15.5.1984.

Mollisia ligni (DESM.) KARST. - complex

Description and illustrations: DENNIS (1978).

Collection examined: 64 Assat, Maison Diriart, on *Phyllostachys bambusoides*, FC 15.6.1986 (FC 4482).

The problems related to the identification of *Mollisia* species have been highlighted by several authors (e.g. AEBI, 1972; NANNFELDT, 1985). We believe that *M. ligni* is a complex species which may be split in different taxa after careful study.

Mollisia ventosa (KARST.) KARST.

Description and illustrations: DENNIS (1978)

Collections examined: 40 Tartas, Bord de la Midouze, on *Phyllostachys mitis*, G. Gilles 24.2.1984; 64 Lavie, on *Arundinaria japonica*, FC 25.3.1984 (FC 4476).

Pezizella pulcherrima B. SPOONER et R.W.G. DENNIS

Description and illustrations: SPOONER & DENNIS (1985)

Collection examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 29.3.1987 (FC 4297).

P. pulcherrima has been described from decorticated wood of *Ulmus* in the Hebrides (SPOONER & DENNIS, 1985). Morphologically our collection cannot be distinguished from the type material. This is the second known record of the species.

Strossmayeria basitricha (SACC.) DENNIS

Description and illustrations: DENNIS (1978)

Anamorph: *Pseudospiropes* sp. fide ITURRIAGA & KORF (1984).

Collection examined: 40 Tartas, Bord de la Midouze, on *Phyllostachys mitis*, FC 22.1.1984 (FC 4302-1).

5. DOTHIDEALES

Astrophaeriella trochus (PENZ. et SACC.) D. HAWKSW.

Description and illustrations: HAWKSWORTH (1981)

Collections examined: 64 Osserain, on *Phyllostachys bambusoides*, FC 25.3.1984 (FC 6170); same site and host, FC 15.1.1989 (FC 6170A).

Aulographum emiliae sp. nov.

Fig. 2 a-d

Ascomata superficialia, solitaria vel gregaria, specie variabili, 500-600 µm longa, 70-80 µm lata adque 60 µm alta. Paries ascomatis ex aequilateralibus, crassitunicatis, brunneis, 5-7 µm diam. cellulis composita. Asci clavati, bitunicati, 30-35 x 8.5-10 µm. Paraphysoides filiformes, 1 µm latae, longiores ascis. Ascospores bicellulatae, hyalinae dum juvenes, pallide brunneae aetate proiecta, cellula basali minime longiori apicali, 9.5-11 x 3-4 µm.

Habitat in culmis emortuis bambusarum.

Holotypus in culmis emortuis *Phyllostachidis bambusoidis*, Gallia, in Pago dicto Assat, domo Diriart, leg. F. Candoussau 11.3.1984 (ZT, FC 149).

Ascomata superficial, single or in groups, variable in shape, 500-600 µm long, 70-80 µm wide and approx. 60 µm tall. Ascomatal wall composed of equilateral, thick-walled, brown cells, 5-7 µm diam.- Asci clavate, bitunicate, 30-35 x 8.5-10 µm.- Paraphysoids filiform, 1 µm wide, longer than asci.- Ascospores two-celled, hyaline when young, light brown when ripe, the

basal cell slightly longer than the apical and often biguttulate, 9.5-11 x 3-4 µm.

Habitat: on dead bamboo culms.

Holotype: 64 Assat, Maison Diriart, on *Phyllostachys bambusoides*, FC 11.3.1984 (FC 149, deposited in ZT)

Etymology of the species name: dedicated to our friend and teacher Emil Müller.

Other collections examined: 64 Gan, Montjoly, on *P. bambusoides*, FC 16.2.1986 (FC 150); 40 Capbreton, Le Boudigau, on *Phyllostachys* sp., FC 9.6.1986 (FC 151); 40 Seignosse, on *P. nigra*, FC 25.9.1986 (FC 152).

Byssosphaeria schiedermayeriana (FUCKEL) BARR

Description and illustrations: BARR (1984)

Collection examined: 40 Capbreton, Monument, on *Arundinaria japonica*, FC 11.6.1984 (FC 6182).

Gloniopsis praelonga (SCHW.) ZOGG

Description and illustrations: ZOGG (1962)

Collections examined: 64 Artix, Réserve, on *Phyllostachys mitis*, FC 5.3.1984 (FC 6370A); 64 Mont, on *Arundinaria japonica*, FC 4.2.1984 (FC 6370B); 64 Assat, Maison Diriart, FC 11.3.1984; 40 Capbreton, Monument, on *A. japonica*, FC 13.5.1984 (FC 6370).

Glonium abbreviatum (SCHW.) LOHMAN

Description and illustrations: ZOGG (1962)

Collection examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 20.3.1983 (FC 6356).

Herpotrichia herpotrichoides (FUCKEL) CANNON

Description and illustrations: BARR (1984)

Collection examined: 40 Capbreton, Monument, on *Arundinaria japonica*, FC 11.6.1984 (FC 6181).

Hysterium angustatum ALB. & SCHW.

Description and illustrations: ZOGG (1962)

Collections examined: 32 Mirande, on *Phyllostachys mitis*, FC 27.11.1983 (FC 6357); 64 Lavie, on *P. nigra*, FC 25.3.1984 (FC 6357A).

Lophiostoma macrostomum (TODE: FR.) CES. et DE NOT.

Description and illustrations: HOLM & HOLM (1988)

Collections examined: 32 Loubersan, La Flourette, on submerged stems of *Arundinaria japonica*, FC March 1972 (FC 6080); same host and site, FC October 1972 (FC 6080A).

A fungus so far found mainly on conifers, *L. macrostomum* has never been recorded on bamboos. Its taxonomic position has been recently discussed by HOLM & HOLM (1988).

Lophiostoma semilibereum (DESM.) CES. et DE NOT.

Description and illustrations: HOLM & HOLM (1988)

Collections examined: 64 Sauveterre, on *Phyllostachys* sp., FC 7.12.1986 (FC 6083); 64 Uhart-Mixe, on *P. bambusoides*, FC 7.12.1987 (FC 6083A).

Rhizodiscina lignyota (FR.) HAFELLNER

Description and illustrations: HAFELLNER (1979)

Collection examined: 64 Biarrotte, on *Phyllostachys bambusoides*, FC 10.3.1985 (FC 4544).

Schizophyrium pomi (MONT.: FR.) VON ARX

Fig. 2e-g

Description and illustrations: VON ARX (1959).

Collection examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 7.4.1985 (FC 4757).

Sporormiella pilosa (CAIN) AHMED et CAIN

Description and illustrations: AHMED & CAIN (1972)

Collection examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 20.3.1983 (FC 6043)

6. SORDARIALES

Chaetomium olivaceum COOKE et ELLIS

Description and illustrations: AMES (1963); DREYFUSS (1975)

Collections examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 28.4.1985 (FC 4802); same site and host, FC 29.3.1987 (FC 4802A).

DEUTEROMYCOTINA

1. HYPHOMYCETES

Acremonium hyalinulum (SACC.) GAMS

Description and illustrations: GAMS (1971)

Collection examined: 40 Tartas, Bord de la Midouze, on *Phyllostachys* sp., FC 29.3.1986 (ZT).

Annellodochium ramulisporum DEIGHTON

Description and illustrations: ELLIS (1971)

Collection examined: 40 Peyrehorade, hyperparasite on *Peroneutypa heteracantha*, on *Phyllostachys* sp., FC 29.11.1987 (FC 174).

Annellodochium cf. ramulisporum DEIGHTON

Collection examined: 64 Uhart et Mixe, on *Phyllostachys bambusoides*, FC 4.11.1984 (ZT)

We were not able to detect any fungus parasitized by this deuteromycete. The size and shape of the sporodochia and of the conidia fit very well in the original description of *A. ramulisporum*: the unusual habitat suggests at least physiological differences for this collection.

Arthrinium anam. *Apiospora montagnei*

See *Apiospora montagnei*

Arthrinium phaeospermum (CDA) M.B. ELLIS

Description and illustrations: ELLIS (1971)

Collection examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 28.3.1986 (ZT).

Chalara inflatipes (PREUSS) SACC.

Description and illustrations: NAGRAJ & KENDRICK (1975); ELLIS (1976)

Collection examined: 64 Biarrotte, on *Phyllostachys bambusoides*, FC 10.3.1985 (ZT).

Corynespora foveolata (PAT.) HUGHES

Description and illustrations: ELLIS (1971)

Collection examined: 40 Tartas, on *Phyllostachys mitis*, FC & G. Gilles 22.1.1984 (ZT).

Fusarium aquaeductum (REDLK. et RABENH.) LAGERBERG

Description and illustrations: BOOTH (1971).

Collection examined: 40 Tartas, on *Phyllostachys* sp., FC & G. Gilles, 22.1. 1984 (ZT)

Fusarium graminearum SCHWÄBE

see *Gibberella zae*

Graphium fasciculatum SACC.

See *Peroneutypa heteracantha*

Gonytrichum anam. *Trichosphaerella inaequalis* (GROVE ex BERL. et VOGL.)

E. MÜLLER

Description and illustrations: ELLIS (1971)

Collections examined: 64 Osserain, on *Phyllostachys bambusoides*, FC 22.4.1984 (FC 002B); 30 Anduze, Bambouseraie de Prafrance, Generargues, on *P. viridiglaucescens*, FC 21.11.87 (FC 174).

Hadrotrichum bambusicola O.PETRINI, L. E. PETRINI et F. CANDOUSSAU sp. nov.

Fig. 1a-b

Conidiomata ad 2 mm diametro, gregaria ad confluentia, disciformia, primum hyalina, dein pallide brunnea aetate provecta, cellulis conidiomatis in textura prismatica ordinatis, 2-4 µm diam. Conidiophora paene hyalina ad pallide brunnea, rare septata, 10-17 x 2.5-4.5 µm, in strato compacto ordinata. Cellulae conidiogenae polyblasticae, terminales, 12 x 4.5 µm. Conidia solitaria, continua, ellipsoidea, hyalina vel aetate provecta pallide brunnea, laevia, basi truncata, 4-8 x 2.5-3.5 µm.

Habitat in culmis emortuis bambusae.

Holotypus: Galliae, in regione pyrenaica, pago dicto Osserain, in culmis emortuis bambusae, legit F. Candoussau, 3.7.1988 (ZT)

Conidiomata 0.7-2 mm diam., circular to slightly elliptical, sporodochial, white when young, later turning light brown in the centre with a white, sterile margin, gregarious, sometimes confluent. Cells of sporodochial tissue of textura prismatica, hyaline, 2-4 µm diam.- Conidiophores 10-17 x 2.5-4.5 µm, unbranched, aseptate or rarely septate, pale brown, in a compact layer.- Conidiogenous cells up to 12 x 4.5 µm, polyblastic, terminal.- Conidia 4-8 x 2.5-3.5 µm, ellipsoidal, with truncate base, hyaline to pale brown, smooth.

Habitat: on dead culms of bamboos.

Holotype: 64 Osserain, on a dead bamboo culm, F. Candoussau 3.7.1988 (ZT)

H. bambusicola differs from the other known species of *Hadrotrichum* [*H. virescens* SACC. et ROUM., *H. globiferum* (ELL. et EVERH.) DAVIS, *H. phragmitis* FUCKEL, *H. pyrenaicum* PETRINI et al.] by the morphology and size of the conidia and of the conidiophores. *H. bambusicola* could eventually prove to be the anamorph of a bambusicolous *Hypoxylon* species, for its conidia are strongly reminiscent of Xylariaceous anamorphs and at least one species of *Hadrotrichum* is known to be xylariaceous (PETRINI & MÜLLER, 1986).

Periconia igniaria MASON et M.B. ELLIS

Description and illustrations: ELLIS (1971)

Collection examined: 64 Sauveterre, on *Phyllostachys* sp., FC 7.12.1986 (FC 154).

Pteroconium anam. *Apiospora camptospora* PENZ. et SACC.

Description and illustrations: ELLIS (1971)

Collection examined: 40 Tartas, on *Arundinaria japonica*, FC 25.12.1984 (ZT).

Torula herbarum (PERS.) LINK ex S.F. GRAY

Description and illustrations: ELLIS (1971)

Collection examined: 64 Assat, Maison Diriart, on *Phyllostachys mitis*, FC 11.03.84 (ZT).

Verticillium lecanii (ZIMM.) VIEGAS

Description and illustrations: GAMS (1971)

Collection examined: 30 Anduze, Bambouseraie de Prafrance, Generargues, on *Phyllostachys pubescens*, FC 21.11.87 (FC174).

Several collections from other bamboo species, same site, were examined but not kept.

2. COELOMYCETES

Amerosporium polynematoides SPEG.

Description and illustration: SUTTON (1980)

Collection examined: 30 Anduze, Bambouseraie de Prafrance, Generargues, on *Phyllostachys nigra*, FC 21.11.87 (FC174).

Dinemasprium strigosum (PERS.: FR.) SACC.

Description and illustrations: SUTTON (1980)

Collections examined: 64 Sauveterre de Béarn, on *Phyllostachys bambusoides*, FC 15.4.1984 (ZT); 30 Anduze, Bambouseraie de Prafrance, Generargues, on *P. pubescens*, FC 21.11.87 (ZT).

Epicoccum purpurascens EHRENB. ex SCHLECHT.

Description and illustrations: ELLIS (1971)

Collections examined: 64 Sauveterre de Béarn, on *Phyllostachys* sp., FC 20.4.1986; 40 Peyrehorade, on *Phyllostachys* sp., FC 29.11.1987.

ACKNOWLEDGEMENTS

We are deeply indebted to following individuals for their help in the identification of our collections:

J. BOISE, Farlow Herbarium, Cambridge, USA; S. CARPENTER, Forestry Sci. Lab., Corvallis, Oregon, USA; Y. DOI, Nat. Sci. Museum, Tokyo, Japan; K. KATUMOTO, Yamaguchi University, Yamaguchi, Japan; S. M. FRANCIS, CMI, Kew, UK; L. HOLM, Uppsala University, Uppsala, Sweden; W. GAMS, CBS, Baarn, NL; F. RAPPAZ, Lausanne, Switzerland; G.J. SAMUELS, NY Bot. Garden, Bronx, USA; M. SHERWOOD-PIKE, University of Oregon, Eugene, USA; B.M. SPOONER, Royal Bot. Garden, Kew, UK; H. ZOGG, Herbarium, ETH Zürich, Switzerland.

Dr. E. MÜLLER, Zürich, Switzerland, confirmed the identification of some collections, critically read the manuscript and made many helpful suggestions.

REFERENCES

- AEBI, B. (1972). Untersuchungen über Discomyceten aus der Gruppe *Tapesia-Trichobelonium*.- Nova Hedwigia 23: 49 - 112.
- AHMED , S.I. & R.F. CAIN (1972). Revision of the genera *Sporormia* and *Sporormiella*.- Can. J. Bot. 50: 419-477.
- AMES, L. M. (1963). A monograph of the Chaetomiaceae.- U.S. Army Res. Dev. Ser. 2, 125 pp.
- ARX, VON J.A. & E. MÜLLER (1954). Die Gattungen der amerosporen Pyrenomyceten.- Beitr. Kryptogamenfl. Schweiz 11(1): 1-434.
- ARX, VON J.A. (1959). Ein Beitrag zur Kenntnis der Fliegenfleckenpilze.- Proc. Kon. Ned. Ak. Wet. Amsterdam, ser. C, 62: 333-340.
- BARR, M. E. (1984). *Herpotrichia* and its segregates.- Mycotaxon 20(1): 1-34.
- BOIDIN, J., F. CANDOUSSAU & G. GILLES (1986). Bambusicolous fungi from the Southwest of France II. Saprobic Heterobasidiomycetes, resupinate Aphyllophorales and Nidulariales.- Trans. Mycol. Soc. Japan 27: 463-471.
- BOOTH, C. (1957). Studies of Pyrenomycetes: I. Four species of *Chaetosphaeria*, two with *Catenularia* conidia. II. *Melanopsamma pomiformis* and its *Stachybotrys* conidia.- Mycol. Papers 68: 1-27.
- BOOTH, C. (1971). The genus *Fusarium*.- C.M.I., Kew, 237 pp.
- CANDOUSSAU, F., K. KATUMOTO & M. SHERWOOD-PIKE (1985). Bambusicolous fungi from the Southwest of France I. Two new species of Pyrenomycetes and a new genus of the Phacidiaceae.- Sydowia 38: 28-34.
- CARPENTER, S.E. (1981). Monograph of *Crocicreas* (Ascomycetes, Helotiales, Leotiaceae).- Memoirs N.Y. Bot. Garden 33: 1-290.
- DENNIS, R.W.G. (1978). British Ascomycetes.- Cramer, Vaduz, 486 pp, 31 fig., 44 pl.
- DREYFUSS, M.M. (1975). Taxonomische Untersuchungen innerhalb der Gattung *Chaetomium* Kunze.- Sydowia 28: 50-133.
- ELLIS, M.B. (1971). Dematiaceous hyphomycetes.- C.M.I., Kew, Surrey, 608 pp.
- ELLIS, M.B. (1976). More dematiaceous hyphomycetes.- C.M.I., Kew, Surrey, 507 pp.

- FRANCIS, S.M. (1975). *Anthostomella* Sacc. Part I.- Mycological Papers 139: 1-97.
- GAMS, W. & V. HOLUBOVÀ-JECHOVÀ (1976). *Chloridium* and some other dematiaceous hyphomycetes growing on decaying wood.- Studies in Mycol. 13: 1-99.
- GAMS, W. (1971). *Cephalosporium*-artige Schimmelpilze.- Fischer, Stuttgart, 262 pp.
- HAFELLNER, J. (1979). *Karschia*.- Beihete Nova Hedwigia 62: 1-248.
- HAWKSWORTH, D.L. (1981). *Astrosphaeriella* Sydow, a misunderstood genus of melanommataceous pyrenomycetes.- Bot. J. Linn. Soc. 82: 35-59.
- HINO, I. & K. KATUMOTO (1955). Illustrationes fungorum bambusicolorum III.- Bull. Fac. Agric. Yamaguti Univ. 6: 29-68.
- HINO, I. & K. KATUMOTO (1958). Illustrationes fungorum bambusicolorum VI.- Bull. Fac. Agric. Yamaguti Univ. 9: 877-908.
- HOLM, L. & K. HOLM (1988). Studies in the Lophiostomataceae with emphasis on the Swedish species.- Symb. Bot. Upsal. 28(2): 1-50.
- ITURRIAGA, T. & R. KORF (1984). Studies in the genus *Strossmayeria* (Helotiales). 4. Connections to its anamorph, *Pseudospriopes*.- Mycotaxon 20(1): 179-184.
- KIRK, P. M. & B.M. SPOONER (1984). An account of the fungi of Arran, Gigha and Kintyre.- Kew Bull. 38(4): 503-597.
- KORF, R. (1981). A preliminary discomycete flora of Macaronesia: Part 2, Hyaloscrophaceae subf. Arachnopezizoideae.- Mycotaxon 13(1): 137-144.
- LE GAL, M. (1966). Contribution à la connaissance du genre *Scutellinia* (Cooke) Lamb. emend. Le Gal (1ère étude).- Bull. Soc. Mycol. Fr. 82: 301-333.
- MUNK, A. (1957). Danish Pyrenomyctes.- Dansk Bot. Arkiv 17: 1-491.
- MÜLLER, E. & J.A. VON ARX (1962). Die Gattungen der didymosporen Pyrenomyceten.- Beitr. Kryptogamenfl. Schweiz 11(2): 1-921.
- NAGRAJ, T.R. & B. KENDRICK (1975). A monograph of *Chalara* and allied genera.- Wilfred Laurier University, Waterloo, Ontario, Canada, 200 pp.
- NANNFELDT, J.A. (1985). *Niptera*, *Trichobelonium* und *Beloniopsis*, drei noch zu erläuternde Gattungen der mollisioiden Discomyceten.- Sydowia 38: 194-215.
- PETRINI, L.E. & E. MÜLLER (1986). Haupt- und Nebenfruchtformen europäischer *Hypoxyylon*-Arten (Xylariaceae, Sphaeriales) und verwandter Pilze.- Mycologia Helvetica 1: 501 - 627.
- PETRINI, L.E. & J.D. ROGERS (1986). A summary of the *Hypoxyylon serpens*-complex.- Mycotaxon 26: 401 - 436.
- POUZAR, Z. (1985). Reassessment of the *Hypoxyylon serpens*-complex II.- Ceska Mykologie 39: 129-134.
- RAPPAZ, F. (1987). Taxonomie et nomenclature des Diatrypacées à asques octosporés.- Mycologia Helvetica 2(3): 285-648.
- SACCARDO, P. (1882). *Sylloge fungorum*.- Padua, 768 pp.
- SAMUELS, G.J. (1988). Fungicolous, lichenicolous and myxomyceticolous species of *Hypocreopsis*, *Nectriopsis*, *Nectria*, *Peristomialis*, and *Trichonectria*. Memoirs N.Y. Bot. Garden 48: 1-78.
- SHERWOOD, M. (1977). The ostropalean fungi.- Mycotaxon 5: 1-277.
- SPOONER, B.M. & F. CANDOUSSAU (1988). Bambusicolous fungi from the Southwest of France III. A new species of *Encoelia*.- Trans. mycol. Soc. Japan 29: 219-223.
- SPOONER, B.M. & R.W.G. DENNIS (1985). New or interesting Ascomycetes from the Highlands and Islands.- Sydowia 38: 294-316.
- SUTTON, B.C. (1980). The coelomycetes. Fungi imperfecti with Pycnidia, Acervuli and Stromata.- C.M.I., Kew, Surrey, 696 pp.
- ZOGG, H. (1962). Die Hysteriaceae s. str. und Lophiaceae.- Beitr. Kryptogamenfl. Schweiz 11(3): 1-190.




Fig. 1.- a,b: *Hadrotrichum bambusicola*. a: conidiophores. b: conidia.-
c: ascospores of *Rosellinia* sp.




Fig. 2.- a-d: *Aulographum emiliae*. a: transverse section of an ascoma. b: ascomata on the host. c: ascus. d: ascospores.- e-g: *Schyzothyrium pomi*. e: ascomatal wall tissue. f: ascus. g: ascospores.

