Zeitschrift: Mycologia Helvetica

Herausgeber: Swiss Mycological Society

Band: 4 (1990-1991)

Heft: 2

Artikel: Pythium folliculosum, a new species from the bank of Lake Zürich

Autor: Paul, Bernard

DOI: https://doi.org/10.5169/seals-1036495

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Pythium folliculosum, a New Species from the Bank of Lake Zürich.

Bernard Paul

Institut de Biologie, Université d'Oran 311000 Es-senia, ORAN, Algérie

Summary: *Pythium folliculosum* sp. nov. has been isolated and identified from soil samples taken at the bank of lake Zürich. This fungus is characterized by its filamentous, inflated type of sporangia, together with both spherical, elongated saclike oogonia, and strictly monoclinous antheridia.

Résumé: Pythium folliculosum sp. nov. a été isolé à partir d'échantillons de sol prélevés au bord du lac de Zürich. Ce champignon est caractérisé par ses sporanges filamenteux enflés: ses oogones sphériques et allongées (en forme de sac) et par ses anthéridies exclusivement monoclinales.

Introduction

The genus *Pythium* comprises about 100 species which are mostly cosmopolitan in distribution. Most of these live in aquatic or other humid habitats. Many species of this genus occur as saprophytes in the soil, while others are parasites that cause a variety of diseases to plants. The most common disease caused by *Pythium* is the "damping-off" of young seedlings in the nurseries, green houses or in the field.

During the course of investigation of these fungi in Switzerland, a number of soil samples, specially forest soils, were collected from different parts of the country. Some humid soil samples were also collected from the banks of "lac Léman" and the lake of Zürich.

A new species of *Pythium* was discovered in one sample taken from the bank of lake Zürich at Zollikon. This fungus belongs to the group of *Pythium* with filamentous inflated sporangia and can easily be distinguished from related species by the presence of sac like oogonia.

Materials and Methods

In July 1989 soil samples were collected at the bank of lake Zürich in sterile capped, disposable plastic bottles. This was done with a spoon which was disinfected with alcohol after each sampling. The samples were brought to the

was transferred into a sterile capped bottle containing 20 ml of sterile distilled water. This was shaken vigorously to obtain a homogeneous soil suspension. Ten ml of this suspension were then placed in a sterile Petri dish containing 10 ml of sterile distilled water. This was, in turn, baited with boiled hemp-seed halves. After 3-4 days the bait was colonized by fungal mycelium. To supress the growth of other quick-growing soil fungi 0,5 grams of benomyl was dissolved in 1 litre of distilled water and autoclaved. This Benomyl water was then used to culture the fungus. Repeated subculturing on hemp seeds in sterile distilled water and benomyl water allowed to produce a bacteria-free colony of *Pythium*. Potato carrot agar (PCA) and corn meal agar (CMA) were used for the study of the isolates (Plaats-Niterink, 1981). Small pieces of fungal mycelium were innoculated on PCA and CMA plates, which were in turn incubated at different temperatures.

The isolates were identified with the help of keys provided by Middleton (1943), Waterhouse (1967) and Plaats-Niterink (1981). A comparative study of this isolate with the descriptions of all the recently discovered species of Pythium was also done (Ali-Shtayeh & Dick 1985, Paul 1986 a,b,c; 1987 a,b; 1988 a,b)

Pythium folliculosum sp. nov. Plates 1-2

Hyphae hyalinae, 4-7 μm diametro. Sporangia filamentosa dilatata. Oogonia terminalia vel intercalaria, globosa, prolata, folliculosa, 10-28 μm diam. et ad 70 μm elongatae. Antheridia monoclinata, cellulae antheridiales clavatae. Oogonia continentia unam, interdum duas, raro tres oosporas. Oosporae apleroticae et pleroticae, 6-24 μm in diametro, paries 0.8-3 μm crassa, globulosa. Temperatura optima 25°C. Isolatum ex terra in Zürich (Zollikon), confoederatione Helvetica. Holotypus in herbario universitatis Oranensis conservatus (S-24).

Etymology :The fungus is being named as *Pythium folliculosum* because of the presence of sac-like oogonia (folliculus = sac).

Mycelium is well branched, measuring 4-7 μ m in diameter. Sporangia are produced at 5-25°C. Formation of sporangial vesicles and liberation of zoospores were not observed either in water cultures or in any of the solid media. The sporangia are of the filamentous inflated type. Toruloid inflated elements line up to form an irregularly swollen contiguous structure with densely granular content.

Oogonia are spherical, pear-shaped to elongated, forming sac-like structures, terminal, intercalary, and sometimes catenulate; when terminal they are borne on oogonial stalks measuring 5-60 μ m in length. Spherical oogonia measure 1028 μ m in diameter (av. 19.5 μ m), when elongated they can be as long as 70 μ m (av. 32.7 μ m).

Paul: Pithyum folliculosum 205




Plate 1: *Pithyum folliculosum*. 1: Filamentous inflated sporangia; 2-5: terminal oogonia; 6-7: intercalary oogonia; 8 -11: oogonia with sessile antheridia; 12-19: oogonia with monoclinous stalked antheridia; 20-25: plerotic oospores; 26-29: aplerotic oospores; 30-38: elongated sac like oogonia containing one to three oospores.

Antheridia are strictly monoclinous, unbranched, stalked or sessile, usually one rarely two per oogonium. Antheridial cells are clavate, making a narrow or wide contact with the oogonia.

Oospores single, often two, rarely three per oogonium, mostly spherical, rarely slightly elongated, measuring 6-24 μ m in diameter (av. 16.7 μ m),and are both plerotic and aplerotic. The oospore wall measures 0.8-3 μ m in thickness.

Pythium folliculosum grows luxuriantly on PCA, CMA and on hemp seeds in water. On both solid media the colony of the fungus is submerged, giving a narrow chrysanthemal pattern on CMA and radiate pattern on PCA. The optimal temperature for its growth was 25°C at which the average daily growth of the fungus was 14.4 mm on CMA and 14 mm on PCA.

Results and Discussion

Pythium folliculosum grows well on hemp seeds in water and on solid media. It belongs to the group of Pythium having filamentously inflated type of sporangia. Plaats-Niterink (1981) in her monograph on the genus Pythium has described about 15 species belonging to this group. However, none have the sac-like oogonia typical of P. folliculosum measuring as long as 70 μm. Most of the species of Pythium have globose, spherical oogonia. Recently, Ali-Shtayeh & Dick (1985) described P. pachycaule from England. This species comes very close to P. folliculosum as it has also sac-like elongated oogonia. A close look to the different characteristics presented by these two fungi makes it clear that they are closely related but not the same. The main differences are underlined in table 1.

TABLE 1 : Differences between P. folliculosum and P. pachycaule

	P. pachycaule	P. folliculosum
Sporangia	non- inflated or	inflated with cont-
	slightly inflated	iguous elements.
Oogonia	24-34 μm in diam.	smaller oogonia 10-
	(av. 26.6)	28 μm (av. 19.5 μm).
Antheridia	mono- and diclinous,	strictly monoclin-
	1-3 per oogonium	ous, usually one per
		oogonium.
Oospores	aplerotic, 18-25 μm	both plerotic and
	in diam. (av. 22 μm).	aplerotic, 6-24 μm
		(av. 16.7μm).

Paul: Pithyum folliculosum 207




Plate 2: *Pithyum folliculosum*. 1-2: filamentous inflated sporangia; 3-6: oogonia and monoclinous antheridia; 7: elongated sac like oogonia; 8: plerotic oospores; 9: oogonia containing multiple oospores; 10-11: elongated oogonia containing aplerotic oospores.

Acknowledgements

The author wishes to thank Rev. father Jean-Louis Declais of Oran for his help in preparing the latin diagnosis and Martine and Pierre Juvet of Cote aux Fées for their help in the collection of soil samples in Switzerland.

Bibliography

- Ali-Shayeh, & M. Dick, 1985: Five new species of *Pythium* (Perenosporomycetidae). Botanical Journal of the Linnean Society 91: 297-317.
- Middleton, J. T., 1943: The taxonomy, host range and geographic distribution of the genus *Pythium*. Mem. Torrey Bot. Club 20: 1-171.
- Paul, B., 1986a: A new species of *Pythium* from Algerian waters. Hydrobiologia 131: 31-38.
- Paul, B., 1986b: An aquatic species *Pythium toruloides* sp. nov. from Algeria. Trans. Brit. Mycol. Soc. 86: 330-334.
- Paul, B., 1986c: A new non zoosporic species of *Pythium* from Algeria. Hydrobiologia 140: 223-235.
- Paul, B., 1987a: A new species of *Pythium* with filamentous sporangia from Algeria. Trans. Brit. Mycol. Soc. 89: 195-198.
- Paul, B., 1987b: A new species of *Pythium* with ornamented oogonia from Algeria. Mycologia 79: 797-802.
- Paul, B., 1988a : Une nouvelle varieté de *Pythium capillosum* dans les sols algériens. Cryptogamie-Mycologie 9: 35-42.
- Paul, B., 1988b : Une nouvelle espèce de *Pythium* isolée d'une saline de l'ouest algérien. Cryptogamie-Mycologie 9: 325-333.
- Waterhouse, G. M., 1967: Key to *Pythium* Pringsheim. Mycological papers 109: 1-15.