

Zeitschrift: Mycologia Helvetica

Herausgeber: Swiss Mycological Society

Band: 6 (1994)

Heft: 2

Artikel: Records of *Ramaria ochraceo-virens* (Jungh.) Donk and *Sphaerobolus stellatus* Pers. from the Arctic island Bjørnøya (Bear Island)

Autor: Skifte, Ola

DOI: <https://doi.org/10.5169/seals-1036342>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Records of *Ramaria ochraceo-virens* (Jungh.) Donk and *Sphaerobolus stellatus* Pers. from the Arctic island Bjørnøya (Bear Island)

Ola Skifte

Tromsø Museum, IMV, University of Tromsø, N-9006 Tromsø, Norway

Summary. In 1983 *Ramaria ochraceo-virens* and *Sphaerobolus stellatus* were observed on the Arctic island Bjørnøya. A review is given of the material which was collected and of the performance of the species on the island.

KEY WORDS: *Ramaria*, *Sphaerobolus*, Bjørnøya (Bear Island).

Introduction

During the expedition of the Tromsø Museum, IMV, to Bjørnøya in 1983 (Engelskjøn 1986, Skifte 1989, Skifte & Høiland 1993) S. Dunfjeld, T. Engelskjøn and the author discovered *Ramaria ochraceo-virens* and *Sphaerobolus stellatus* on the island. As these species have not previously been published from the archipelago of Svalbard, I will give a brief account on these collections.

Bjørnøya (Fig. 1) is situated about halfway between the northernmost coast of Norway and the southernmost tip of Spitsbergen. The position for the weather station (and Bjørnøy Radio) is 74°31' N 19°01' E (Steffensen 1982).

The southern and southeastern parts of Bjørnøya consist mostly of mountains, the highest, Miseryfjellet, is 536 m a.s.l. The northern and northwestern parts of the island are flat but rugged and stony with many small lakes (Steffensen 1982). Bedrock varies considerably. Sandstone areas are represented in different parts of the island., and in northeastern Bjørnøya deposits of coal are found in these Devonian strata. Limestone and lime-rich bedrock are also encountered in several areas, especially in southern parts of the island (Engelskjøn 1986).

Bjørnøya has a maritime middle Arctic climate (Elvebakk 1985) with low temperatures and humid conditions. A special feature of the island, as well as other Arctic islands, is fog in summer.

The first contribution to the flora of Bjørnøya was a collection brought back from the island in 1827 by the Norwegian geologist, B. M. Keilhau (Rønning 1959). Thanks to the publications by Rønning (1959), Engelskjøn (1986) and Engelskjøn & Schweitzer (1970), our knowledge about the vascular flora of

This contribution is dedicated to Prof. Dr. M. Moser (Innsbruck, Austria),
on occasion of his 70th birthday.




Fig. 1.—Map of Bjørnøya (Bear Island) showing the localities where the material has been collected.—1. Northeastern Bjørnøya (Bear Island): Tunheim: Locality for *Sphaerobolus stellatus* Pers. (OS 19/Bjø 83).—2. Southern Bjørnøya (Bear Island): Between the lake Ellasjøen and the valley Ymerdalen. Locality for *Ramaria ochraceo-virens* (Jungh.) Donk (OS 88/Bjø 83).

Bjørnøya today is rather good, both from the floristic as well as from the eco-geographical point of view.

From the mycological point of view, however, the situation is quite different. Except for several records of fungi made by B. M. Kilhau and the Danish botanist J. E. Vahl (published e.g. in *Botaniska Notiser* 1840), the Finnish mycologist P. A. Karsten ought to be considered as the first contributor to our knowledge of the macrofungi in the archipelago of Svalbard (Woldmar 1969). The list of species from Bjørnøya published by Karsten (1972) includes seven agarics, one Gasteromycete and one discomycete.

As a participant in the expeditions to Bjørnøya from Tromsø Museum in 1957 and 1983 (Rønning 1959; Engelskjøn 1986), the author had the opportunity to gather specimens of macrofungi on the island. From these collections material representing the genera *Russula* (Skifte 1989) and *Cortinarius* (*Dermocybe*) (Skifte & Høiland 1993) has been published previously.

The material mentioned in this paper is deposited at Tromsø Museum, IMV, University of Tromsø (TROM).

Description of the taxa

***Ramaria ochraceo-virens* (Jungh.) Donk 1933**
Rev. Niederl. Homobas. Aphyll. 2: 112.

During a study tour to Copenhagen in 1985, the material of this *Ramaria* from Bjørnøya has been identified by H. Knudsen and J. H. Petersen.

The basidiomes are compact and irregularly branched, about 20 mm high, and about 10 mm wide the top. The stems measure about 2,7 mm at the apex, and about 2 mm at the base. The colour at the top of the basidiomes is yellow-orange (Kornerup & Wanscher 1961: 5A7), but dull ochreous (4A5) elsewhere. The lower part of the stem is yellow-green, especially at the base, which often stains greenish. The actual base is tomentose due to white mycelium, which still can be observed on dry material.

Spores 6.5–8(8.5)×3.5–4.5 μm , rough or slightly verrucose, ovoid to egg-shaped or oblong, often clumped together as stated by Corner (1950). Basidia about 40 × 5 μm , clavate, narrow with 4 sterigmata. Hyphae with clamps on septa.

Material examined: Southern BJØRNØYA: Between the lake Ellasjøen and the valley Ymerdalen (see Fig. 1). In *Salix polaris* – community dominated by moss vegetation, about 150 m a.s.l. The specimens were growing in deep moss-carpet forming a fairy-ring with a diameter of 90 cm. 15 August 1983, S. Dunfjeld, T. Engelskjøn & O. Skifte (TROM: OS 88/Bjø 83).

This contribution is dedicated to Prof. Dr. M. Moser (Innsbruck, Austria),
on occasion of his 70th birthday.

In Europe this species has been reported both from humus-rich soil (Corner 1950, Breitenbach & Kränzlin 1986, Bon 1987) and on litter in coniferous woods (Breitenbach & Kränzlin 1986, Bon 1987).

***Sphaerobolus stellatus* Persoon 1801**
Syn. Fung. 115.

Fruitbodies 1–2 mm in diameter, globose to subglobose, weakly tomentose at first, becoming smooth, white to pale yellow. On the substratum immersed in a web of white mycelium.

Spores 8–9.5×5 µm, broadly ellipsoid, hyaline, smooth.

Material examined: Northeastern BJØRNØYA: Tunheim, in an old cow-stable constructed during the coal-mining time (1916–1925), in large numbers on a board of *Pinus sylvestris*, 10 August 1983, S. Dunfjeld, T. Engelskjøn & O. Skifte (TROM: OS 19/Bjø 83).

The distribution of *Sphaerobolus stellatus* seems to be very wide. Eckblad (1955) and Lange (1948) claim it is a cosmopolitan species. It seems, however, to be rare in (sub-)arctic areas of Fennoscandia. So far, only few records have been published from northern Finland (Kallio & Kankainen 1966) and Greenland (Lange 1948). According to Herbarium specimens in TROM, we know this species from northern Norway (Nordreisa and Lyngen districts in Tromsø county). *Sphaerobolus stellatus* has not previously been observed in the archipelago of Svalbard.

Discussion

Ramaria ochraceo-virens: The habitats supporting this species at Bjørnøya are quite different from those recorded elsewhere in Europe. Perhaps the species is not very substratum-dependent, and for its occurrence both climate and hydrology may be important as well. In Scandinavia this species is recorded from different ecological niches, e. g. on rotting needles of *Juniperus* growing on calcareous soil (Petersen & Vesterholt 1990).

At Bjørnøya *Ramaria ochraceo-virens* is growing in a habitat on limestone and dolomite (Engelskjøn 1986), dominated by several flowering plants demanding eutrophic soil, such as: *Equisetum scirpoides*, *Alopecurus alpinus*, *Poa alpina* var. *vivipara*, *Phippsia concinna*, *Salix polaris*, *S. reticulata*, *Ranunculus sulphureus*, *Papaver dahlianum*, *Arabis alpina*, *Draba alpina*, *Saxifraga hirculus*, *S. nivalis*, *S. oppositifolia*, and *S. tenuis*.

As regards to the climate, Elvebakk (1985) classifies Bjørnøya as middle Arctic and Engelskjøn (1986) considers the island as being situated in the typical maritime Arctic, having a longer growing season (extending into September) than Spitsbergen.

Accordingly, both the peculiar substrate and the rather mild climate on Bjørnøya may be responsible for the occurrence of *Ramaria ochraceo-virens* in an otherwise Arctic environment.

Sphaerobolus stellatus: During the coal-mining time a horse was kept at Tunheim (Hagnæs Kjelldahl 1973). After the coal-mining was given up (1925), this horse ("Bjørnøya Blakka") still lived there until 1941 as a mascot for the crew of Bjørnøy Radio which had been established in the Tunheim area. It is difficult to imagine how *Sphaerobolus stellatus* reached Bjørnøya. Perhaps it has been accidentally introduced with the wooden board on which we found it or more likely arrived on the island with fodder (grass, straw) for the before-mentioned horse, or other domestic animals.

Acknowledgements

I am grateful to Dr. H. Knudsen and his staff for identifying my material of *Ramaria* from Bjørnøya. For help during the preparation of this manuscript I am indebted to T. Engelskjøn, A. Granmo, and to O. Kvalheim for assistance with the illustration. The Norwegian Research Council for Science and Humanities covered the expenses for the expedition in 1983.

References

Bon, M. 1987. *Mushrooms and toadstools of Britain and north-western Europe*. London, 352 pp.

Breitenbach, J. & F. Kränzlin. 1986. *Pilze der Schweiz*. Vol. 2. *Heterobasidiomycetes, Aphylophorales, Gastromycetes*. Luzern, 416 pp.

Corner, E. J. H. 1950. *A monograph of Clavaria and allied genera*. London, 740 pp.

Eckblad, F.-E. 1955. *The Gasteromycetes of Norway*. *Nytt Mag. Bot.* 4, 19–86.

Elvebakk, A. 1985. Higher phytosociological syntaxa on Svalbard and their use in subdivisions of the Arctic. *Nord. J. Bot.*: 263–284.

Engelskjøn, T. 1986. Eco-geographical relations of the Bjørnøya vascular flora (Svalbard). *Polar Research* 5, 79–127.

Engelskjøn, T. & H. J. Schweitzer. 1970. Studies on the flora of Bear Island (Bjørnøya) I. Vascular plants. *Astarte* 3, 1–36.

Hagnæs Kjelldahl, D. 1973. *Bjørnøya. På utpost i ishavet*. Oslo, 143 pp.

Kallio, P. & E. Kankainen. 1966. Additions to the mycoflora of northernmost Finnish Lapland. *Annls. Univ. Turku Ser. A. II*, 36, 172–210.

Karsten, P. A. 1872. *Fungi in insulis Spetsbergen et Beeren Eiland collecti*. *Öfv. Kongl. Vetsk. Akad. Förh.* 2, 91–108.

Kornerup, A. & J. H. Wanscher. 1961. *Farver i Farver*. København, 248 pp.

This contribution is dedicated to Prof. Dr. M. Moser (Innsbruck, Austria),
on occasion of his 70th birthday.

Lange, M. 1948. The Gasteromycetes of Greenland. Medd. Grønl. 147(4), 1–32.

Petersen, J. H. & J. Vesterholt. 1990. Danske storsvampe (basidiesvampe). Copenhagen, 588 pp.

Rønning, O. I. 1959. The vascular flora of Bear Island. Acta Boreal. Ser. A. Scientia No. 15, 1–53.

Skifte, O. 1989. *Russula* of the island Bjørnøya (Bear Island), Svalbard. Opera Bot. 100, 233–239.

Skifte, O. & K. Høiland. 1993. *Cortinarius* subgen. *Dermocybe* on Bear Island. Blyttia 51, 151–153.

Steffensen, E. 1982. The climate at Norwegian Arctic stations. Klima 5, 1–44.

Survey of Bjørnøya (Bear Island) 1922–1931. Norges Svalbard og Ishavs Undersøkelser. Skrifter 86, 1–82.

Woldmar, S. 1969. Svalbards storsvampar i litteratur och natur. Friesia 9, 282–288.