

Zeitschrift: Mycologia Helvetica
Herausgeber: Swiss Mycological Society
Band: 10 (1998-1999)
Heft: 1

Artikel: Cloning and molecular biology of fungal allergens
Autor: Cramer, R. / Hemmann, S. / Blaser, K.
DOI: <https://doi.org/10.5169/seals-1036393>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Cloning and molecular biology of fungal allergens

Cramer, R., Hemmann, S. and K. Blaser

Swiss Institute of Allergy and Asthma Research (SIAF),
Obere Strasse 22, CH-7270 Davos, Switzerland

Endemic and opportunistic fungal infections have become an important medical problem, concomitantly with the increasing number of patients undergoing immunosuppressive forms of medical treatment and with increases in congenital and acquired immunodeficiency (Murphy et al. 1993). Morbidity and mortality rates associated with fungal species that are normally harmless to immunocompetent hosts, such as *Candida*, *Aspergillus*, *Fusarium* and *Trichosporum*, are high (Levitz, 1992) and diagnosis and treatment of these infections remain difficult. These fungal species are, together with *Cladosporium* and *Alternaria*, leading allergenic moulds (Karlsson-Borga et al., 1989). The ability to diagnose fungal infections and allergies, as well as to study immune responses to fungi has been so far hampered by the lack of well defined antigens (Levitz, 1992). The same reason also hinders the precise diagnosis of allergic bronchopulmonary aspergillosis (Cramer, 1996). Significant advances in our understanding of immune defence mechanisms to fungal infections and fungal sensitisation can be expected from the application of molecular biology leading to improved definition of fungal antigens/allergens. Molecular cloning and production of recombinant antigens/allergens provide modern methods to produce standardised reagents with a great potential to improve both, diagnosis of fungal complications (Cramer et al. 1996a) and basic research (Scheiner & Kraft, 1995). Classical cloning methods (Achatz et al. 1995) as well as the development of a new cDNA cloning system based on phage surface display technology (Cramer 1997) allow fast isolation and characterisation of cDNAs encoding fungal allergens. Most of the cloned allergens correspond to phylogenetically highly conserved proteins and include enolase, aldehyde dehydrogenase, alcohol dehydrogenase, P2 acidic ribosomal protein (Achatz et al. 1995) and manganese superoxide dismutase (MnSOD, Cramer et al. 1996b), proteins which occur in all eukaryotic organisms. The homologous proteins isolated from different fungal sources share a high degree of sequence identity and similarity when compared with each other and also to the corresponding human proteins (Achatz et al. 1995; Cramer et al. 1996b). In the case of MnSOD it has been shown that both, the fungal and the human enzyme are recognised by IgE antibodies from subjects allergic to the *A. fumigatus* protein and elicited specific immediate type allergic skin reactions in these individuals (Cramer et al. 1996b). Whether these autoimmune reactions are caused by

molecular mimicry between conserved T and B cell epitopes present on the fungal and on the structurally related human enzyme or by sensitisation to human MnSOD due to an inflammatory process, remains to be elucidated. Besides these allergenic proteins which can be clearly classified in terms of biochemical function, a number fungal cDNAs encoding IgE-binding proteins with unknown function have been reported (Cramer, 1996; De Vouge et al. 1996; Banerjee et al. 1996). None of the fungal allergens that have been sequenced so far correspond with respect to sequence and/or enzymatic function to previously sequenced non-fungal allergens, although many of these allergens do occur as homologous allergens in more than one fungal species. These results suggest that fungal sensitisation could be a phenomenon related to cross-reactive fungus-specific molecular structures among phylogenetically closely related organisms.

Acknowledgement

This work was supported in part by the Swiss National Science Foundation (grant 31-39429.93).

References

- Achatz, G., Oberkofler, H., Lechnauer, E., Simon, B., Unger, A., Kandler, D., Ebner, C., Prillinger, H., Kraft, D. & Breitenbach, M. 1995. Molecular cloning of major and minor allergens of *Alternaria alternata* and *Cladosporium herbarum*. *Mol. Immunol.* 32:213–227.
- Banerjee, B., Kurup, V. P., Phadnis, S., Greenberger, P. A. & Fink, J. N. 1996. Molecular cloning and expression of a recombinant *Aspergillus fumigatus* protein Asp f II with significant immunoglobulin E reactivity in allergic bronchopulmonary aspergillosis. *J. Lab. Clin. Med.* 127:253–262.
- Cramer, R. 1997. pJuFo: a phage surface display system for cloning genes based on protein-ligand interaction. In: B. Schaefer (ed). *Gene Cloning and Analysis: Current Innovations*. Horizon Scientific Press, Wymondham, U.K. 29–42.
- Cramer, R. 1996. Recombinant *Aspergillus fumigatus* allergens: from the nucleotide sequences to clinical applications. *Habilitationsschrift Universität Salzburg*.
- Cramer, R., Blaser, K. & Menz, G. 1996a. Rekombinante Allergene und ihr Potential für die allergologische Diagnostik. *Pneumologie* 50:387–393

- Crameri, R., Faith, A., Hemmann, S., Jaussi, R., Ismail, C., Menz, G. & Blaser, K. 1996b. Humoral and cell-mediated autoimmunity in allergy to *Aspergillus fumigatus*. J. Exp. Med. 184:265–270.
- De Vouge, M.W., Thaker, A. J., Curran, I.H. A., Zahng, L., Muradia, G., Rode, H. & Vijay, H. M. 1996. Isolation and expression of a cDNA clone encoding an *Alternaria alternata* Alt a 1 subunit. Int. Arch. Allergy Immunol. 111:385–395.
- Karlsson-Borga, A., Jonsson, P. & Rolfsen, W. 1989. Specific IgE antibodies to 16 wide-spread mould genera in patients with suspected mould allergy. Ann. Allergy 63:521–526.
- Levitz, S. M. 1992. Overview of host defences in fungal infections. Clin. Infect. Dis. 14: 37–42.
- Murphy, J. W., Friedman, H. & Bendinelli, M. 1993. Fungal infections and immune responses. Plenum Press, New York.
- Scheiner, O. & Kraft, D. 1995. Basic and practical aspects of recombinant allergens. Allergy 50:384–391.

