

Zeitschrift: Bulletin der Schweizerischen Akademie der Medizinischen Wissenschaften = Bulletin de l'Académie suisse des sciences médicales = Bollettino dell' Accademia svizzera delle scienze mediche

Herausgeber: Schweizerische Akademie der Medizinischen Wissenschaften

Band: 8 (1952)

Heft: 1-2: Symposium über die Beeinflussung des reaktiven Geschehens durch Hypophyse und Nebennierenrinde = Symposium on the influence of the hypophysis and the adrenal cortex on biological reactions = Symposium sur l'influence de l'hypophyse et de la corticostéroïde dans les réactions biologiques

Artikel: Influence of ACTH and cortisone on the antibody production and the plasma cell reaction

Autor: Moeschlin, S. / Báguena, R. / Báguena, J.

DOI: <https://doi.org/10.5169/seals-307070>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Medical Department, University of Zurich – Director Prof. W. Löffler

Influence of ACTH and Cortisone on the Antibody Production and the Plasma Cell Reaction

By S. Moeschlin, R. Báguena and J. Báguena

In four series of rabbits, the influence of ACTH and cortisone on the antibody production was studied.

All the animals were sensitized with a subcutaneous injection of 3 cm³ of typhoid-paratyphoid-vaccine. Approximately 3 weeks later, when a positive agglutination titer could be demonstrated in the blood, an intravenous dose (2.5 cm³ of vaccine) was administered. The antibody titer was studied every day, the animals were killed 3, 4, 5, 6 and 7 days after the revaccination and the spleen, the bone marrow and the liver examined by smears, histologically and especially the development of the plasma cells by the phase contrast microscope.

In the first series of 20 rabbits, the investigation of *Fagraeus* about the relation between the rise of the antibody titer and the plasma cells was confirmed. Moreover, by phase contrast microscopy the appearance of drop-like, dark granules in these cells could be demonstrated which correspond to the same granulations as shown by *Moeschlin* in myeloma cells producing pathological β - or γ -globulins.

In the second series of 6 rabbits, 3 received 20 mg of ACTH simultaneously with the revaccination and later every day until they were killed. We saw no difference between the antibody titer and the plasma cell number in the spleen of these animals and the control rabbits. Cortisone in a third series administered in the same way (8 rabbits) gave the same result.

Thus ACTH and cortisone, if given simultaneously with the reinjection of vaccine and on the following days, did not change the antibody formation or the plasma cell reaction. In the third series ACTH and cortisone were started 7 days before the reinjection and produced a marked decrease of the plasma cell reaction and the antibody production. Probably the difference in response is rather due to a change in the break-

down of the antigen, which in some way may be inhibited by the preliminary treatment with ACTH and cortisone, than to a depression of the antibody formation itself.

Bibliography: see *Moeschlin et al.*: *Acta Haematol.* **6** (1951), and *Proc. of Internat. Congress of Allergy, Zurich 1951.*