

Zeitschrift: Bulletin der Schweizerischen Akademie der Medizinischen Wissenschaften = Bulletin de l'Académie suisse des sciences médicales = Bollettino dell' Accademia svizzera delle scienze mediche

Herausgeber: Schweizerische Akademie der Medizinischen Wissenschaften

Band: 13 (1957)

Heft: 1-4: Symposium über Arteriosklerose = Symposium sur l'artérosclérose
= Symposium on arteriosclerosis

Artikel: Nutritional factors in experimental atherosclerosis of the rabbit

Autor: Sala, G. / Di Marco, A.

DOI: <https://doi.org/10.5169/seals-307326>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Research Laboratories, Farmitalia, Milan

Nutritional Factors in Experimental Atherosclerosis of the Rabbit

By G. Sala and A. di Marco

Studies carried out on experimental cholesterol atherosclerosis in the rabbit showed the presence of different degrees of aortic atherosclerotic lesions in groups of animals fed with different diets.

This paper deals with the investigation of the role of nutritional factors and with the importance of endogenous factors in the development of experimental atherosclerosis. The complete work has been published elsewhere (1).

Material and methods

The detailed methods used in the present study are described elsewhere (1, 2).

Atherosclerosis was obtained in the male rabbit after the daily oral administration of 1 g of cholesterol and 3 ml of olive oil, mixed into the diet, for 60 days.

The atherosclerotic process was evaluated both by a subjective examination of the aorta and by chemical determination of cholesterol in the ether extract of the adventitia free aorta.

Serum cholesterol, total fats, phospholipids, turbidity and lipoproteins have been determined according to the methods already described. Thyroid function has been determined by histological methods and by the thyroid uptake of a tracer dose of I¹³¹.

Results

1. A significant difference of atherosclerosis is evident in groups of rabbits fed different diets (Fig. 1). The lower degree of atherosclerosis meets with a decreased body weight and a lower concentration of fats in the serum. Thyroid function is higher in animals with inhibited atherosclerosis than in controls presenting a high degree of atheromasic process (Fig. 2).

2. The crude composition of the two diets is different (Table 1), but the




Fig. 1. Atherosclerotic pattern in two groups of rabbits fed different diets.

	I Diet A	II Diet B
Fixation of I^{131} thyroid %	9 ± 0.8	35 ± 8
	20 ± 2.2	32 ± 5.6
Weight of the thyroid gland mg/kg	60 ± 23	184 ± 31
Follicle diameter μ	50 ± 3	43 ± 4
Histological aspect	calm	activated

Fig. 2. Relationship between atherosclerosis and thyroid function.

chemical analysis does not show marked differences in carbohydrates, cellulose, proteins, fats, unsaponifiable matter and moisture content; the caloric value of the non-atherogenic diet (diet B) is eventually higher (Table 2). Diet A, however, is enriched with a salt mixture, vitamins and antibiotics.

3. When diet A is lacking in antibiotics or vitamins, the atheromasic process is not prevented; likewise the addition of single vitamines to the same diet appears to have no effect on atherosclerosis (Fig. 3).

4. In preliminary observations we could establish that rabbits fed diet B with low incidence of atherosclerosis, ingested a reduced quantity of food. Since this finding could be connected with atherosclerosis inhibition, it seemed necessary to consider separately cholesterol + olive oil and total food ingestion.

Table 1
Crude composition of diet A and B

Diet A	%	Diet B	%
Corn	26	Bran	50
Wheat meal	20	Corn	20
Alfa-alfa	22	Oats	20
Bran	30	Dry bread	10
Salt mixture*	2	Carrots	g 100
Vitamins**		Salad	g 50
Antibiotics***			

* Salt mixture = $\text{CaO} = 46.8$; $\text{CO}_2 = 27.9$; $\text{Na} = 5.6$; $\text{Cl} = 8.7$; $\text{P}_2\text{O}_5 = 9.4$; moisture = 1.2%.

** Vitamins = $\text{B}_1 = 2810 \text{ U/kg}$; $\text{B}_2 = 3435 \gamma/\text{kg}$, $\text{A} = 6600 \text{ U/kg}$; $\text{D}_3 = 235 \text{ U/kg}$.

*** Antibiotics = Bacitracin = 1.73 γ/kg ; Procaine Penicillin = 40 mg/kg.

Table 2
Chemical analysis and caloric equivalent of diet A and B

	Diet A %	Diet B %
Total carbohydrates	46	54.60
Cellulose	5.75	4.20
Proteins	15	13
Fats	3.03	4.50
Unsaponifiable matter	0.57	0.43
Total phosphorus	0.60	0.80
Ash	5.78	4
Moisture	12.60	11.80
Caloric equivalent	3800 Cal/kg	4130 Cal/kg

a) If groups of rabbits are obliged to eat amounts of cholesterol + olive oil equivalent to those of controls but still reduced quantity of food, an inhibition of atherosclerosis occurs, in spite of serum cholesterol and phospholipids being increased (Fig. 4).

b) An equivalent intake of food and of cholesterol + olive oil was obtained in both groups after the addition of a salt mixture to the non-atherogenic diet (diet B): finally the gravity and the incidence of atherosclerosis was the same in the groups fed different diets (Fig. 5).

Comment

From these results, it appears that atherosclerosis is inhibited in rabbits which were fed a lower amount of food and which consequently received a reduced caloric intake. These animals also showed an increased thyroid function, whose origin is still difficult to establish.




Fig. 3. Vitamins and antibiotics do not modify the atherogenic effect of diet A.




Fig. 4. Increased ingestion of cholesterol + olive oil does not favour atherosclerosis, when the food intake is low.




Fig. 5. High food intake is necessary for the development of atherosclerosis.

The increase of caloric intake seems to be responsible for the development of atherosclerosis, while the increased cholesterol + olive oil ingestion was not able per se, in absence of enough food intake, to favour the atherosclerotic process, in spite of the increased levels of serum fats.

In conclusion, these findings suggest that cholesterol experimental atherosclerosis in the rabbit depends both upon the exogenous cholesterol ingestion and upon metabolic and endogenous factors: prominent among them are the total caloric intake and thyroid function.

Summary

From the study of experimental atherosclerosis in groups of rabbits fed on different diets, the authors think that cholesterol atherosclerosis is determined both by exogenous cholesterol intake and by nutritional factors: low total caloric intake and increased thyroid function may inhibit the atheromastic process in the rabbit, even in presence of a high cholesterol intake.

Zusammenfassung

Auf Grund von Versuchen mit Kaninchen, bei denen durch verschiedenartige Fütterung eine experimentelle Atherosklerose erzeugt worden war, glauben die Verfasser, daß die Cholesterinatherosklerose sowohl durch exogene Cholesterinaufnahme als auch durch nahrungsbedingte Faktoren verursacht wird. Niedrige Kalorienzufuhr und steigende Schilddrüsenfunktion können den atheromatösen Prozeß beim Kaninchen hemmen, selbst bei gleichzeitiger reichlicher Cholesterinzufuhr.

Résumé

Se fondant sur une étude sur l'athérosclérose expérimentale chez des lapins soumis à des régimes différents, les auteurs pensent que l'athérosclérose cholestérolique dépend, d'une part, de l'apport exogène du cholestérol, d'autre part, de facteurs nutritifs; une ration alimentaire hypocalorique associée à une hyperfonction thyroïdienne peut inhiber l'athérosclérose, même si l'alimentation est riche en cholestérol.

1. *Sala, G., and di Marco, A.: Aterosclerosi umana e sperimentale.* CEA, Milano 1956, p. 189. – 2. *Sala, G.: Minerva med. (Torino) 2, 251 (1956).*