Zeitschrift: Schweizer Archiv für Tierheilkunde SAT : die Fachzeitschrift für

Tierärztinnen und Tierärzte = Archives Suisses de Médecine Vétérinaire

ASMV : la revue professionnelle des vétérinaires

Herausgeber: Gesellschaft Schweizer Tierärztinnen und Tierärzte

Band: 132 (1990)

Heft: 8

Artikel: Comparison of lesions in one- and ten-day-old gnotobiotic calves

inoculated with a rotavirus of low virulence

Autor: Hall, G.A. / Bridger, J.C. / Parsons, K.R.

DOI: https://doi.org/10.5169/seals-593503

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 16.11.2025


ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Table 1: Cause of death

Neoplasia	15	
Salpingo-peritonitis	12	
Chronic hepatopathy	11	
Cardiac failure	5	
Uricosis	2	
Marek's disease	1	
Coligranulomatosis	1	
Tuberculosis	2	
Aspergillosis	- 1	
Intestinal volvulus	1	

Table 2: Percent	tage of lesions	s of the different	apparatuses

8	TT
Digestive tract	93%
Cadiovascular system	84%
Urogenital system	82%
Endocrine and Lymphatic system	45%
Musculo-skeletal system	32%
Respiratory tract	24%






Fig. 2: Liver: Hepatoma

animals showed lesions of the cardiovascular system and 51% of the birds inspected were affected by atherosclerosis, mainly localized to the abdominal part of the aorta. The urogenital system showed an high degree of changes (82%), but most of these consisted of atrophy of the ovaries an oviducts, which could be considered as almost physiological changes due to the aging. Differently from what has been well established in poultry raised under intensive conditions, the pathology of the respiratory tract does not seem to play a major role in aged poultry (24%).

One of the most interesting observations standing out from this study is represented by the high rate of the non viral tumours (22%), which shows a considerable discrepancy with the values reported by diffe-

rent authors about the percentage of neoplastic lesions observed in broilers at slaughtering (.08 to 3%) (Hemsley, 1966; Bergmann et al., 1984)

Therefore our data show a direct correlation between aging and the development of non viral tumours in poultry, a phenomenon otherwise well established in mammals.

References

1. Andrew W. (1971): Heinemann Medical Books Limited, London. — 2. Bergmann V. et al. (1984): Mh. Vet.-Med. 39, 82. — 3. Hemsley L. A. (1966): J. Path. Bact. 92, 91.

AFRC Inst. Animal Health, Compton Laboratory, Berkshire (UK)

COMPARISON OF LESIONS IN ONE- AND TEN-DAY-OLD GNOTOBIOTIC CALVES INOCULATED WITH A ROTAVIRUS OF LOW VIRULENCE

G. A. Hall, J. C. Bridger, K. R. Parsons, K. C. Varshney
Calves usually experience infection with rotavirus in the first few
weeks of life, but many of these infections are subclinical (McNulty
& Logan, 1983). Fully virulent bovine rotaviruses are pathogenic in
calves of any age, whereas strains of low virulence only cause diarrhoea in day-old calves (Bridger, 1988). Explanations for this variation
in virulence have been sought by studying pathogenesis in one-dayold and ten-day-old calves inoculated with a strain of low virulence.

Material and methods

Three, one-day-old gnotobiotic calves were inoculated orally with 10^6 TCID₅₀ of a cloned rotavirus of low virulence (strain C3-160) and three were inoculated similarly at ten-days-old. Two groups of three calves were inoculated with virus-free medium and were age-matched controls. Calves were examined at least twice daily for clinical

signs of disease and faeces were collected and the virus content assayed by inoculation of MA-104 cells. Calves were killed within 24 hours of peak virus excretion. They were inoculated with colchicine prior to induction of terminal anaesthesia and small intestinal tissue was taken for measurement of villus height and crypt cell production rate (MacDonald & Ferguson, 1978; Hall et al., 1988a), for histological examination (Hall et al., 1988b) and for immunoperoxidase detection of rotavirus antigen (Parsons et al., 1984). The area of immunostaining in the epithelium was measured with an image analyser and expressed as the mean percentage of total epithelial area.

Results

Control calves and ten-day-old calves inoculated with rotavirus C3-160 remained healthy. Calves inoculated at one-day-old became depressed and developed explosive watery diarrhoea approximately




Fig. 1: Mean percent area immunostained for rotavirus antigen in the epithelium of the abomasum, duodenum and 9 small intestinal sites of one-day-old (*) and ten-day-old (o) gnotobiotic calves.

24 hours after inoculation, when the peak titre of virus in faeces $(10^{8.8}-10^{9.2})$ occurred. The peak titre of virus in faeces of calves inoculated at ten-days-old $(10^{5.7}-10^{7.9})$ was detected between 48 and 72 hours after inoculation.

The mean percent area of epithelium immunostained in the small intestine was greater in calves inoculated at one-day-old than in the calves inoculated at ten-days-old; immunostaining was present throughout the small intestine in one-day-old calves but absent from the upper small intestine of calves inoculated at ten-days-old (Fig.). In the calves inoculated at one-day-old, enterocytes were exfoliated and villi stunted and fused and enterocytes remaining in the epithelium were cuboidal. Villus height was reduced to 50% of control

values in the mid and distal small intestine. Foci of enterocyte exfoliation and cuboidal enterocytes in the epithelium were seen in calves inoculated at ten-days-old, but villus height was not affected. Crypt cell production rates were increased in both groups of calves, compared with controls, in the proximal, mid and distal small intestine.

Discussion

This study explains why the outcome of rotavirus infection differs in calves of different ages. In calves of both ages, the rotavirus used here infected and killed enterocytes but the inability of this strain to cause diarrhoea in ten-day-old calves appeared to be related to its inability to cause severe intestinal damage; it was not able to kill enterocytes faster than they could be replaced. There were differences in susceptibility of the gut to damage by this rotavirus in the two age-groups of calves; these host differences require further investigation.

References

1. Bridger JC. (1988): J Med Micro 27, xi-xii. — 2. Hall GA. et al. (1988a): Proc 15th World Buiatrics Congress, Palma de Mallorca. 915–920. — 3. Hall GA. et al. (1988b): Res Vet Sci 45, 240–250. — 4. MacDonald TT., Ferguson A. (1978): Gastroenterology 74, 496–500. — 5. McNulty MS., Logan EF. (1983): Vet Rec 113, 333–335. — 6. Parsons KR. et al. (1984): J Clin Pathol 37, 645–650.

Institut für Zoologie und Hydrobiologie, Universität München, FRG

LIGHT- AND ELECTRONMICROSCOPICAL EFFECTS OF ATRAZINE IN RAINBOW TROUT (ONCORHYNCHUS MYKISS) AFTER SUBCHRONIC EXPOSITION

R. Hoffmann, Th. Fischer-Scherl, A. Veeser

Atrazine has become one of the most frequently used ingredients of herbicides in the last decade. It enters the aquatic environment by run-offs from agriculturally used areas. Morphological investigations of fish with atrazine have primarily been performed with very high concentrations. These high concentrations, however, normally occur in the environment only after accidents for a short period. To achieve more information on this water pollutant and its impacts on fish in sublethal concentrations, rainbow trout (*Oncorhynchus mykiss*) were used to assess the effects of various atrazine concentrations.

Atrazine was dissolved to a saturated mixture, filtered and added constantly to aquaria with a capacity of $36\,l$ by mixing the original solution with tapwater by means of a dispenser. Atrazine concentration was measured regularly by a gaschromatograph at the beginning, during and at the end of the experiment to ensure the constant test concentration. Water temperature was $10.5^{\circ}C$ at a flow rate of 2 1/h, pH 7.6 and conductivity was $744~\mu$ S/cm. *Chronic exposure* to atrazine was performed at concentrations of $5,\,10,\,20,\,40$ and $80~\mu$ g/l for 28~days and *acute exposure* was done at concentrations of 1.4~and~2.8~mg/l for 96~h. For each concentration two tests with 10~fish each were performed simultaneously. Additionally, for each test two control

groups each with ten fish were examined. The mostly affected organs were gills and kidney.

Chronic exposure resulted in gills in hypertrophia as well as hyperplasia of chlorid cells. A low percentage of them had first signs of degeneration in lower concentrations (5–20 μ g/l) whereas in higher concentrations (40–80 μ g/l) degeneration was more prominent than hyperplasia. Acute exposition induced an immense proliferation of chlorid cells which were degenerated in a high percentage.

In kidney both the excretory as well as the hemopoietic parts were affected. At chronic exposure, podocyte proliferation, mesangial proliferation and thickening of glomerular basement membrane took place indicating a membrano-proliferative glomerulonephritis. Furthermore, a periglomerular fibrosis was seen in higher concentrations. Acute exposure induced an obliteration of Bowman's space due to proliferated mesangial cells.

The tubular system was especially affected at acute exposure showing extensive vacuolar degeneration.

In the hemopoietic tissue an increase of mitotic figures was the result of low concentrations. Higher concentrations at chronic exposure as well as acute intoxication resulted in a degeneration of sinus endothelia and hemopoietic cells.