Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =

Bulletin de la Société Entomologique Suisse = Journal of the Swiss

Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 44 (1971)

Heft: 1-2

Artikel: Stereospecific chemical synthesis of juvenile hormones

Autor: Corey, E.J.

DOI: https://doi.org/10.5169/seals-401640

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 02.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Stereospecific chemical synthesis of juvenile hormones

E. J. Corey

The major objectives of our research on the synthesis of juvenile hormones have been twofold: first, the development of stereospecific synthetic routes which would yield the pure hormones uncontaminated by isomeric structures, and secondly, the discovery of new methods for the stereospecific formation of the trisubstituted olefinic units which represent the major obstacle to the successful synthesis of the juvenile hormones and other acyclic isoprenoids. Such synthetic methods would fill a void which has been conspicuously displayed in those synthetic studies in the isoprenoid area which have relied on conventional processes of olefin formation. This lecture will be concerned, therefore, both with new synthetic reactions applicable to the stereospecific synthesis of acyclic isoprenoids and with the application of these methods to juvenile hormone synthesis in particular. Whatever success we have achieved in this enterprise is largely to the credit of the very able collaborators with whom I have been associated. In particular, the outstanding contributions of Hisashi Yamamoto, John Katzenellenbogen, Norman Gilman, and Steven Roman must be mentioned.

Attention will first be centered on several new processes for the stereospecific synthesis of trisubstituted olefins which have been developed at Harvard since 1966. These methods allow access to the four different arrangements which are possible about a carbon-carbon double bond with two different alkyl (or functionalized alkyl) groups,

a methyl group and a hydrogen atom as substituents.

Figure 1 shows a route which depends on the stereospecific generation of a vinyl aluminum intermediate by the action of lithium aluminum hydride on a propargylic alcohol. The underlying rationale for the investigation of this method arose from a number of previous findings including (1) the well-known unique lability of $C \equiv C$ to reduction by LiAlH₄ in propargylic alcohols, (2) the stereospecific formation of trans allylic alcohols of type RCH = CHCH₂OH from propargylic alcohols and LiAlH₄, and (3) the demonstration by Hochstein and Brown of an organo aluminum intermediate in the reduction of cinnamyl alcohol by LiAlH₄ (1). It was found by experimentation that

NEW METHODOLOGY APPLICABLE TO STEREOSPECIFIC SYNTHESIS OF ACYCLIC ISOPRENOIDS

 $\underline{e} \cdot \underline{g} \cdot , R = geranyl, n-hexyl$

Fig. 1

NEW METHODOLOGY APPLICABLE TO STEREOSPECIFIC SYNTHESIS OF ISOPRENOIDS

 $\underline{e} \cdot \underline{g}$., R = n-hexyl, π -tricyclyl Fig. 1a

the reaction of propargylic alcohols with LiAlH4 in tetrahydrofuran (THF) followed by iodination afforded a mixture of 3- and 2-iodinated alcohols in proportion which varied with the history and nature of the hydride reagent. It was further ascertained that the addition of sodium methoxide to the reagent led to exclusive formation of 3-iodo alcohol, whereas the addition of small amounts of aluminum chloride greatly favored formation of the 2-iodo alcohol (2). In connection with a program of research on carbon-carbon bond formation between unlike groups using organo-transition metal reagents (3), an excellent new process was at hand for the stereospecific replacement of iodo by methyl (or other alkyl) leading to the overall route to trisubstituted olefins shown in Figure 1. A method was also developed (4) for the oxidative one-flask conversion of primary allylic alcohols to methyl esters by the scheme shown in Figure 1 which proceeds via aldehyde, cyanohydrin, and acyl cyanide intermediates with manganese dioxide serving as the effective oxidizing agent. This method is efficient and stereospecific and obviously well suited to the synthesis of juvenile hormones.

Figure 1a demonstrates a highly efficient method for the conversion of a propargylic alcohol into a 2-iodo-trans-allylic alcohol using the reaction of diisobutylaluminum hydride on the lithium alkoxide (5). This alternative to the LiAlH₄-AlCl₃ reagent leads to both stereospecific and position specific reduction and contrasts to the cis course of reduction observed for the reaction of dialkylaluminum hydrides with acetylenic hydrocarbons. As shown in Figure 1a, this approach coupled with the use of either organocopper (3) or organonickel (5, 6) reagents allows the stereospecific synthesis of 2-methyl cis- or trans-

allylic alcohols.

Figure 2 shows still another new approach to trisubstituted olefins which has been developed at Harvard (7). This method, mentioned earlier in Dr. Siddall's lecture, was also studied by him at Syntex Corp. Figure 3 shows the synthesis of a 4,5-dehydrojuvenile hormone which rests partly on the methodology just discussed. The final product has been found to possess high juvenile hormone activity in studies by Prof. Carroll Williams. Interestingly, he observed even higher activity than that of C₁₈-cecropia hormone in certain bugs. The reduction of this dehydro compound to the C₁₈-cecropia juvenile hormone did not proceed cleanly even with the most favorable reagent studied (HN = NH), and so this sequence, though completely stereospecific, does not provide an efficient route to the natural hormone.

Figure 4 shows how the intermediate C₁₂-dienol, a synthesis of which appears in Figure 3, was converted to the C₁₈-cecropia hormone in the first stereospecific synthetic route to this substance (8). The approach demonstrates the applicability of the methods outlined earlier

in the lecture.

Figures 5 (9) and 6 (10, 11) outline two additional syntheses of the C₁₂-dienol intermediate which involve novel chemistry.

NEW METHODOLOGY APPLICABLE TO STEREOSPECIFIC SYNTHESIS OF ISOPRENOIDS

R = geranyl, n-hexyl

Fig. 2

A DEHYDRO CECROPIA HORMONE

Fig. 3

$\underline{\text{FIRST}} \ \underline{\text{SYNTHESIS}} \ \underline{\text{OF}} \ \underline{\text{C}}_{18} \underline{\text{-CECROPIA}} \ \underline{\text{HORMONE}}$

1.
$$PBr_3$$

2. $LiCH_2C = CSi(CH_3)_2$
3. Ag^+ then CN^-
4. $BuLi$ then CH_2O
from 2, $X = Si(CH_3)_3$
from 3, $X = H$
from 4, $X = CH_2OH$

from 2, R = Ifrom 3, $R = CH_3$

50%

Fig. 4

$\underline{\mathtt{ALTERNATE}} \ \underline{\mathtt{SYNTHESIS}} \ \underline{\mathtt{OF}} \ \underline{\mathtt{C}}_{12} \ \underline{\mathtt{DIENOL}}$

Fig. 5

ANOTHER SYNTHESIS OF C12-DIENOL

The latest synthesis of juvenile hormones to be completed at Harvard is stereospecific, simple, and direct. It depends heavily on the application of β -oxido phosphonium ylides to olefin synthesis as recently developed by us (12). Figure 7 outlines the new method in general form. It is apparent that the method allows the joining of three structural units in one synthetic step. The reaction is remarkable in that an

STEREOSPECIFIC SYNTHESIS OF TRISUBSTITUTED OLEFINS FROM β -OXIDO PHOSPHONIUM YLIDES AND ALDEHYDES

RCHO +
$$(C_6H_5)_3P$$
=CHCH₃

THF

RCHCH-P(C₆H₅)₃

RCHCH-P(C₆H₅)₃

I

RCHCH-P(C₆H₅)₃

I

RCHCH-P(C₆H₅)₃

I

RCHCH-P(C₆H₅)₃

I

RCHCH-P(C₆H₅)₃

I

RCHCH-P(C₆H₅)₃

I

RCHCHO

RCHCH-P(C₆H₅)₃

I

RCHCHO

RCHCH-P(C₆H₅)₃

II

R'CHO

CH₃

R' = alkyl or R | C=C CH₂OH | CH₃

R' = H

V

Fig. 7

intermediate containing three new stereocenters is generated stereospecifically and, further, a directional specificity operates in the final olefin-forming elimination.

Figure 8 shows the application of the oxido ylide method to the stereospecific synthesis of 5-methyl-cis-4-hepten-1-ol, a valuable intermediate for juvenile hormone synthesis (13). Figure 9 shows still another stereospecific route to the same alcohol (after borohydride reduction)

OXIDO YLIDE SYNTHESIS OF C8 ALCOHOL

OTHP

Collins

CHO

OTHP

1.
$$(C_6H_5)_3$$

PCHCH₃

2. BuLi
3. CH_2O

1. $OH \rightarrow Br$

2. Me_3

Feli

OTHP

CH₂OH

OTHP

NEW SYNTHESIS OF C7 AND C8 ALDEHYDES BY REARRANGEMENT

involving a 1,5-hydrogen shift of a *cis*-vinylcyclopropylcarbinol (13). This process is based on earlier findings of several groups (14–17) on the thermal decomposition of simple *cis*-2-alkylvinylcyclopropanes. The use of one of the intermediates for the efficient synthesis of 4-methyl-*cis*-3-hexenal (or -ol) is also outlined in Figure 9 (18).

Next, an application of the 3-component oxido ylide method to the stereospecific synthesis of juvenile hormones is outlined in Figure 10 (19). A single intermediate, generated in one step as shown, leads by a simple and short route to both C_{17} - and C_{18} -cecropia hormones. This is certainly the simplest synthesis of these substances now known which is stereospecific and leads directly to the pure substances. The method has also been used to generate the methyl-ethyl transposed position isomer of the C_{17} -hormone and the interesting chloro isostere of the C_{18} -hormone which are shown in Figure 10.

β-OXIDO YLIDE SYNTHESIS OF CECROPIA HORMONES

Finally, it seems appropriate to amplify the results reported during an earlier discussion at this meeting with regard to the biological properties of certain imino analogs of the juvenile hormones, for example, structure I in which R may be CH₃ or C₂H₅. These substances per se have only very low juvenile hormone type activity. However, Mr. Alfred Ajami, working with Prof. Riddiford at Harvard, has shown that the injection into cecropia pupae of a mixture of C₁₈-cecropia hormone

in small amount and an imino analog (typically 0.01 µg hormone and 1 µg imino compound) results in a level of hormonal activity which would correspond to that expected from much larger doses of the hormone. Enhancements of up to 10-fold have been observed. This result is understandable on the basis that the imino compound inhibits an enzyme which catalyzes the destruction of the juvenile hormone by a proton-catalyzed heterolytic cleavage of the oxido function. Such an enzymatic reaction of the juvenile hormones could lead to a 1,2-glycol by hydration or to cyclization involving the proximate double bond. Enzymic processes of both types are known (20, 21), and inhibition by imino analogs has been demonstrated (21). The rapid metabolism of the juvenile hormones reported by Dr. Siddall substantiates this supposition. It is clear that the epoxide cleavage reaction could be an important part of the natural process which regulates juvenile hormone levels in the insect (22).

The high activities of mixtures in juvenile hormone assays, e.g., the Williams-Law mixture, might be due to the presence of two types of active substances — one which has juvenile hormone activity and another which inhibits breakdown of the hormonally active substances. Further investigation of the inhibition of hormone metabolism is clearly desirable, and such studies are now being continued at Harvard (22).

Acknowledgement. The research reported herein received financial assistance from the U.S. National Science Foundation, the National Institutes of Health, and the Hoffmann-La Roche Company.

REFERENCES

- 1. Hochstein, F. A. and W. G. Brown, J. Am. Chem. Soc., 70, 3484 (1948).
- 2. Corey, E. J., J. A. Katzenellenbogen, and G. H. Posner, ibid., 89, 4245 (1967).
- 3. Corey, E. J. and G. H. Posner, ibid., 89, 3911 (1967); 90, 5615 (1968).
- 4. Corey, E. J., N. W. GILMAN, and B. E. GANEM, ibid., 90, 5616 (1968).
- 5. Corey, E. J., H. A. Kirst, and J. A. Katzenellenbogen, *ibid.*, **92**, 6314 (1970).
- 6. Corey, E. J. and L. S. Hegedus, ibid., 91, 1233 (1969).
- 7. Corey, E. J. and J. A. Katzenellenbogen, ibid., 91, 1851 (1969).
- 8. Corey, E. J., J. A. Katzenellenbogen, N. W. Gilman, S. A. Roman, and B. W. Erickson, *ibid.*, **90**, 5618 (1968).
- 9. COREY, E. J. and J. A. KATZENELLENBOGEN, unpublished; see J. A. KATZEN-ELLENBOGEN, Ph.D. thesis, Harvard University, 1969.
- 10. Corey, E. J., J. A. Katzenellenbogen, S. A. Roman, and N. W. Gilman, Tetrahedron Lett., 1821 (1971).
- 11. Corey, E. J. and H. A. Kirst, Tetrahedron Lett., 5041 (1968).
- 12. Corey, E. J. and H. Yamamoto, J. Am. Chem. Soc., 92, 226, 3523 (1970).
- 13. Corey, E. J., H. Yamamoto, D. Herron, and K. Achiwa, *ibid.*, **92**, 6635 (1970).

- 14. ROTH, W. R. and J. KÖNIG, Ann., 688, 28 (1956).
- 15. ELLIS, R. J. and H. M. FREY, Proc. Chem. Soc., 221 (1964).
- 16. Oloff, G., Tetrahedron Lett., 3795 (1965).
- 17. JORGENSON, M. J. and A. F. THACHER, ibid., 4651 (1969).
- 18. Corey, E. J. and D. HERRON, Tetrahedron Lett., in press
- 19. Corey, E. J. and H. Yamamoto, J. Am. Chem. Soc., 92, 6636, 6637 (1970).
- 20. Corey, E. J., K. Lin, and M. Jautelat, ibid., 90, 2724 (1968).
- 21. Corey, E. J., P. R. Ortiz de Montellano, K. Lin, and P. D. G. Dean, *ibid.*, **89**, 2797 (1967).
- 22. L. M. RIDDIFORD, A. M. AJAMI, E. J. COREY, H. YAMAMOTO, and J. E. ANDERSON, J. Am. Chem. Soc., 93, 1815 (1971).

Prof. E. J. COREY
Department of Chemistry
Harvard University
12 Oxford Street
Cambridge, Massachusetts 02138
USA