Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =

Bulletin de la Société Entomologique Suisse = Journal of the Swiss

Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 44 (1971)

Heft: 3-4

Artikel: Lepidopterous larvae and pupae and their parasites, collected on

pomaceous fruit trees in Israel

Autor: Plaut, H.N.

DOI: https://doi.org/10.5169/seals-401654

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 29.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Lepidopterous larvae and pupae and their parasites, collected on Pomaceous Fruit trees in Israel *

by

H. N. PLAUT

Introduction

The Lepidoptera Cydia pomonella L. (Tortricidae), Zeuzera pyrina L. (Cossidae), Spodoptera littoralis Boisd. (Noctuidae) and Lyonetia clerkella L. (Lyonetidae) are common pests of pome trees in Israel.

In the literature up to 1950, Bodenheimer (1930) mentioned 19 and Schweig (1950) two additional species and subspecies of Lepidoptera which had been occasionally encountered feeding on apple or pear trees. Most of those mentioned by Bodenheimer were collected at Qirjat Anavim (700 m), in the mountains near Jerusalem.

Since about 1960, farmers have occasionally complained of different types of feeding injury to apple and pear fruit, which in some instances reached proportions of economic damage. These cases were, and still are, usually reported after the causal factor has disappeared.

Small lepidopterous larvae often infest grafts in orchards and prevent their acceptance. They are called graft-moths by the growers, who are well aquainted with their delicate dark frass. They have not been identified.

Since the nineteen-fifties, the habit of keeping the soil as clear as possible of weeds has gradually given way to different degrees of non-tillage, which has given rise to abundant undergrowth of diverse floral composition, in many groves. The fauna which subsists on this undergrowth is an additional comparatively new factor in the ecosystem of the commercial grove, which must be taken into consideration when the possibilities of integrated pest control are being examined. Lepidoptera feeding on the weeds may play the useful role of some « indifferent » organisms, being in themselves neither directly harmful or useful, but

^{*} Contribution from The Volcani Institute of Agricultural Research, Bet Dagan, Israel. 1971 Series, No. 1330 -E.

serving as food for predators and parasites which have the potential of reducing pests of the trees.

Other possible causes for current changes in abundance and species-composition of Lepidoptera in pome-fruit groves are changing irrigation and pest control practices, changes in the nature of neighbouring crops, areal spread, and biological adaptation.

Thus, it seemed worthwhile to update our knowledge of the lepidopterous fauna of pome trees. A summary of the findings since 1962 is herewith presented.

Methods

Foliage of apple and pear trees was scanned visually, and sometimes branches were tapped. Growers and extension personnel were encouraged to report cases of feeding on the trees, especially the fruit. A current research program on the use of burlap bands on trunks of trees for the indication of population density and flight of the codling moth, also yielded information on other species of lepidoptera, which were collected systematically at Newe Ya'ar and Nahalal, and occasionally sent in from commercial groves.

Most material was collected within, or in the vicinity, of the Yizre'el Valley, but some of it came from as far north as the Huleh Valley in Upper Galilee, or as far south as Sa'ad in the southern coastal plain and Tse'elim in the western Negev. Elevations were usually lower, and in a few cases a little higher, than 200 m above sea level.

Whenever the collected material could not be directly identified, it was reared or kept, to obtain imagines. Emerging parasites were also identified, but in most cases they could not with certainty be related to their host species.

Most Lepidoptera were identified by Mr. J. Palmoni, of the Gordon Institute, Deganya Alef, and by Dr. W. Sauter, of the Department of Entomology, E.T.H., Zuerich. Dr. Sauter was very helpful in updating some of the zoological nomenclature.

Dr. Aubert, Dr. Delucchi, Dr. Fisher and Mr. J. R. Stephen, through the Center of Identification of Entomophagous Insects of the C.I.L.B., Geneva, identified most of the Ichneumonidae, Pteromalidae, Braconidae and Chalcididae, respectively.

Dr. D. Rosen, of the Hebrew University of Jerusalem, and Dr. Y. Kougler of the Tel-Aviv University, helped with the identification of some of the parasites.

Results

The results are summarized in tables 1 and 2.

Table 1

Lepidoptera collected in the larval or pupal stage on pome fruit trees during 1963–1970 (excluding common pests)

Species	Habitat of the collected specimens (burlap traps are on trunks)	Frequency of encounter (+ rare, + + + + common)
Gracillariidae Lithocolletis blancardella F.	Apple, pear : leaves (minor)	++++
TORTRICIDAE Lobesia botrana Den. & Schiff. Crocidosema plebejana Z. Acleris variegana Den. & Schiff.	Pear : fruit Apple : burlap traps Quince : burlap traps	+++++++++++++++++++++++++++++++++++++++
Pyralidae Cryptoblabes gnidielia MILL.	Apple : fruit and burlap traps Pear : fruit, burlap traps, and feeding at base of young growth	++++
Euzophera bigella Z.	Quince : fruit and burlap traps Pear : callus on bark wounds Apple : burlap traps	+++
Ectomyelois ceratoniae Z.	Quince : fruit Apple : fruit mummies	+
Lamoria anella Schiff. Antiercta ornatalis Dup. Herculia fulvocilialis Dup. Ulotricha egregialis H. + S.	Apple: burlap traps " " " Rat nests on pear trees " " " "	+ + + + + + + + +
Pterophoridae Stenoptilia bipunctatidactyla Hw.	Apple : burlap traps	+++
Drepanidae Cilix asiatica BH.	Apple : leaves	+++
GEOMETRIDAE Sterrha allongata STGR. Gymnoscelis pumilata HB. Rhodometra sacraria L. Boarmia selenaria DEN. & SCHIFF. Nychiodes palaestinensis WAGN.	Apple : fruit Apple : fruit and burlap traps Apple : burlap traps	+ + + + +
	Apple : leaves, burlap traps Pear : leaves Apple : burlap traps	+++++
Noctuidae Euxoa temera f. viliersi Gn. Scotia segetum Schiff.	Apple : seedlings Apple : burlap traps	+++

Species	Habitat of the collected specimens (burlap traps are on trunks)	Frequency of encounter (+ rare, + + + common)
Noctuidae (continued) Scotia crassa lata Tr. Scotia trux Schiff.	Apple: burlap traps Apple: stock-sucker	+ +
Peridroma saucia ochreacosta Haw. Mamestra persicariae accipitrina Esp. Mythimna unipuncta Haw. Mythimna loreyi Dup.	Apple: burlap traps " " " " " " " " " "	+ + + + + + + + + + +
Apatele psi L. Apatele rumicis L.	Apple : burlap traps Apple : fruit, burlap traps Pear : leaves, burlap traps	+++++
Apatele rumicis pallida ROTHSCH. Cryphia ochsi BRSN. Spodoptera exigua HB. Chloridea peltigera SCHIFF. Chloridea armigera HB. Chloridea nubigera H. S. Porphyrinia cochylioides GUEN. Trichoplusia ni HBN. Autographa gamma L. Syngrapha circumflexa L. Chrysodeixis chalcytes ESP. Grammodes bifasciata PET. Dysgonia torrida GN.	Apple: fruit Pear: burlap traps Apple: leaves, burlap traps Pear: leaves Apple: leaves, fruit Apple: leaves, fruit Apple: leaves, fruit Apple: burlap traps Apple: burlap traps Apple: burlap traps Apple: leaves, burlap traps Apple: burlap traps	+ + + + + + + + + + + + + + + + + + +
Lymantriidae Lymantria dispar L.	Apple : leaves	++
Arctiidae Phragmatobia fulginosa fervida Stgr.	Apple, pear : burlap traps	+

Table 2 Parasites reared from larvae and pupae of Lepidoptera collected on pome fruit trees

ICHNEUMONIDAE:

Pimpla spuria GRAV. nilotica SCHMK. Exochus coronatus GRAV. meridionator AUBERT Herpestomus arridens GRAV. Diadegma sp.

Braconidae:

Microplitis mediana Ruthe M. brachycera Thoms. Apanteles ruficrus Hal. Meteorus rubens Nees Rogas ductor Thnbg. R. bicolor Spin.

ICHNEUMONIDAE (continued):

Campoplex ensator GRAV.
C. orientalis SCHMK.
Meloboris velox HOLM.
Gelis meridionator AUBERT
Nepiera collector THBG.
Hyposoter didymator THBG.
Campoletis annulata GRAV.
f. maculipes TCHEK.

Braconidae (continued):

R. testaceus Spin.
Bracon erythrostictus Marsh.

CHALCIDIDAE:

Invreia rufitarsis ILL.

PTEROMALIDAE:

Dibrachys sp.
D. brucheanus RATZEBURG

TACHINIDAE:

Voria ruralis FALL.

Discussion

In their relation to the fruit trees, the listed Lepidoptera can be divided into three groups — those feeding on the tissue of the trees (fruit, leaves or bark), those feeding on material positioned on the trees (rat nests, lichens), and those seeking shelter, without feeding on the trees.

From the agricultural-economic viewpoint, the most immediately significant findings concern the fruit-feeders, of which *L. botrana*, *C. gnidiella* and *G. pumilata* were found to cause some economic dam-

L. botrana is a common pest of the grapevine in Israel and many other Mediterranean countries. During recent years, it has been found by the author causing some economic damage to Japanese plums, and by Kottler (1966) to roses in commercial greenhouses. The recent pressure of L. botrana on new hosts may be a result of the uprooting of a large percentage of the winegrape acreage, which forces whole populations of the pest to search for alternative hosts. In this survey it was found causing shallow scarring to pear fruit.

C. gnidiella in Israel is a well-known pest of citrus, and common on the fruits of grapevine and pomegranate and on corn cobs. It is of Mediterranean distribution. Its seasonal occurrence has been traced by AVIDOV and GOTTHILF (1960) on these and other hosts, leaving a gap for April and May. Our findings close this seasonal gap, at least partly, and add apple and pear to the list of known hosts. C. gnidiella seems to be responsible for much of the shallow scarring found on apple and pear fruit which is caused during May and June. In the Negev, it was also found to cause damage at the bases of young pear shoots (as reported by Mr. M. J. Berlinger).

G. pumilata had previously been found to be common among the pistils ("silk") of corn cobs in Israel (Plaut, unpublished data). It has

been recorded as destroying flowers of many hosts, (e.g. citrus, olives, roses, dalias, sorghum, corn) in Mediterranean countries (Marocco, Italy, Spain, Turkey). In this survey, it was found in five localities on apple trees; in one case it caused some economic damage by boring

deeply into fruits of the Golden Delicious variety.

Three species of *Chloridea* were collected on apple trees, feeding on leaves, and mainly on fruit, but no case of anything approaching economic damage was actually encountered. Growers and extension workers have repeatedly reported damage to apple fruit by what they consider to be *Chloridea* spp., but no such case has been verified by a specialist. The possibility of such damage is plausible, in the light

of our findings.

E. bigella was collected in considerable numbers from infested quince fruits, from burlap traps on quince and apple trees, and from callus on healing wounds on pear trees. It was not found feeding on apple fruit in orchards, but in the laboratory several generations were raised on otherwise unimpaired apple fruit (Plaut, 1965). It has been reported as cortiphagous on olive and other trees in Europe and northwestern Africa (Silvestri 1943), and as noxious on apple, apricot and other fruit in southern Russia (Gerasimov, 1930; Pavlowsky and Stakelberg, 1955). It is probably one of our "graft moths".

In a few isolated cases, S. allongata, A. rumicis and A. rumicis pallida

were found feeding on apple fruit.

Of the leaf feeders, B. selenaria warrants some attention. It was found in small numbers, but widely distributed on apple trees, voraciously feeding on the leaves. In the laboratory it was raised on apple fruit. According to Shoham (1965) the number of adults of B. selenaria, caught in a mercury vapor light trap near an apple grove in upper Galilee, was negligible in 1960–1962, but rose steeply from year to year during 1963–1965. The species has been mentioned from Japan, Burma, India, Madagascar, South Africa and Sicily on a variety of hosts.

In a few cases, S. exigua and L. dispar were found feeding in large

numbers on apple leaves.

The small leafminer L. blankardella was common on apple and pear in the Yizre'el Valley, but does not yet seem to have been identified from this region.

L. anella was found feeding heavily on the bark of apple trees under

burlap bands.

Of species found in burlap bands and not feeding on arboreal tissue C. ochsi Brsn. probably subsists on lichens. The host of A. ornatalis and S. bipunctatidactyla was Convolvulus sp., and of Mythimna spp. unidentified species of Gramineae. Most of the others — mainly Noctuidae — are well known polyphagous species which feed mainly on weeds and reach the burlap bands searching for shelter. This group of Lepidoptera could play a useful role in an integrated control program, being harmless to the trees but serving as an alternative food supply for

predators and parasites of fruit tree pests. Steiner (1962) has expounded on the probable usefulness of "indifferent" arthropods in apple orchards in Germany.

Acknowledgements

The author gratefully acknowledges the assistance extended by the specialists mentioned in the "Methods" section with the identification of the material. Thanks are due to Mrs. Lydia Rauf, who did the rearing and the preparations. Also, it is a pleasure to acknowledge the cooperation of agricultural extension officers and of fruit growers, who sent in material and directed attention to instances of activity of lepidopterous larvae in orchards.

Summary

Lepidopterous larvae and pupae were studied on pomaceous fruit trees during 1962–1970, by visual examination, branch tapping, collection from burlap bands around tree trunks, and on information on unusual types of fruit injury.

Of 42 species identified, apart from common pests, ten were found feeding on fruit, most of them recorded for the first time as such in Israel.

Of the fruit feeders, Lobesia botrana DEN. and SCHIFF., Cryptoblabes gnidiella MILL. and Gymnoscelis pumilata HB. caused economic damage, and Chloridea spp. are suspected of doing so. Euzophera bigella Z. seems to be a potential pest of fruits, and Boarmin selenaria DEN. and SCHIFF. of leaves.

Lamoria anella Schiff. was found feeding heavily on the bark of apple trunks, under burlap traps. The leaf minor Lithocolletis blancardella F. is

recorded for the first time from pometrees in Israel.

Among the species coming to burlap brands from weeds for shelter, Mythimna unipuncta HAW., Mythimma loreyi DUP. and Stenoptilia bipunctatidactyla Hw. were common. This group is considered potentially useful—especially if integrated control is attempted—as a reservoir of food for parasites and predators, which is little affected by insecticide applications to the trees.

Dense populations of *Herculia fulvociliatis* Dup. and *Ulotricha egregialis* H. S. were found in rat nests on pear trees. Twenty-three species of parasites were reared from the collected material, among which were twenty two Hymenoptera and one Dipteron.

REFERENCES

- 1) AVIDOV, Z. and GOTTHILF, S. (1960). The citrus peel moth, Cryptoblabes gnidiella Mill., in Israel. Ktavim 10 (3-4): 119-137 (in Hebrew).
- BODENHEIMER, F. W. (1930). Die Schädlingsfauna Palästinas. 465 pp. Berlin, P. Parey.
- GERASIMOV, (1930). Russk. Entom. Obozr. 24: 179, as cited in: SILVESTRI, F. (1943). Compendio di entomologia applicata, Parte speciale, Vol. 2, 699 pp. Portici.

- 4) Kotter, E. (1966). Plant protection experiments and observations. 25 pp. Israel Ministry of Agriculture, Akko Regional Extension Bureau. (Hebrew, with English summary).
- 5) PAVLOVSKY, E. N. and STAKELBERG, A. A. (Eds.) (1955). Forest Pest Handbook. 1097 pp. Adademija Nauk S.S.S.R., Zoologicheskii Institute, Moskva. (In Russian).
- 6) PLAUT, H. N. (1965). The Quince Moth, Euzophera bigella Z., on pomaceous fruit trees in Israel. Prelim. Rep. Nat. Univ. Inst. Agric., Rehovot, Israel. No. 196, 14 pp. (Hebrew, with English summary).
- 7) Schoham, Z. (1965). A new pest of deciduous fruit trees, Boarmia selenaria Schiff. Hassadeh 45 (9): 1073–1075. (In Hebrew).
- 8) Schweig, K. (1950). The pests of grapevine and fruit trees. 167 pp. Tel-Aviv, Hassadeh.
- 9) Sorauer, P. (1953). Handbuch der Pflanzenkrankheiten. Band 4, 518 pp. Berlin, P. Parey.
- 10) Steiner, H. (1962). Bedeutung der indifferenten Arthropoden in Obstanlagen. Entomophaga 7 (3): 263–267.

Dr H. N. PLAUT Newe Ya'ar Regional Experiment Station P.O. Haifa, Israel