

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 48 (1975)

Heft: 3-4

Artikel: A new species of Capniioneura from Yugoslavia (Plecopera, Capniidae)

Autor: Baumann, Richard W. / Kaanski, Dragica

DOI: <https://doi.org/10.5169/seals-401796>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A New Species of *Capnioneura* from Yugoslavia (Plecoptera, Capniidae)

RICHARD W. BAUMANN

Department of Entomology, Smithsonian Institution, Washington, D.C., 20560, USA

DRAGICA KAĆANSKI

Biološki institut Univerziteta, 7100 Sarajevo, M. Tita 114, Yugoslavia

While studying the Plecoptera fauna of the Morača River drainage in southern Yugoslavia we unexpectedly found an undescribed species of *Capnioneura*. This genus is restricted to the western Palearctic with most species occurring in central and southern Europe (ILLIES, BERTHÉLEMY). *Capnioneura petitpierreae* AUBERT (1960) is known only from North Africa and *Capnioneura caucasicus* ZHILTZOVA (1964) is found in the Caucasus Mountains.

Capnioneura balkanica n. sp.

Holotype male (Figs. b-f): Macropterous; length of forewings 7.5 mm; length of body 5.5 mm. General color dark brown; legs brown. Head wider than prothorax. Pronotum longer than wide. Wings hyaline, lightly infuscated around veins; venation typical for genus (Hanson, 1946). Tenth tergum brown at apex and along lateral margins, with large light triangle at base. Epiproct long and narrow, apex oval and elongate dorsally, lateral aspect broadest at base, tapering to thin pointed tip. Paraprocts quite broad, distal section tapering gradually from broad base to thin apex, outer membranous portion extending to nearly half the length of sclerotized portion. Specillum thin at apex, with narrowly rounded tip, sclerotized patch thin and narrow, situated near dorsal margin in lateral aspect. Cerci short and flat, dorsal inner corner elongate and pointed at apex.

Female (Fig. a): Macropterous; length of forewings 7.5-8.5 mm; length of body 5.5-6.5 mm. Color and general morphology similar to male. Seventh sternum prolonged along posterior margin at midline into bluntly rounded triangle which joins eighth sternum by narrow sclerotized band. Eighth sternum heavily sclerotized, median area lightly colored around long thin triangular patch which joins seventh sternum at base. Ninth sternum with arch-like sclerotized area, bases of sclerotized arch at lateral posterior corners and apex along anterior margin near genital opening.


Nymph: unknown.

Material

Holotype ♂, allotype and 2 ♀ paratypes, Slatina Creek, tributary of Morača River, Mioska (500 m), Montenegro, Yugoslavia, 22-II-1973, Baumann and Kaćanski; paratypes: small left tributary of Morača River, between highway tunnels 20 and 21, 23-III-1974, D. Kaćanski, 1 ♀; left tributary Morača River, at

bridge below Morača Monastery (300 m), 23-II-1974, D. Kaćanski, 1 ♀; Morača River, at bridge below Morača Monastery (300 m), 3-XI-1973, D. Kaćanski, 1 ♀; Meljestak Creek, near junction Morača River, 23-II-1973, R. W. Baumann, 1 ♀.

Holotype (No. 73289), allotype and three paratypes deposited at the National Museum of Natural History, Smithsonian Institution, Washington, D. C., USA. One paratype deposited in the following collections: D. Kaćanski, Sarajevo, Yugoslavia; Biological Institute, Titograd, Yugoslavia; Max-Planck Limnological Station, Schlitz, Germany.



Figs. a-f: *Capnioneura balkanica* n. sp. a. ♀ terminalia, ventral. b. ♂ terminalia, dorsal. c. ♂ cercus, left. d. ♂ paraproct, left. e. epiproct, lateral. f. epiproct and distal part of tenth tergum.

Discussion

Capnioneura balkanica is a large species that is most similar to *C. petitpierreae* Aubert and *C. petricola* Giudicelli (1967). The males can be separated by the shape of the epiproct: short and blunt, appearing as a stout rounded triangle in dorsal aspect in *C. petricola*; long and thin in both *C. petitpierreae* and *C. balkanica* but more delicate and elongate in *C. balkanica*. The females of both *C. petricola* and *C. petitpierreae* exhibit a broad triangular junction at the midline of sterna seven and eight. In *C. balkanica* this junction is very narrow and results in triangular sclerotized areas on both segments instead of one large triangular area.

Ecology

All collections to date are from small creeks and streams at elevations above 300 meters. The single collection from the Morača River below Morača Monastery is from near the mouth of a small tributary stream. Adults emerge from November to March during the coldest time of the year.

Acknowledgements

This study was part of the Skadar Lake Project, Smithsonian Foreign Currency Program (Grant No. SFG-2-6308), which includes studies on the ecology of Skadar Lake and its major tributaries in the southern Yugoslavian republic of Montenegro. Thanks are given to Dr. Bozina Ivanovic, Biological Institute of Titograd and Dr. Robert P. Higgins, Oceanography and Limnology Program, Smithsonian Institution, the principal investigators, for their help and encouragement. We are also indebted to Dr. Gerald Z. Jacobi, University of Wisconsin, Stevens Point, for arranging field collecting trips and to Dr. Catherine J. Kerby, Oceanography and Limnology Program, Smithsonian Institution, for valuable administrative assistance. The drawings were made by Rebecca F. Surdick.

Literature Cited

- AUBERT, J., (1960). *Contribution à l'étude des Plécoptères du Maroc*. Mitt. Schweiz. Ent. Ges. Lausanne, 33 (4): 213-222.
BERTHÉLEMY, C., (1969). *Les Capnioneura des Pyrénées (Plecoptera)*. Annls. Limnol., Toulouse, 5 (1): 25-47.
GIUDICELLI, J., (1967). *Capnioneura petricola n. sp., Plécoptère nouveau de Corse*. Annls. Limnol., Toulouse, 3 (2): 249-266.
HANSON, J. F., (1946). *Comparative Morphology and Taxonomy of the Capniidae (Plecoptera)*. Amer. Midl. Natural., Notre Dame (Indiana), 35 (1): 193-249.
ILLIES, J., (1966). *Katalog der rezenten Plecoptera*. Das Tierreich, Berlin, 82: I-XXX; 1-632.
ZHILTOVA, L. A., (1964). *On the knowledge of Plecoptera from the Caucasus, VI*. New species of Taeniopterygidae, Nemouridae and Capniidae. Ent. Obozr., Moscow, 43: (2): 347-362.

