

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 57 (1984)

Heft: 2-3

Artikel: Indian species of the genus *Stollia* Ellenrieder (Heteroptera, Pentatomidae)

Autor: Azim, M. Nayyar / Shafee, S. Adam

DOI: <https://doi.org/10.5169/seals-402125>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Indian species of the genus *Stollia* ELLENRIEDER (Heteroptera, Pentatomidae)

M. NAYYAR AZIM & S. ADAM SHAFEE

Section of Entomology, Department of Zoology, Aligarh Muslim University, Aligarh, India

The genus *Stollia* ELLENRIEDER is redefined. *S. spinosa* sp. n. is described and illustrated. *S. guttiger* (THUNBERG) is also studied and illustrated. A key for the separation of Indian species is given. Material deposited in the Zoological Museum, Aligarh Muslim University, Aligarh.

Genus *Stollia* ELLENRIEDER

Stollia ELLENRIEDER, 1862: 149.

Type-species: *S. filiginosa* ELLENRIEDER.

Bainbriggeanus DISTANT, 1918: 135 (LESTON, 1956: 702).


Type-species: *Bainbriggeanus fletcheri* DISTANT.

Diagnosis: Head distinctly wider than long; juga as long as tylus, lateral margins sinuate before eyes; rostrum extending slightly beyond hind coxae; bucculae shorter than first rostral segment; antennae 5-segmented, first segment not reaching apex of head; scutellum broadly rounded apically; mesosternum sulcated with indistinct longitudinal carina medially; metasternal scent gland ostiole with peritreme small, apex of which slightly raised and directed anteriorly, evaporatoria well developed; abdominal venter unsulcated and unarmed at base; last tergum in female (fig. D) with anterior and posterior margins convex. *Female genitalia*: external plates (fig. C), first gonocoxae broadly triangular, paratergites 8th and 9th triangular; spermatheca (fig. E), dilation deeply constricted at proximal one third, bulb semicircular without tubular outgrowths. *Male genitalia*: Claspers (fig. F) L-shaped and with fine ridges; pseudoclaspers (fig. G) spatulate; subgenital plate (fig. H) horse-shoe-shaped, anterior margin with a prolongation medially.

The genus is represented by two species from India including a new species. The two species can be separated on the basis of the following key characters:

Key to Indian species of *Stollia* ELLENRIEDER

1. Pronotum (fig. A) with humeral angles obtuse, anterolateral margins smooth and straight; antennae (fig. B) with second and third segments subequal in length; membrane of hemelytra reaching beyond apex of abdomen *S. guttiger* (THUNBERG)
- pronotum (fig. I) with humeral angles much acute, anterolateral margins serrated and sinuate; antennae (fig. J) with second segment distinctly shorter than third; membrane of hemelytra reaching apex of abdomen. *S. spinosa* sp. n.



Figs. A-H. *Stollia guttiger* (THUNBERG), ♀, ♂; A. Entire body in dorsal view, ♀; B. Antenna, ♀; C. External genitalia, ♀; D. Last tergum, ♀; E. Spermatheca, ♀; F. Clasper, ♂; G. Pseudoclasper, ♂; H. Subgenital plate, ♂.

Figs. I-J. *Stollia spinosa* sp. n., ♀; I. Head and Thorax in dorsal view, ♀; J. Antenna, ♀.

Stollia guttiger (THUNBERG)
(figs. A-H)

Cimex guttiger THUNBERG, 1783: 32.

Stollia guttiger (THUNBERG); Stal, 1876: 81.

Bainbriggeanus fletcheri DISTANT, 1918: 135 (LESTON, 1956: 702).

Material examined: -5 ♀, 6 ♂, India: Uttar pradesh, Aligarh, University Botanical garden, on *Croton sp.*, 5.VII.1978 (M. NAYYAR AZIM); 5 ♀, 7 ♂, Tamil Nadu, Madras, Myalapore, on grass, 15.XI.1979 (M. NAYYAR AZIM).

Stollia spinosa sp. n.

Female (figs. I-J).

Head dark, densely and darkly punctate, distinctly wider than long (2.4 : 1.8 mm); juga as long as tylus, truncated apically, lateral margins sinuate before eyes; eyes reddish brown; ocelli red, space between ocellus and inner orbital margin about one-fifth the inter-ocellar space. Rostrum brownish except apical segment dark, extending just beyond hind coxae, segments I, II, III and IV, 0.79, 1.27, 0.4 and 0.57 mm in length respectively. Antennae yellowish brown except apical segment dark, segments I, II, III, IV and V, 0.42, 0.74, 0.82, 0.9 and 1.18 mm in length respectively.

Thorax dark brown, densely and darkly punctate except two triangular patches on anterior area of pronotum and one on each basal angle of scutellum laevigate; pronotum with anterior margin deeply concave, antero-lateral margins sinuate and serrated, humeral angles produced laterally into acute spines; maximum width of pronotum about three times as wide as long (6.9 : 2.2 mm); scutellum gradually narrowing apically, as wide as long (3.7 : 3.7 mm); membrane of hemelytra reaching apex of abdomen. Legs yellowish except two dark patches near apex of femora.

Abdomen dark except connexivum with alternating dark and yellow bands, densely punctate. Since the female genitalic characters have generic significance, there is no remarkable difference in the genitalia between the females of *S. guttiger* and *S. spinosa*.

Body length: 8 mm.

Holotype. 1 ♀, India: Uttar pradesh, Agra, on grass, 30.XI.1983 (M. NAYYAR AZIM).

ACKNOWLEDGEMENTS

The authors are deeply indebted to Prof. NAWAB H. KHAN, Chairman, Department of Zoology, Aligarh Muslim University, Aligarh, for providing research facilities. One of us (M. N. AZIM) is thankful to U. G. C., New Delhi for providing financial help.

LITERATURE

DISTANT, W. L. 1918. *The fauna of British India including Ceylon and Burma*. Rhynchota Vol. 7 (London, Taylor & Francis).

ELLENRIEDER. 1862. *Natuurkundig Tijdschrift voor Nederlandsch Indie*. Batavia.

LESTON, D. 1956. *Pentatomoidea (Hemiptera) New species and synonymy*. Ann. Mag. Nat. Hist. 12: 698-704.

STAL, C. 1876. *Enumeratio Hemipterorum 5. Kongl. Svenska Vetensk. Akad. Hand. 14*: 1-162.

THUNBERG, C. P. 1783. *Dissertatio Entomologica Novas Insectorum species*. 2 Upsaliae.

(received April 17, 1984)

