

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 57 (1984)

Heft: 4: Festschrift Prof. P. Bovey

Artikel: First report of Zaga Girault and Oligositoides Doult (Hymenoptera :
Trichogrammatidae) from India, with descriptions of three new species

Autor: Yousuf, Mohammad / Shafee, S. Adam

DOI: <https://doi.org/10.5169/seals-402133>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 31.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

First report of *Zaga* GIRAULT and *Oligositoides* DOUTT
(Hymenoptera: Trichogrammatidae) from India, with descriptions of
three new species

MOHAMMAD YOUSUF & S. ADAM SHAFEE

Section of Entomology, Department of Zoology, Aligarh Muslim University, Aligarh, India.

The genera *Zaga* GIRAULT and *Oligositoides* DOUTT are reported for the first time from India. Three new species: *Z. angustipennis* sp. n., *O. terebratus* sp. n., and *O. gudurensis* sp. n. are described and illustrated. Types deposited in Zoological Museum, Aligarh Muslim University, Aligarh, India.

Genus *Zaga* GIRAULT

Zaga GIRAULT, 1911: 31.

Type-species: *Zaga latipennis* GIRAULT.

Comment: The genus differs from other genera of Trichogrammatidae in having antennae without funicle and fore wings with discal setae arranged in lines. The genus is known to contain two species including one new species. The two species are separated by the following key characters:

1. ♀: Fore wings (DOUTT & VIGGIANI, 1968: fig. 52 B) broad, twice as long as wide; marginal vein about as long as premarginal vein; marginal fringe very short, less than one-tenth of wing width; antennae with pedicel one and a half times as long as wide, club 4-segmented. *Z. latipennis* GIRAULT
- ♀: Fore wings (fig. B) moderately narrow, slightly less than three times as long as wide; marginal vein much longer than premarginal vein; marginal fringe long, one-fourth of wing width; antennae with pedicel two times as long as wide, club 5-segmented. *Z. angustipennis* sp. n.

Zaga angustipennis sp. n. (Figs. A, B)

Female


Head yellowish brown; eyes dark; malar space slightly shorter than eye width. Antennae (fig. A) yellowish brown, inserted at lower level of eyes; scape gently narrowing apically, three and a half times as long as wide; pedicel twice as long as wide; single ring segment distinct; funicle absent; club 5-segmented, three times as long as wide.

Thorax dark brown, sparsely setose. Fore wings (fig. B) hyaline, slightly less than three times as long as wide; disc with setae arranged in lines; costal cell narrow, longer than marginal vein; marginal vein much longer than stigmal and premarginal veins separately; postmarginal vein rudimentary; marginal fringe long, one-fourth of wing width; legs yellowish brown; hind coxae much longer than fore and middle coxae.

Abdomen yellowish brown, distinctly longer than thorax; ovipositor slightly exerted, arising from base of abdominal venter.

Body length: 0.52 mm.

Holotype: ♀, India: Uttar Pradesh, Aligarh, 10.X.1983 (M. YOUSUF).



Figs. A, B: *Zaga angustipennis* sp. n., ♀: (A) Antenna; (B) Fore wing.

Figs. C, D: *Oligositoides terebratus* sp. n., ♀: (C) Antenna; (D) Fore wing.

Figs. E, F: *Oligositoides gudurensis* sp. n., ♀: (E) Antenna; (F) Fore wing.

Genus *Oligositoides* DOUTT

Oligositoides DOUTT in DOUTT & VIGGIANI, 1968: 541.

Type-species: *Oligositoides fletcheri* DOUTT.

Comment: The genus is allied to *Oligosita* WALKER in possessing a single funicle segment and a 3-segmented club, but differs from that genus in having forewings broadened apically, with discal setae arranged in lines. The genus is represented by three species including two new species. A key for their separation is given below:

Key to species of *Oligositoides* DOUTT, based on females

1. Fore wings with marginal vein distinctly longer than stigmal vein; marginal fringe long, one-sixth or more the wing width. 2
- Fore wings with marginal vein slightly shorter than stigmal vein (DOUTT and VIGGIANI, 1968: fig. 42 B); marginal fringe very short, less than one-tenth the wing width. *O. fletcheri* DOUTT
2. Ovipositor much exserted; funicle segment about as long as wide (fig. C); fore wings with marginal fringe one-third the wing width (fig. D). . *O. terebratus* sp. n.
- Ovipositor slightly exserted; funicle segment wider than long (fig. E); fore wings with marginal fringe about one-sixth the wing width (fig. F) *O. gudurensis* sp. n.

Oligositoides terebratus sp. n. (Figs. C, D)

Female

Head yellowish brown, slightly wider than long in facial view; ocelli red, lateral ocellus separated by more than its diameter from inner orbital and less than its diameter from occipital margins. Antennae (fig. C) yellowish brown; pedicel distinctly longer than ring and funicle segments together; funicle 1-segmented, as long as wide; club 3-segmented, about four times as long as wide, apical end terminating into a rod-like projection.

Thorax dark brown. Fore wings (fig. D) two and a half times as long as wide, outer margin broadly rounded; disc with setae arranged in lines; costal cell well developed, longer than marginal vein; marginal vein distinctly longer than stigmal vein, with 3 long setae; postmarginal vein absent. Hind wings long and narrow; marginal fringe twice the wing width. Legs yellowish brown.

Abdomen brown, distinctly longer than thorax; ovipositor much exserted, arising from basal one-fourth of abdominal venter, exserted part of ovipositor slightly less than one-half the length of abdomen.

Body length, excluding exserted part of ovipositor: 0.51 mm.

Holotype: ♀, India: Andhra Pradesh, Gudur, 30.I.1984 (M. YOUSUF).

Oligositoides gudurensis sp. n. (Figs. E, F)

Female

Head brown, slightly wider than long in facial view; eyes dark; malar space shorter than eye width. Antennae (fig. E) yellowish brown, inserted at lower level of eyes; scape about three times as long as wide; pedicel slightly less than twice as long as wide, longer than ring and funicle segments together; funicle 1-segmented, slightly wider than long; club 3-segmented, slightly less than four times as long as wide.

Thorax dark brown. Fore wings (fig. F) hyaline except three small infuscated patches arranged in triangle beneath the venation; slightly less than two and a half times as long as wide, broadly rounded apically; disc with setae arranged in lines; costal cell longer than marginal vein, marginal vein thrice the length of stigmal vein, postmarginal vein absent, stigmal vein slightly shorter than premarginal vein; marginal fringe about one-sixth the wing width. Legs yellowish except hind femora dark.

Abdomen dark brown, slightly longer than thorax; ovipositor slightly exerted, arising from base of abdominal venter.

Body length excluding exerted part of ovipositor: 0.63 mm.

Holotype: ♀, India: Andhra Pradesh, Gudur, 30.I.1984 (M. YOUSUF).

ACKNOWLEDGEMENTS

The authors are greatly indebted to Prof. NAWAB H. KHAN, Chairman, Department of Zoology, for providing research facilities. One of us (M. YOUSUF) is thankful to C.S.I.R., New Delhi, for providing financial assistance.

LITERATURE

DOUTT, R. L. & VIGGIANI, G. 1968. *The classification of Trichogrammatidae (Hymenoptera: Chalcidoidea)*, Proc. Calif. Acad. Sci. 35: 477-586.