

<b>Zeitschrift:</b>	Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society
<b>Herausgeber:</b>	Schweizerische Entomologische Gesellschaft
<b>Band:</b>	58 (1985)
<b>Heft:</b>	1-4: Fascicule-jubilé pour le 80e anniversaire du Prof. Dr. Paul Bovey = Festschrift zum 80. Geburtstag von Prof. Dr. Paul Bovey
<b>Artikel:</b>	Cryphalus abietis (Ratz.) and Ips typographus (L.) new for Turkey, and a note on the tree killing capacity of Pityophthorus pityographus (Ratz.)
<b>Autor:</b>	Benz, G.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-402161">https://doi.org/10.5169/seals-402161</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

*Cryphalus abietis* (RATZ.) and *Ips typographus* (L.) new for Turkey,  
and a note on the tree killing capacity of *Pityophthorus*  
*pityographus* (RATZ.)

G. BENZ

Departement of Entomology, Swiss Federal Institute of Technology, CH-8092 Zurich, Switzerland.

The granulated spruce bark beetle, *Cryphalus abietis* (RATZ.), and the eight-toothed spruce bark beetle, *Ips typographus* (L.), are reported for the first time on oriental spruce, *Picea orientalis* L., in the Artvin District in northeastern Turkey.

The oriental spruce, *Picea orientalis* L., growing in large forests in the eastern Pontic Mountains, south of the Black Sea shore of Turkey, is attacked by several bark beetles (Scolytidae). In the Artvin District *Dendroctonus micans* (KUG.) and *Ips sexdentatus* BÖRN. are the most dangerous species at present. On a FAO mission to Turkey for the study of *D. micans* in 1984 the author found *P. orientalis* with BHDS of 20 cm or less killed by the bark beetles *Cryphalus abietis* (RATZ.) and/or *Pityophthorus pityographus* RATZ. in the region of Ardanuç. *P. pityographus* was known to occur in Turkey (recorded by SCHIMITSCHEK (1944) as *P. micrographus* L.); but *C. abietis* is a new record for Turkey.

Many of the trees looking dead from a distance turned out to be alive in the lower quarter but were losing or lacking needles in the upper parts. In the dying or dead sections, the stems were attacked by one or the other of the two bark beetle species, *P. pityographus* being found more often than *C. abietis* in the fully dead trees.

It was interesting to note that the trees infested by *P. pityographus* were losing their needles on all the branches at the level of the attacks and higher up as soon as the males had finished the nuptial chamber and the females started to bore the egg galleries. Since the ascending sap flow is unlikely to be hindered at such an early stage of attack, a «toxic» action of *P. pityographus* on *P. orientalis* must be considered and should be further investigated.

In July 1984 *Ips typographus* (L.) was first found to attack *P. orientalis* on Turkish territory in the Hicaköy/Demirkapi region near the border line to the USSR. The present distribution of the species suggests that it has recently invaded Turkey from Sovjet Georgia, where its association with *P. orientalis* is of old record and produces mass outbreaks in stands weakened by attacks of *D. micans* (KOBACHIDZE, 1967). *I. typographus* is therefore a new threat against the Turkish spruce forests.

REFERENCES

- KOBACHIDZE, D. N. 1967. *Der Riesenbastkäfer (Dendroctonus micans KUGEL.) in Georgien (UDSSR)*. Anz. Schädlingeskde. 40, 65–68.  
SCHIMITSCHEK, E. 1944. *Forstinsekten in der Türkei und ihre Umwelt*. Volk und Reich Verlag, Prag, 371 pp.

(received February 11, 1985)