

**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

**Herausgeber:** Schweizerische Entomologische Gesellschaft

**Band:** 58 (1985)

**Heft:** 1-4: Fascicule-jubilé pour le 80e anniversaire du Prof. Dr. Paul Bovey = Festschrift zum 80. Geburtstag von Prof. Dr. Paul Bovey

**Artikel:** Descriptions of two new species of Leptanusia (Hymenoptera : Encyrtidae) from Patna, India

**Autor:** Shamim, S.M. / Shafee, S. Adam

**DOI:** <https://doi.org/10.5169/seals-402167>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Descriptions of two new species of *Leptanusia* (Hymenoptera: Encyrtidae) from Patna, India

S.M. SHAMIM<sup>1</sup> & S. ADAM SHAFEE<sup>2</sup>

<sup>1</sup> Post-Graduate Department of Zoology, Ranchi University, Ranchi, India.

<sup>2</sup> Department of Zoology, Aligarh Muslim University, Aligarh, India.

Two new species: *Leptanusia brevicornis* sp. n. and *Leptanusia varicornis* sp. n. from Patna are described and illustrated. Revised key to species of *Leptanusia* DE SANTIS is also given. Types deposited in Zoological Museum, Aligarh Muslim University, Aligarh, India.

Genus *Leptanusia* DE SANTIS


*Leptanusia* DE SANTIS, 1963: 80.

Type-species: *Leptomastidea pseudococci* BRETHES, by monotypy.

The generic characters proposed by DE SANTIS (1963) and ALAM & SHAFEE (1981) for the genus *Leptanusia* apply well on the species including two new species. HAYAT *et al.* (1975) key to species of *Leptanusia* DE SANTIS has been revised in order to accommodate the new species.

*Revised key to species of Leptanusia* DE SANTIS, based on females.

1. Funicle of uniform colour; funicle segment first more than two and a half times as long as wide, as long as or longer than pedicel; fore wings with postmarginal vein as long as or longer than stigmal vein . . . . . 2  
– Funicle with black and white segments; funicle segment first at most twice as long as wide, shorter than pedicel; fore wings with postmarginal vein much shorter than stigmal vein . . . . . 4
2. Club brownish, three times as long as wide; postmarginal and stigmal veins subequal . . . . . 3  
– Club white, two and a half times as long as wide; postmarginal vein much longer than stigmal vein . . . . . *pseudococci* (BRETHES)
3. Head dorsum slightly less than two times as wide as long; width of frontovertex more than one-third head width; scape brownish with a long whitish strip in middle . . . . . *Szelenyii* HOFFER  
– Head dorsum more than two times as wide as long; width of frontovertex about one-third head width; scape whitish with a long patch on dorsal side at distal half, and an irregular patch on ventral surface . . . . . *quadri* HAYAT *et al.*
4. Funicle segments 1,4 and 5th white, remaining dark; pedicel as long as following two funicle segments together; funicle segments 1–6 quadrate; disc of forewings with five patches of transparent setae . . . . . *brevicornis* sp. n.



Figs. A-F, A-C, *Leptanusia brevicornis* sp. n., ♀: A, Antenna; B, forewing; C, part of forewing venation. D-F, *Leptanusia varicornis* sp. n., ♀: D, antenna; E, fore wing; F, part of fore wing venation.

- Funicle segments 2-4 white, remaining dark; pedicel distinctly shorter than following two funicle segments together; funicle segments 1-6 longer than wide; disc of fore-wings with semicircular band of transparent setae *varicornis* sp. n.

*Leptanusia brevicornis* sp. n. (Figs. A–C)

*Female*

Head orange yellow, finely punctate; frontovertex slightly less than twice as long as wide; ocelli white, arranged slightly in obtuse triangle, lateral ocellus separated by its own diameter from inner orbital and occipital margins separately; eyes silvery white; malar sutures absent; malar space shorter than eye width; antennae inserted below ocular line, interantennal space slightly less than width of frons between eyes at median ocellus; mandibles bidentate. Antennae (Fig. A) white except funicle segments 2nd, 3rd, 6th and club dark; scape lightly infuscated, basal three-fourth and inner margin dark; scape flattened, two and a half times as long as wide; pedicel about twice as long as wide, as long as basal two funicle segments together; funicle segments quadrate; club 3-segmented, slightly less than three times as long as wide, about as long as preceding four funicle segments together.

Thorax yellowish except dorsum infuscated; mesoscutum without parapsidal furrows. Fore wings (fig. B) with patches of coarse and hyaline setae; slightly less than three times as long as wide; costal cell narrow; submarginal vein with 15 setae; marginal vein slightly longer than stigmal vein; postmarginal vein rudimentary (fig. C).

Abdomen orange yellow except dorsum infuscated, longer than thorax; cercal plates located on basal one-third, paratergites present; subgenital plate reaching apex of abdomen; ovipositor hidden, arising from apical one-third of abdominal venter.

Body length: 1.22 mm.

Holotype ♀, INDIA: Bihar, Patna, Danapur, ex *Ferrisia virgata* COCKERELL, 10.I.1984 (S.M. SHAMIM).

*Leptanusia varicornis* sp. n. (Figs. D–F)

*Female*

Resembles *L. brevicornis* sp. n. except in the following characters:

Antennae (fig. D) dark except apical one-third of scape, apical half of pedicel, funicle segments 2–4 white; pedicel much shorter than following two funicle segments together; funicle segment first twice as long as wide, 2–6 distinctly longer than wide; fore wings (fig. E) with semicircular band of transparent setae apically, slightly more than twice as long as wide; marginal fringe short.

Body length: 1.4 mm.

Holotype ♀, INDIA: Bihar, Patna, Danapur, 10.I.1984, (S.M. SHAMIM).

ACKNOWLEDGEMENTS

The authors are deeply indebted to Prof. NAWAB H. KHAN, Chairman, Department of Zoology, Aligarh Muslim University, Aligarh for providing research facilities. One of us (S.M. SHAMIM) is also thankful to Dr. P. N. MEHROTRA, Head, Post-graduate, Department of Zoology, Ranchi University, for encouragement and for providing research facilities.

LITERATURE

- DE SANTIS, L. 1963. *Encyrtidos de la Republica Argentina*. An. Comm. Invest. Cient. Prov. Bs Aires 4: 9–422.  
HAYAT, M., ALAM, S.M. & AGARWAL, M.M. 1975. *Taxonomic survey of encyrtid Parasites (Hymenoptera: Encyrtidae) in India*. Alig. Mus. Univ. Publ. (Zool. Ser.) Indian Ins. Typ. 9: 1–112.  
ALAM, S.M. & SHAFEE, S.A. 1981. *Significance of morphological structures in the classification of Indian Encyrtidae (Chalcidoidea: Hymenoptera)*. Proc. Indian nat. Sci. Acad. B 47:775–798.

(received March 20, 1985)

