

Zeitschrift:	Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society
Herausgeber:	Schweizerische Entomologische Gesellschaft
Band:	59 (1986)
Heft:	3-4
Artikel:	The puparium of <i>Placopsidella cynocephala</i> Kertész (Diptera : Ephydidae) and notes on its biology
Autor:	Kirk-Spriggs, A.H.
DOI:	https://doi.org/10.5169/seals-402247

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The puparium of *Placopsidella cynocephala* KERTÉSZ (Diptera: Ephydriidae) and notes on its biology

A. H. KIRK-SPRIGGS

Department of Zoology, National Museum of Wales, Cathays Park, Cardiff CF1 3NP, Wales, United Kingdom

The puparium of *Placopsidella cynocephala* KERTÉSZ (Diptera: Ephydriidae: Psilopinae) is described and figured, from material developing in decaying tube worms in stacked coral found on the beach at Bacan, Indonesia. Other larvae found in the same situation are discussed.

During September 1985 when on the island of Bacan, Indonesia, I observed on a heap of stacked coral on the sandy beach at Labuha huge numbers of feeding *Placopsidella cynocephala* KERTÉSZ which almost blanketed the heap. This was stinking from the decay of the marine animals it contained. In the decaying matter of the tube worm (Polychaete) holes I observed large numbers of small Cyclorrhaphous larvae feeding. A sample of these was put into spirit and other larvae put into tubes of sawdust for pupation. Unfortunately only a single larva succeeded in pupating, from which a *P. cynocephala* emerged. It became evident when examining the puparium that the other larvae preserved in spirit represented an entirely different taxon, possibly a tethinid, which were also common in the same locality.


It is not unusual to find such heaps of coral in such situations. As burnt coral provides lime for house building it is possible that the coral had been stacked there with such a purpose in mind.

Recently MATHIS (1986) has revised the genus *Placopsidella*, dividing it into 8 species, 2 of which were new to science. The material listed above has been identified with the aid of the key to species and figures given in that publication.

DESCRIPTION


Placopsidella cynocephala KERTÉSZ (figs. 1–10)

Puparium (fig. 1) overall length 3.37 mm; breadth at widest point (mid-length) 0.94 mm. Shape, thickened centrally, convex, slightly dorso-ventrally flattened, anteriorly and posteriorly truncate; surface very slightly shining, colour pale tan-brown, either side laterally with a series of two protuberances, the more ventral of which is scale-like and the more dorsal having a fan of seven long pointed spicules (fig. 10); situated dorso-laterally is another spike-like protuberance and situated on either side of mid-line dorsally is a less prominent protuberance, all these others lacking the apical spicules present on the latero-dorsal protuberance, all the above situated on the abdominal segments; ten spicular zones present, encircling the puparium, those on the ventral surface (fig. 7) closely grouped together, more so on the first few anterior rows; rows on dorsal surface with



Figs. 1–4. *Placopsidella cynocephala* puparium (1) whole puparium, dorsal aspect; (2) pupal cephalopharyngeal skeleton from above; (3) the same, in profile; (4) posterior end of puparium from behind.

spicules (fig. 8) not contiguous at their bases but widely spaced, encircling lateral and dorsal protuberances; all spicules of the coarse-pointed type. Anterior spiracles (fig. 5) of the fan type, consisting of 8 short digitations on a semicircular base, distance between them 0.24 mm. Posterior spiracles (figs. 4 and 9) with the usual three spiracular openings (not visible in figure 4), darkened, short. Anal plate (fig. 6) membranous and of indistinct form. Cephalopharyngeal skeleton (figs. 2



Figs. 5–10. *Placopsidella cynocephala* puparium (5) anterior spiracle, left side in profile; (6) anal plate, ventral aspect (7) ventral spicular zone eight; (8) dorsal spicular zones; (9) posterior spiracle, from above (10) latero-dorsal protuberance, from above. Scales: peak lines at base of each photograph 5, 6, 7, 9 = 100 μ m; 8 = 1000 μ m; 10 μ m.

and 3); mandibular sclerite completely black with a single tiny window situated at mid-length, unserrated with anterior half sickle-shaped; hypostomal sclerite black, with a small scale-like dentary sclerite; pharyngeal sclerite brown/black in colour, ventral wing paler in colour at its posterior end, having regular longitudinal grooves (pharyngeal sieving mechanism), both dorsal and ventral wings with a forwardly-directed spur, which is shorter in the dorsal wing and more curved

forward and longer in the ventral; dorsal hood (fig. 2) having its surface pale in colour with rows of holes giving a lacey appearance.

MATERIAL EXAMINED

Adult and associated puparium of the following: 1 ♂ Indonesia: Maluku Utara, Bacan, Labuha, 26.IX.1985, ex rotten tube worms in stacked coral on beach (A. H. KIRK-SPRIGGS). Deposited in the collections of the National Museum of Wales NMW.Z.1985.078. A specimen in British Museum (Natural History) New Guinea; ex logs, 31.VII.1968 (P. W. HOWARD) has a puparium mounted with it. This puparium in my opinion is not that from which the adult fly emerged, but is probably of a chloropid.

DISTRIBUTION

This species is widespread, occurring in Australia, Belau, Federated States of Micronesia, Gilbert Islands, Guam, Indonesia, Kiribati, Malaysia, Marshall Islands, Papua New Guinea, and the Philippines, and appears to be a coastal species.

BIOLOGY

Little is known of the biology of *Placopsidella*. Tenorio (1980: 270–271), gives information on the biology of *P. marquesana* (Malloch, 1933: 13) (as *P. cynocephala*) and gives figures of female ventral receptacle, larva and puparium (reproduced in MATHIS, 1986: 14). He states that this species occurs commonly in seaweed and other debris especially sugarcane on beaches of the Hawaiian Islands. It is not known, however, as Mathis points out, whether the flies were feeding on the seaweed itself or as scavengers on crustacea or other animals among the seaweed.

MATHIS, 1986: 10) gives a review of the biological information of *P. cynocephala* where he sights material from Cape York Peninsula (Australia) "breeding in dead mussels *Fluviolanahus subtortus* (BUNKER)" and specimens collected at Ulebaehel Island (Belau) taken from barnacles attached to a stick. Further to this BOHART & GRESSITT (1951: 87) found this species being attracted in small numbers to traps baited with echinoderms and sea-cucumbers on the island or Guam.

The present record is from a very similar breeding medium. This would strongly suggest that this species develops in dead marine invertebrates which have become stranded. Furthermore the pharyngeal sieving mechanism of the pharyngeal sclerite indicate bacterial feeding. Such a protein-rich substance in such temperature would have a high bacterial content.

COMMENTS

The immature stages of this species show close affinities to those of *P. marquesana*, having very similar pupal shape and spiracular formation. The cephalopharyngeal skeleton, however, varies greatly in the shape of the mandibular, dentary and hypostomal sclerites, the pharyngeal sclerite in *P. marquesana* lacks the

hook on the upper wing and from the figures given by Tenorio, 1980: 350–351 lacks the pharyngeal sieving mechanism on the lower wing.

ACKNOWLEDGEMENTS

I am most grateful to J. C. DEEMING (National Museum of Wales) for his kind help and encouragement, Shell Research Ltd, ICI Research Ltd, The British Ecological Society, The Professor Hering Memorial Research Fund, University College, Cardiff, Colemans Ltd, my father Mr. K. R. KIRK-SPRIGGS and the National Museum of Wales who kindly provided funds for this research. I would also like to thank the Director Dr. D. W. DYKES and Mr. P. J. MORGAN (National Museum of Wales) for the use of S. E. M. and other facilities.

LITERATURE

BOHART, G. E. & GRESSITT, J. L. 1951. *Filth inhabiting Flies of Guam*. Bulletin Bernice P. Bishop Museum No. 204, 152 pp.

MATHIS, W. N. 1986. *Studies of Psilopinae (Diptera: Ephydriidae), I. A Revision of the Shore Fly Genus Placopsidella Kertész*. Smithsonian Contributions to Zoology No. 430: 30 pp. 35 figures, Washington D. C.

TENORIO, J. A. 1980. *Family Ephydriidae*. In: D. E. HARDY & M. D. DELFINADO, eds. *Diptera: Cyclorrhapha III, Insects of Hawaii, 13*: 255–351, 141 figures. Honolulu: University Press of Hawaii.

(received March 17, 1986)

