

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 60 (1987)

Heft: 3-4

Artikel: Two new species of Uroleucon Mordv. (Homoptera : Aphididae) from Iran

Autor: Rezwani, A. / Lampel, G.

DOI: <https://doi.org/10.5169/seals-402281>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 03.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Two new species of *Uroleucon* MORDV. (Homoptera: Aphididae) from Iran

A. REZWANI¹ & G. LAMPEL²

¹ Plant Pests and Diseases Research Institute, 19395-Tehran, P.O. Box 1454, Iran

² Zoological Institute of the University, Entomological Department, CH-1700 Fribourg, Switzerland

The apterous viviparous females of two new species of *Uroleucon* MORDV. living on *Scorzonera tortuosissima* BOISS. and *Hymenocephalus rigidus* JAUB. & SPACH. are described from specimens collected in Iran.

1. *Uroleucon (Uroleucon) tortuosissimae* sp. n.

Apterous viviparous female (Figs. 1 and 2)

Colour in living specimens: Mat brown, cauda pale, siphunculi dark brown. Macerated specimens: Head and coxae brownish, antennal joints I and II dark brown, basal part of third and whole length of fourth antennal joint pale or brown, medial part of third antennal joint often darker than the rest of the same joint. The last two segments of the rostrum brown, thorax and abdominal segments colourless, without sclerites, postsiphuncular sclerites not developed, cauda colourless, genital plate smoky, siphunculi brown to dark brown. Distal part of femora, second half of tibiae and tarsi dark brown.

Morphological characters: Body oval, rather broad, body length 1.86–2.60 mm. Dorsal and apical hairs slender and acute, longest hair 1.0–1.2 times as long as the basal diameter of antennal segment III. Rostrum reaching to the third pair of coxae, ultimate rostral segment 0.12–0.15 mm, 0.80–1.03 times as long as the hind tarsal joint II, bearing 7–9 secondary hairs. Total length of the antennae 2.57–2.98 mm, antennal segment III 0.66–0.86 mm, IV 0.60–0.74 mm, V 0.43–0.55 mm, VI 0.19–0.22 + 0.45–0.52 mm. Processus terminalis 2.13–2.63 times as long as the basal part of antennal segment VI, with 3–7 smooth hairs. Longest hair on antennal segment III 0.75 times as long as the basal diameter of the same segment, number of secondary rhinaria on it 7–19, rather variable in size and distributed over 40–60% of antennal segment III (see fig. 1 A). Secondary rhinaria in an alatiform specimen 29/30, covering the whole length of the segment. Siphunculi 0.58–0.74 mm, 0.28–0.38 times the body length, reticulated area covering the distal 30–40%. Cauda slender, acuminate, 0.37–0.52 mm, nearly 3 times as long as the width of the base, 60–80% of the length of the siphunculi, bearing 11–18 hairs, longest hair on cauda (whole series) 0.075 mm, 1.7 times as long as basal diameter of antennal segment III. Abdominal segment VIII with 4 and VII with 6 hairs. First tarsal joints bearing 3:3:3 hairs. Second tarsal joint of hind legs 0.14–0.17 mm, 0.67–0.91 times as long as the basal part of antennal segment VI.




Fig. 1: *Uroleucon tortuosissimae* sp. n., apterous viviparous female. A = antennal segment III, B = antennal segment IV, C = antennal segment V, D = antennal segment VI, E = siphunculus, F = cauda, G = head.




Fig. 2: *Uroleucon tortuosissimae* sp. n., apterous viviparous female. A = hind tarsus, B = ultimate rostral segment.

Tab. 1: *Uroleucon (Uroleucon) tortuosissimae* sp. n., apterous viviparous females. Measurements in mm. Urs. = ultimate rostral segment, H. tars. = hind tarsus, Pt. = processus terminalis, B = basis.

No	Body, length	Antennae, tot. leng.		Siphunculi, length		Cauda	Urs.	H. tars. II, length		Antennal segments, length												Rhin. on III		Index Pt./B VI		Index Siph./Body	
		l	r	l	r			l	r	l	r	III	IV	V	VI		l	r	l	r	l	r					
1	2,60	2,98	2,94	0,732	0,744	0,520	0,148	0,186	0,136	0,863	0,843	0,744	0,719	0,472	0,488	0,205+0,473	0,208+0,482	13	12	2,31	2,32	0,28	0,29				
2	2,10	2,85	2,82	0,644	0,644	0,508	0,136	0,167	0,161	0,806	0,768	0,644	0,682	0,501	0,433	0,210+0,498	0,202+0,489	7	8	2,37	2,42	0,31	0,31				
3	2,29	2,86	-	0,707	0,719	0,471	0,149	0,149	0,149	0,770	0,769	0,657	0,709	0,508	-	0,198+0,506	-	16	15	2,56	-	0,31	0,31				
4	1,86	2,77	2,75	0,657	0,649	0,409	0,124	0,153	0,148	0,719	0,682	0,620	0,632	0,496	0,488	0,223+0,508	0,198+0,508	14	15	2,28	2,57	0,35	0,35				
5	2,29	-	-	0,682	0,694	0,496	0,131	0,161	0,161	-	0,768	-	0,649	-	-	-	-	-	15	-	-	-	0,30	0,30			
6	2,10	2,76	2,80	0,644	0,649	0,422	0,125	0,149	0,149	0,744	0,719	0,620	0,670	0,471	0,508	0,223+0,521	0,211+0,484	10	13	2,34	2,29	0,31	0,31				
7	2,17	-	2,74	0,707	0,682	0,434	0,149	0,149	0,144	-	0,769	-	0,682	-	0,446	-	-	0,188+0,471	-	12	-	2,51	0,33	0,31			
8	2,48	2,86	-	0,719	0,719	0,434	0,148	-	0,161	0,781	0,794	0,670	0,682	0,520	-	0,210+0,466	-	11	12	2,22	-	0,29	0,29				
9	1,92	2,60	2,57	0,595	0,582	0,372	0,124	0,153	0,149	0,682	0,657	0,595	0,620	0,471	0,434	0,210+0,471	0,210+0,447	15	14	2,24	2,13	0,31	0,30				
10	2,05	2,89	-	0,682	0,683	0,447	0,137	0,149	0,149	0,806	0,769	0,645	-	0,546	-	0,198+0,496	-	9	9	2,51	-	0,33	0,33				
11	1,92	-	2,88	0,732	0,719	0,459	0,149	0,153	0,161	0,768	0,769	-	0,694	-	0,508	-	-	0,198+0,508	15	13	-	2,57	0,38	0,37			
12	2,29	-	-	0,670	0,682	0,471	0,134	0,167	0,161	0,744	0,769	0,670	0,709	-	-	-	-	-	12	13	-	-	0,29	0,30			
13	2,17	2,84	2,81	0,744	0,744	0,447	0,137	0,167	0,162	0,770	0,768	0,682	0,620	0,508	0,496	0,198+0,488	0,198+0,484	12	14	2,46	2,44	0,34	0,34				
14	1,92	-	-	0,694	0,682	0,496	0,136	0,161	0,160	0,769	0,767	0,683	0,682	-	-	-	-	-	18	19	-	-	0,36	0,36			
15	1,93	2,68	-	0,632	0,620	0,494	0,136	0,149	0,161	0,806	0,794	0,670	0,657	0,471	0,496	0,223+0,508	0,198+	10	12	2,28	-	0,33	0,32				
16	1,86	2,96	2,94	0,657	0,670	0,434	0,149	0,144	0,149	0,794	0,781	0,688	0,682	0,520	0,496	0,198+0,520	0,211+0,521	16	14	2,63	2,47	0,35	0,36				

Tab. 2: *Uroleucon (Uroleucon) hymenocephali* sp. n., apterous viviparous females. Measurements in mm, abbreviations as in tab. 1.

No	Body, length	Antennae, tot. leng.		Siphunculi, length		Cauda	Urs.	H. tars. II, length		Antennal segments, length												Rhin. on III		Index Pt./B VI		Index Siph./Body	
		l	r	l	r			l	r	III	IV	V	VI		l	r	l	r	l	r							
1	3,22	3,00	3,09	0,781	0,769	0,620	0,150	0,180	0,174	0,806	0,769	0,637	0,682	0,521	0,608	0,223+0,540	0,236+0,546	29	30	2,42	2,31	0,24	0,24				
2	2,98	3,26	3,34	0,769	0,744	0,710	0,172	0,192	0,198	0,942	0,930	0,769	0,781	0,595	0,595	0,235+0,570	0,248+0,533	16	18	2,43	2,15	0,26	0,25				
3	2,73	3,40	3,45	0,893	0,905	0,682	0,160	0,186	0,180	0,930	0,918	0,776	0,830	0,628	0,657	0,230+0,598	0,233+0,583	29	29	2,60	2,50	0,33	0,33				
4	2,60	-	-	0,905	0,908	0,676	0,164	0,186	0,184	0,885	0,866	0,788	-	0,578	-	-	-	-	21	23	-	-	0,35	0,35			
5	2,67	3,25	3,39	0,918	0,930	0,692	0,170	0,174	0,181	0,893	0,930	0,756	0,802	0,585	0,632	0,211+0,568	0,229+0,556	26	24	2,69	2,43	0,34	0,35				

Taxonomic note: The morphological observation of *Uroleucon tortuosissimae* shows that this species resembles *U. scorzonerae* DAN. (DANIELSSON, 1973) collected from *Scorzonera humilis* L., but it differs from it mainly by the following characters: Cauda pale, secondary rhinaria only 7–19, distributed at most over 60% of the antennal segment III, first tarsal joints bearing 3 : 3 : 3 setae, no post-siphuncular sclerites.

Hostplant: *Scorzonera tortuosissima* BOISS. (Compositae = Asteraceae). This plant is found only in Iran and Afghanistan (RECHINGER, 1977).

Aphids collected from Saveh (Saleh-Abad, 1500 m), 140 km south-west of Tehran, on 25.5.1985. Types in the collection of the first author.

2. *Uroleucon (Uroleucon) hymenocephali* sp. n.

Apterous viviparous female (Figs. 3 and 4)

Colour in living specimens: Greyish brown, mat, cauda pale. Cleared specimens: Thorax and abdomen colourless without sclerites, postsiphuncular scler-




Fig. 3: *Uroleucon hymenocephali* sp. n., apterous viviparous female. A = head, B = antennal segment III, C = antennal segment IV, D = antennal segment V, E = antennal segment VI, F = cauda, G = siphunculus.




Fig. 4: *Uroleucon hymenocephali* sp. n., apterous viviparous female. A = ultimate rostral segment, B = hind tarsus.

ites not developed, head and coxae brownish. Distal half of femora, whole length of tibiae and tarsi as well as the siphunculi and the last two segments of the rostrum dark brown to blackish. Basal part of third and basal half of fourth antennal segment pale, the remainder parts of the antennae brownish to brown. Cauda pale, genital plate brownish.

Morphological characters: Body oval, total length 2.60–3.22 mm. Dorsal and apical hairs slender and acute, longest hair 1.0–1.2 times as long as basal diameter of antennal segment III. Rostrum reaching to the third pair of coxae, ultimate rostral segment 0.15–0.17 mm, 0.83–0.98 times as long as the hind tarsal segment II, bearing 6–7 secondary hairs. Total length of the antennae 3.00–3.45 mm, 0.93–1.27 times the body length. Antennal segment III 0.77–0.94 mm, IV 0.64–0.83 mm, V 0.52–0.66 mm, basal part of the VI. antennal segment 0.21–0.25 mm, 1.24–1.57 times as long as the ultimate rostral segment and 1.16–1.36 times as long as the hind tarsal segment II. Processus terminalis 0.53–0.60 mm, 2.15–2.69 times as long as the basal part of antennal segment VI, bearing 5–7 hairs. Antennal segment III bearing 16–30 secondary rhinaria, distributed in an irregular manner over nearly the whole length of the segment (see fig. 3 B). Longest antennal hair on segment III 0.75–0.85 times as long as the basal diameter of this segment. Siphunculi straight, 0.74–0.93 mm, 0.24–0.35 times the body length and 1.05–1.34 times as long as the cauda, 0.33–0.45% of the apical part reticulated. Cauda dirkshaped, pale, mostly with

a distinct constriction, 0.62–0.71 mm, 0.19–0.26 times the body length and 0.75–0.95 times as long as the siphunculi, bearing 18–26 hairs. Longest hair on the cauda (whole series) 0.09 mm, about 1.75 times as long as the basal diameter of antennal segment III. Hind tarsal segment II 0.17–0.20 mm, first tarsal segments bearing 5 : 5 : 5, exceptionally 5 : 3 : 3 setae. Abdominal segment VIII with 4 and VII with 6 hairs.

Taxonomic note: This species differs from that mentioned above in having a greater body length and a higher number of secondary rhinaria. In addition the whole length of the tibiae is dark brown to blackish and the first tarsal segments mostly bear 5 : 5 : 5 setae.

Hostplant: *Hymenocephalus rigidus* JAUB. & SPACH. (Compositae = Asteraceae). This plant is only found in Iran (WAGENITZ, 1980).

Aphids collected from Saveh (Saleh-Abad, 1500 m), 140 km south-west of Tehran, on 25.5.1985. Types in the collection of the first author.

REFERENCES

- DANIELSSON, R. 1973. *Uroleucon (Uromelan) scorzonerae*, a new aphid species from Sweden (Hom. Aphididae). *Ent. scand.* 4: 249–253.
- EASTOP, V. F. 1985. Key to the Middle Eastern species of *Uroleucon* MORDVILKO (Aphididae: Homoptera). *Syst. Entomol.* 10: 395–404.
- RECHINGER, K. H. 1977. *Scorzonera tortuosissima* BOISS. In: RECHINGER, K. H. (Herausg.): *Flora iranica Lfg. 122: Compositae II – Lactuceae*, pp. 49–50. Akademische Druck- und Verlagsanstalt, Graz.
- WAGENITZ, G. 1980. *Hymenocephalus rigidus* JAUB. & SPACH. In: RECHINGER, K. H. (Herausg.): *Flora iranica Lfg. 139: Compositae III – Cynareae (2)*, p. 421. Akademische Druck- und Verlagsanstalt, Graz.

(received June 10, 1987)