**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft =

Bulletin de la Société Entomologique Suisse = Journal of the Swiss

**Entomological Society** 

Herausgeber: Schweizerische Entomologische Gesellschaft

**Band:** 60 (1987)

**Heft:** 3-4

Artikel: An aberrant new genus and new species of Carpocorini Stål (Hemiptera

: Pentatomidae : Pentatominae) from Pakistan

Autor: Ahmad, Imtiaz / Rana, Nazeer Ahmad

**DOI:** https://doi.org/10.5169/seals-402284

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 28.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

An aberrant new genus and new species of Carpocorini Stål (Hemiptera: Pentatomidae: Pentatominae) from Pakistan<sup>1</sup>

# IMTIAZ AHMAD & NAZEER AHMAD RANA<sup>2</sup>

Department of Zoology-Entomology, University of Karachi.

A new genus to accomodate a new species from Pisheen in Baluchistan is described with special reference to their metathoracic scent gland ostioles and male and female genitalia. Their relationships within Pyrrhocoridae is also briefly discussed.

During various expeditions to northern areas of Pakistan from 1967 through 1983, five unique specimens were collected on a wild bush from Pisheen in Baluchistan. These certainly belong to the tribe Carpocorini Stål (Ahmad 1979) for they possess the characters noted presently under the section "comparative note" and "relationships". However, they appear isolated in the entire tribe Carpocorini in their simple sickleshaped paramere and much abbreviated metathoracic scent gland ostiolar complex. These are presently described as a new genus and new species with reference to their metathoracic scent gland ostiole and male and female genitalia. The new genus is named *Qadriana* in honour of Late Professor Dr. M. A. H. Qadri, University of Karachi in acknowledgement of his valuable contributions in the field of systematics and biology of Hemiptera. For the dissections of male and female genitalia, and for measurements and diagrams the conventional procedures specially those used by the present authors (1981) were generally followed.

## Oadriana Gen. nov.

General shape: Body elongately ovate; longer than broad, broadly rounded in the posterior abdominal half; pronotum medially raised, much deflected in the anterior half.

Head: Distinctly longer than broad; anterior of head excluding eyes distinctly shorter than remainder of head; paraclypei not passing beyond clypeus, with their outer margins entire, distinctly sinuate infront of eyes, accuminate and slightly reflexed in the anterior half; clypeus distinctly prominent; labium short not passing hind coxae; antenniferous tubercles visible from above, antennae with second segment distinctly longer than the 3rd.

Thorax: Pronotum much deflected in the anterior half, slightly shorter than head length with anterior margin broader than the width of head across the eyes, anterior anlges subacute, lateral margins entire, humeral angles broadly

<sup>&</sup>lt;sup>1</sup> Financially supported by PARC/USDA Research Project No. FG-Pa-361 (PK-SEA-155)

<sup>&</sup>lt;sup>2</sup> Government Degree College, Liaquatpur, Punjab, Pakistan.

rounded, mesosternum obsoletely sulcate; metathoracic scent gland ostioles indistinct with peritremes extremely reduced.

Abdomen: Connexiva distinctly visible at repose; in females 8th spiracles located on ventro-posterior margin of the 8th paratergites, 7th abdominal sternum with ventro-posterior margin distinctly concave, lateral angles subacute.

Male genitalia: Pygophore much broader than long, proctiger somewhat quadrangular with rounded apex; paramere sickle-shaped with elongate blade, inner margin serrate; inflated aedeagus with pair of dorsal membranous conjunctival appendages, vesica remarkably long.

Female genitalia: 1st gonocoxae placed wide apart, somewhat triangular; 2nd gonocoxae broader than long; 9th paratergites elongated, reaching margins of fused 8th paratergites; spermathecal bulb distinct but small with distal and proximal flanges prominent; pump region tubular and distinct.

Comparative note: The genus appears quite abberrent in the entire tribe Carpocorini in having body obovate, short, scutellum distinctly broad at apex, metathoracic scent gland ostioles reduced with poorly marked long impression of peritremes and parameres sickle-shaped with irregulary, serrate inner margins.

Colour: Body uniformly dull straw-coloured with greenish sheen; antennae yellowish brown, entire body with uniform reddish brown punctures; membrane of hemelytra hyaline, yellowish brown; eyes dark ochraceous with blackish tinge; ocelli dark pink.

Head: Distinctly longer than broad; paraclypei not enclosing clypeus, slightly passing beyond clypeus; latter prominently raised at apex; length of head equal to length of pronotum, antennae with apical segment longer than 3rd, length of segments. I  $0.3 \, \text{mm}$ , II  $0.5 \, \text{mm}$ , III  $0.45 \, \text{mm}$ , IV  $0.67 \, \text{mm}$ , V0.72 mm, antennal formula 1 < 3 < 2 < 4 < 5; labium with ¼ of the basal segment extending beyond bucculae, reaching to hind coxae, length of segments, I  $0.57 \, \text{mm}$ , II  $0.95 \, \text{mm}$ , III  $0.55 \, \text{mm}$ , IV  $0.47 \, \text{mm}$ , labial formula 4 < 3 < 1 < 2 <; length of anterior portion of head excluding eyes  $0.7 \, \text{mm}$ , length posterior of head including eyes  $0.9 \, \text{mm}$ , width  $1.5 \, \text{mm}$ , interocular distance  $0.9 \, \text{mm}$ , interecellar distance  $0.7 \, \text{mm}$ .

Thorax and abdomen: Pronotum with anterior angles subacute, broadly rounded, anterolateral margins entire, on anterior half two dark ochraceous callosities present, nearly 3x broader than length, length 1.6 mm, width 4.0 mm, scutellum with apical half portion, broader but gradually tappering towards apex with two black impressions at basal margin in the middle, distinctly longer than broad, length 2.7 mm, width 2.3 mm; metathoracic scent gland ostiole (Fig. 2) indistinct, peritreme extremely reduced, evaporatoria quite large but less rugulose; distance base scutellum-apex clavus 1.2 mm; apex clavus-apex corium 1.5 mm, apex corium-apex abdomen including membrane 1.2 mm, connexiva exposed at repose; female's posterior margin of 7th abdominal sternum broadly concave with lateral angles subacute. Total length ♂ 7.1 mm, ♀ 8.7 mm.





Fig. 1. Quadriana pisheenensis sp.nov.

Male genitalia: Pygophore (Figs. 4-8) having dorsomendian surface broadly concave, anterolaterally much broad with anterior angles roundly projected, dorsolateral inner processes knoblike, ventroposterior margins sinuate; paramere (Fig. 5) sickleshaped, pointed at apex with outer margin convex, inner margin concave with inner dentation, blade slightly showing lamination; aedeagus with (Figs. 6-8) a pair of dorsal membranous conjunctival appendages, semisclerotized at apex, tubular vesica very much dilated before the apical curved tubular half, gonopore with opening located dorsad.

Female genitalia: First gonocoxae (Fig. 10) placed wide apart, somewhat triangular with their inner angles rounded; 8th paratergites triangular, never reaching posterior margin of 1st gonocoxae; 9th paratergites elongated, placed wide apart; 2nd gonocoxae quadragular with anterior margin straight; spermathecal bulb small but distinct without processes, distal and proximal flanges distinct, pump region uniformly tubular (Fig. 9).

Material examined: Holotype ♂, Baluchistan, Pisheen, wild bush; 14-10-1978, University of Karachi. Paratypes 3 ♂, 1 ♀ with the same data as holotype, in the above museum, at Unites States National Museum, Washington, D.C., USA, at Natural History Museum, Basel, Switzerland and in Ahmad's collection.



Figs. 2–10 Quadriana pisheenensis sp.nov. 2, scent gland ostiole; 3, pygohpore, dorsal view; 4 pygophore, ventral view; 5, paramere; 6, aedeagus, dorsal view; 7, aedeagus, ventral view; 8 aedeagus, lateral view; 9, spermatheca; 10, female terminalia. per., peritreme; ev., evaporatoria; o. ostiole; 11th abd. seg., 11th abdomina segment; dl.l. dorsolateral lobe; p., paramere; in.pl., innet plate; proc., proctiger; dis.spd. distal spermathecal duct; prx.f. proximal flange; sp.p., spermathecal pump; dis.f., distal flange; spb. spermathecal bulb; m.dl. median dilation; scl. md., sclerotized me dian duct; prx. spd., proximal spermathecal duct; gp., gonopore; ves., vesica; dmc. app., dorsal membranous conjunctival appendage; th., theca; piv., pivot; bp., basal plate; inf.p., inflatory pump gp., gonopore; 1st gox., 1st gonocoxa; tr., triangulin; 9th. pt., 9thparatergite; 2nd gox., 2nd gonocoxae; 8th pt., eighth paratergite.

## Relationships of the new taxa

Quadriana pisheenensis, new species, possesses the characters of scutellum which is distinctly longer than broad with apical portion narrow, tapering near middle and sides of prontum entire, which are also shared by the species of Brachynema Muslant& Rey, Cappaea Ellner, Carpocoris Kolenati, Codophila Muslant& Rey, Croantha Stål Dolycoris Muslant& Rey, Halyomorpha Mayr, Holcostethus Fieber, Manoriana Ahmad & Kamaluddin, Mormidella Horv. and Palomena Muslant& Rey found in Pakistan.

GROSS (1976) although considered the most important characteristic feature of his *Carpocoris* group, the strong and thick F-shaped parameres but in the present new taxon the parameres are simple sickleshaped with irregular serrations on inner margins, showing little affinity towards any carpocorine known from Pakistan and Azad Kashmir (Ahmad 1979, 1980). The present taxon was also found isolated in the tribe Carpocorini in having the characters of scutellum distinctly borader in the apical half, narrow at apex and metathoracic scent gland ostioles reduced with ill-marked long impression of peritremes and other characters noted in the comparative note of the genus. The reduced ostiolar penitreme and the simple sickleshaped parameres also relate it to the members of some extant Strachiini Stål.

#### REFERENCES

Ahmad, I. 1979. A revision of the check-list of Coreidae and Pentatomidae of the superfamilies Coreoidea and Pentatomoidea (Heteroptera: Pentatomomorpha) from Pakistan with phylogenetic considerations *Karachi ent. Soc. Suppl. 4* (1): 1–113.

AHMAD, I. 1980. Insect fauna of Pakistan and Azad Kashmir: Some groups within the order Hemiptera. *Proc. 1st. Pakistan. Congr. Zool. A:* 115–155.

AHMAD, I. & RANA, N. A. 1981. A new genus and a new species of Antestrini DISTANT (Pentatominae) and their Relationships. *Pakistan. J. Zool. 13* (1&2): 45-50.

GROSS, G. F. 1976. Plant-feeding and other bugs (Hemiptera) of South Australia-Heteroptera. Part II. Government Printer, South Australia.

(received October 3, 1987)

