

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 61 (1988)

Heft: 1-4

Artikel: Strategy for the integrated control of pear pests by the use of an insect
growth regulator as a key element

Autor: Sechser, B. / Engelhardt, M.

DOI: <https://doi.org/10.5169/seals-402311>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Strategy for the integrated control of pear pests by the use of an insect growth regulator as a key element

B. SECHSER & M. ENGELHARDT

CIBA-GEIGY Ltd., Agricultural Division, CH-4002 Basel, Switzerland

A new insect growth regulator, CGA 45 128, was used for three consecutive years against the last instar larvae of *Adoxophyes orana* F.v.R. in spring and caused a fruit damage reduction to 9, 0.8 and 1% respectively. Typical deformations for this type of compound could be observed on all sampled larvae. CGA 45 128 was fully selective to the anthocorids which helped to keep psyllid populations at an acceptable level.

INTRODUCTION

The pear production in Europe is threatened by two pests mainly: the fruit tortrix moth *Adoxophyes orana* F.v.R. and several psyllid species, in particular *Psylla piri* L. While the tortricid can be well controlled, psyllids caused serious problems in several regions over the past decade (IOBC, 1983).

One of the reasons for these problems was insufficient coverage of big pear trees, resulting in sublethal dosage rates, which initiate a quick build-up of resistance with psyllids. In addition, certain insecticides are deleterious to anthocorids which are the most important group of psyllid predators.

New strategies in pear pest control aim at the introduction of more selective insecticides and a shift of the treatments against *A. orana* from the summer generation to the overwintering larvae. A growth regulator type of compound with activity against the last instars would suit such a system very well because the damage to the foliage in springtime by this leafroller is acceptable. Anthocorids are not vulnerable to this type of substance because they are only present in the adult stage at this time of the year. An insect growth regulator (IGR) with the code number CGA 45 128 displays the required properties. To test its suitability for the suggested system was the purpose of the present study.

MATERIALS AND METHODS

The experiments were carried out in 1985, 1986, and 1987 in pear orchards in the Swiss Rhone valley. The locations and relevant treatments are listed in Table 1.

A. orana was treated with CGA 45 128. The systematic chemical name is N-ethyl O[2-(4-phenoxyphenoxy)-ethyl] carbamate.

It has morphogenetic action on the last larval instars of this leafroller species, resulting in deformed larvae, pupae and adults. It was compared to check orchards where only psyllids were controlled. All pesticides were applied with a turbo sprayer at 1400 l water/ha against overwintering pests and at 400 l/ha during the growing season.

Table 1. Treatment list in pear insect control trial

	Trial site	Pest	Active ingredient (a.i.) and formulation	Rates g a.i./100 l	Dates of treatment
1985	Riddes A 0.5 ha	PP ^{a)}	IGR ^{c)} EC 100 + mineral oil (50 %)	50 + 300	April 3 / 13
		AO ^{b)}	CGA 45 128 EC 250	5	May 16 / 27
		PP	Amitraz EC 200	50	June 27
	Riddes B 0.3 ha	PP AO	DNOC SC 500 No treatment	375	Febr. 26
1986	Riddes A 0.5 ha	PP AO	DNOC SC 500 CGA 45 128 EC 100	375 1.25 & 2.5	March 6 May 12 / 23
	Saxon B 0.5 ha	PP AO	Mineral oil (99 %) No treatment (?)	200	March
1987	Riddes A 0.5 ha	PP AO	DNOC SC 500 & mineral oil (50 %) CGA 45 128 EC 100	375 + 350 2.5	March 23 & April 2/24 May 8/20
	Saxon A 0.4 ha	PP	Mineral oil	350	April 7
		AO	CGA 45 128 EC 100	2.5	May 9/25
		PP	Amitraz EC 200	50	July 7
	Saxon B 0.5 ha	PP AO	Mineral oil (99 %) & soap No treatment	200	March & several times

a) PP-*Psylla piri* b) AO-*Adoxophyes orana*

c) Insect growth regulator 5-(4-phenoxyphenoxy-l-pentin)

The damage caused by all pests was rated at harvest by counting all fruits from 4 trees per treatment.

The development of psyllids and anthocorids during the growing season was followed up by visual inspections on 100–400 relevant organs (inflorescences, terminals) but the figures indicated here apply to 100 organs.

RESULTS AND DISCUSSION

The fruit tortrix moth was well kept under control in all three years (Table 2). Larvae treated with CGA 45 128 sometimes tend to stay longer on the foliage which may result in some nibbling damage on the newly formed fruits. When 60 *A. orana* last instar larvae were caged in 1986 on treated branches, all of them demonstrated the deformations typical for growth regulators. The total leafroller

Table 2. Damage rating (%) of *Adoxophyes orana* and *Psylla piri* on pear fruits at harvest

Trial sites		Adoxophyes		Psylla		Other pests a)
		spring	summer	slight	severe	
1985	Riddes A (IGR)	7	2	22	0	2
	Riddes B (Check)	No rating at harvest				
1986	Riddes A (IGR)	0.4	0.4	10	0.2	2
	Saxon (Check)	2	2	10	2	5
1987	Riddes A (IGR)	0.2	0.8	6	0	4
	Saxon A (IGR)	0	0.2	15	0.4	2
	Saxon B (Check)	1	1	14	1	6

a) Noctuid Larvae, mirids, San José scale, sawflies

damage was commercially acceptable in the first year since all these fruits went for distillation. The leafroller population had decreased in the following two years in the same orchard to a negligible level which is a general experience with this type of compound (CHARMILLOT & BLASER, 1985).

A strong reduction of *Psylla piri* could be achieved by the proper timing of another growth regulator or DNOC against overwintering adults (Table 3). Re-surfing populations in 1985 and 1987 were kept under control during the growing season by applying amitraz against the eggs. The sooty aspect of mould growing on honey dew excretions of psyllid larvae was only slight (Table 2).

Complementary selective control measures were pirimicarb against aphids, diflubenzuron against noctuid larvae, and a mixture of penconazol and captan for the control of apple scab and mildew.

Anthocoris nemoralis F. and *A. nemorum* L. were present all through the growing season except in general from May to mid-June (Table 3). This is a critical period and most often the moment when corrective treatments against psyllids become necessary. A selective product such as CGA 45 128 allows the full survival of anthocorids at the beginning of the growing season keeping the psyllid population at a low level.

The application of a single compound against both important pear insect pests would be the most convenient control method for the fruit grower. This

Table 3. Seasonal development of *Psylla piri* and anthocorid larvae and adults by visual inspection in pear orchards (% occupation)

Trial site			March	May	June			July				Aug.		
			2	28	10	14	28	3	10	18/19	25	2	8	17
1985	Riddes A (IGR)	ppa)	49	37	43	98	82	35	55	11	16	70	6	11
		AN ^{b)}	2	0	0	1	0	5	6	10	14	8	32	27
	Riddes B (Check)	PP	1	6	68	12	42	44	14	100	98	-	10	10
		AN ^{b)}	0	0	0	0	0	0	0	4	48	10	-	0
			May		June			July						Aug.
			22	28	10/12	16	24/25	2/3	6	8/10	14/16	22/24	29/30	5/12
1986	Riddes A (IGR)	PP	1	3	7	5	8	18	5	10	42	60	32	4
		AN ^{b)}	0	0	0.3	0.3	1	1	5	1	2	8	17	5
	Saxon B (Check)	PP	4	25	21	-	32	17	-	20	17	21	22	14
		AN ^{c)}	0.3	0	2	-	10	10	-	4	3	18	10	5
			May		June			July						Aug.
			20	26	2	11/12	17	23/24	6	15	23	29	5	19
1987	Rides A (IGR)	PP	0	4	4	22	11	27	35	52	66	50	42	21
		AN ^{b)}	0	0	0	1	2	2	3	3	16	3	13	10
	Saxon A (IGR)	PP	14	7	18	14	16	39	90	99	33	22	54	10
		AN ^{b)}	0	0	1	3	0	5	4	6	10	7	14	6
	Saxon B (Check)	PP	83	88	74	75	98	44	100	84	57	25	40	17
		AN ^{c)}	0	0	1	19	21	7	11	9	11	7	12	7

a) PP-*Psylla piri* b) AN-*Anthocoris nemoralis* c) *Anthocoris nemorum*

worked very well in the past with organophosphates until the appearance of serious resistance problems with psyllids in northern Italy, which were caused by many successive generations and sublethal rates of insecticides in the crown canopy of traditionally high grown pear trees (PICO & PICO, 1978). The problem was aggravated by the use of pyrethroids. Psyllid control became difficult in parts of Switzerland (STAEUBLI, 1983), France (BASSINO, 1983), and the Netherlands (BLOM *et al.*, 1985). Pyrethroids are detrimental to anthocorids as most important natural enemies of psyllids (VEZ, 1978; HAGLEY & SIMPSON, 1983). Growth regulators such as CGA 45 128 or phenoxycarb allow the control of *A. orana* already in early springtime, but are harmless to anthocorid adults which are the only stage of this predator present at this time of the year (MONNET, 1986, REEDE *et al.*, 1985). This is one of the rare cases of a really efficient insect pest antagonist. Its mid-season absence for several weeks from pear orchards can cause a critical resurgence of psyllid populations. Fortunately, there is amitraz available as a selective compound for corrective treatments as long as there is no resistance to this product. It would be interesting to know where anthocorids stay during the months of May and June and whether it would be possible to manipulate the envi-

ronment in such a way (e.g. by intercrop planting of attractive host plants as suggested by FYE, 1983) to make pear orchards attractive to anthocorids also during this period.

CONCLUSIONS

In these experiments the insect growth regulator CGA 45 128 proved its good activity against leafrollers and its full selectivity on anthocorids. Therefore, by the use of such a type of compound these predators could display their full potential as biological control agents of psyllids.

ACKNOWLEDGEMENT

We appreciate the help by T. HOPPE who did most of the visual inspection in 1985 and to M. A. RUZETTE for the damage rating.

RÉSUMÉ

Stratégie de la lutte intégrée contre des ravageurs sur poiriers par l'emploi d'un régulateur de croissance comme élément principal – Un nouveau régulateur de croissance, CGA 45 128, a été utilisé en deux traitements contre le dernier stade larvaire d'*Adoxophyes orana* en printemps. Les fruits endommagés ont été ramenés au cours des trois années à 9, 0.8 et 1% respectivement. Des malformations typiques ont été constatées sur toutes chenilles retrouvées sur les poiriers traités. Aucune action négative du CGA 45 128 n'a été observé sur les anthocorides qui ont largement contribué à maintenir les psylles à un niveau tolérable.

REFERENCES

- BASSINO, J.P. 1983. Evolution et aménagement de la lutte chimique en verger de poiriers. *Bull. IOCB/WPRS VII* (5): 281–85.
- BLOM, J., VAN DER DRUKKER, B. & BLOMMERS, L.H.M. 1985. The possible significance of various groups of predators in preventing pear psylla outbreaks. *Meded. Fac. Landbouwwet. Rijksuniv. Gent* 50: 419–24.
- CHARMILLOT, P.J. & BLASER, C. 1985. Le fenoxycarb, un régulateur de croissance d'insectes homologué contre la tordeuse de la pelure *Adoxophyes orana* F.v.R. *Revue suisse Vitic. Arboric. Hortic.* 15: 195–201.
- FYE, R.E. 1983. Cover crop manipulation for building pear psylla (Homoptera: Psyllidae) predator populations in pear orchards. *J. Econ. Entomol.* 76: 306–10.
- HAGLEY, E.A.C. & SIMPSON, C.M. 1983. Effect of insecticides on predators of the pear psylla, *Psylla pyricola* (Hemiptera: Psyllidae), in Ontario. *Can. Entomol.* 115: 1409–14.
- IOBC, 1984. Integrated control of pear psyllids. Toulouse, 27.–29.9.83. *Bull. IOCB/WPRS VII* (5), 388 pp.
- MONNEZ, Y. 1986. Contre les tordeuses de la pelure: Une substance perturbant la métamorphose des insectes. *Phytoma* 375: 28–30.
- PICO, D. & PICO, A.M. 1978. Nella difesa delle pomacee con la «lotta complementare» anche la psilla del pero (*Psylla pyri* L.) posta sotto il controllo del suo naturale «complesso biotico antagonista». *Notiziario delle Malattie delle Piante* 98/99: 101–17.
- REEDER, R.H. DE, ALKEMA, P. & BLOMMERS, L.H.M. 1985. The use of the insect growth regulators fenoxycarb and epofenonane against leafrollers in integrated pest management in apple orchards. *Entomol. Exp. Appl.* 39: 265–72.
- STAEUBLI, A. 1983. Aménagement d'une lutte raisonnée contre le psylle du poirier (*Psylla pyri* L.) en Suisse particulièrement en Valais et dans le Bassin lémanique. *Bull. IOCB/WPRS VII* (5): 258–64.
- VEZ, F. 1978. Etude des antagonistes du psylle de poirier (*Psylla pyri* L.) et en particulier du prédateur *Anthocoris nemoralis* (F.) dans les vergers de Suisse romande. Travail de diplôme, ETH Zürich, 98 pp.

(received June 10, 1988)