

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 63 (1990)

Heft: 1-2

Artikel: Drunella andaluciaca sp.n. (Ephemeroptera, Ephemerellidae) from Spain

Autor: Kazanci, Nilgün

DOI: <https://doi.org/10.5169/seals-402393>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 03.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Drunella andaluciaca sp.n. (Ephemeroptera, Ephemerellidae)
from Spain

NILGÜN KAZANCI

Hacettepe Üniversitesi, Fen Fakültesi Biyoloji Bölümü
Hidrobiyoloji Beytepe – Ankara, Türkiye

This paper includes the nymphal description of a new *Drunella* species from Spain, and a nymphal key for the Western Palaearctic *Drunella* species.

INTRODUCTION

In the Western Palaearctic Region the genus *Drunella* has been represented by *D. paradinasi* GONZALES DEL TANAGO et GARCIA DE JALON, 1983 from Spain, *D. euphratica* KAZANCI, 1987 and *D. karia* KAZANCI, 1989 from Turkey. The new species *Drunella andaluciaca* sp.n. is another representative of the genus *Drunella* in the Western Palaearctic Region.

DESCRIPTION

Drunella andaluciaca sp.n.

Mature nymph (female)

Length of body: 4–8 mm. Length of cerci: 4–6 mm.

Head: Head with distinctive tubercles, located in the middle of the vertex and covered with short setae (Figs. 1 and 2). Genae with slight projections (Fig. 1). Head yellowish brown with light brown spots. Second segment of antennae brown, remainders are yellow.

Mouth parts (Figs. 4–8): Terminal segment of maxillar palpus with four spines and second segment with one lateral seta (Fig. 5). Glossae a little longer than broad, paraglossae broad and slightly pointed at the apex, anterolateral edges of submentum pointed (Fig. 8). Labrum and hypopharynx are drawn in Figs. 3 and 7. The hypopharynx of *D. euphratica* (Fig. 9) is given for comparison.

Thorax: Pro- and mesothorax light brown with brown patterns. Pronotum with projections on the anterolateral edges. Pronotum with a single low median and low paired submedian tubercles with low lateral ridges. Mesonotum with a single low anteromedian, paired median, and single posterior tubercles between wing pads (Figs. 10 and 11).

Legs: All legs light brown without distinctive patterns, dorsal margin of femora without long hairs, insertion of stout spines on femora with slight protuberances (Figs. 12 and 13), tarsal claws with 9–11 denticles (Figs. 14 and 15).

Abdomen: Tergites 3–10 light brown median pattern (Fig. 16) and sternites with lateral stripes and median small rounded spots. Dorsal tubercles blunt and covered with spines between tergites 5–7 (Figs. 16 and 17). Whole tergite surface with mushroom-shaped structures without chagrin. Spines on the tergite margin stout and covered with hairs (Fig. 18). Cerci light brown without annulation and with long and thick spines on the posterior margin of each segment. Imago unknown.

Affinities: This new species belongs to the genus *Drunella* according to nymphal characteristics (ALLAN & EDMUNDS, 1962). *D. andaluciaca* sp.n. belongs to the *D. grandis* EATON group, and it seems to be similar to *D. paradinasi* GONZALES DEL TANAGO & GARCIA DE JALON, 1983 from Spain and *D. euphratica* KAZANCI, 1987





Fig. 1–8. *Drunella andaluciaca* sp.n.; Fig. 9. *Drunella euphratica* KAZANCI, 1987.
1. Head, 2. Head in lateral view, 3. Labrum, 4. Maxilla, 5. Maxillary palpus, 6. Right mandible, 7. Hypopharynx, 8. Labium, 9. Hypopharynx of *D. euphratica*.



Figs. 10–19. *Drunella andaluciaca* sp.n.; 10. Thorax in dorsal view, 11. Thorax in lateral view, 12. First leg, 13. Third leg, 14. Tarsal claw of first leg, 15. Tarsal claw of third leg, 16. Abdomen in dorsal view, 17. Abdomen in lateral view, 18. Structure of tergite surgave and marginal spines, 19. First gill.

from Eastern Turkey. The new species differs from *D. paradinasi* and *D. euphratica* in the body coloration and patterns, in the body tubercles and ridges, and in the shape of the mouth parts. The structure of the tergite surface is similar to that of *D. euphratica*; by the lack of star-like hairs it differs from *D. paradinasi*.

Material: Holotype and 15 paratypes, nymphs. Rio Trevelez 1600 m. (Granada, Spain), 28.7.1989.

KEY FOR THE NYMPHS OF *DRUNELLA* SPECIES FOR THE WESTERN PALAEARCTIC REGION

1. Head tubercles located near the base of compound eyes, gena without protuberance, prothorax narrow, a light and slender species, legs with distinctive dark spots, femora slender (see KAZANCI, 1989, Figs. 1, 9, 11, 12) *karia*
- Head tubercles located in the middle of the vertex, gena with slight protuberance (Fig. 1), prothorax shield-shaped (Fig. 10), dark and stout species, legs without distinctive dark spots, femora broad (Figs. 12 and 13) 2
2. Whole body is covered, with star-like hairs (see STUDEMANN & TOMKA, 1987, Figs. 2 and 3) *paradinasi*
- Body without star-like hairs 3
3. Abdominal tubercles are small almost equal in size in different segments and present in segments 3–9 (Fig. 16) *andaluciaca*
- Abdominal tubercles are well developed, but small between segments 2–4 and almost equal in size between segments 5–9 (see KAZANCI, 1987, Figs. 8, 9) *euphratica*

REFERENCES

ALLEN, R.K. & EDMUNDS, G.G. 1962. A revision of the genus *Ephemerella* (Ephemeroptera) V. The subgenus *Drunella* in North America, *Misc. Publ. Entomol. Soc. Amer.*, 3: 147–179.

GONZALES DEL TANAGO, M. & GARCIA DE JALON, D. 1983: New Ephemerellidae from Spain (Ephemeroptera). *Aquatic Insects*, 5 (3): 147–156.

KAZANCI, N. 1987. New *Drunella* (Ephemeroptera; Ephemerillidae) species from Turkey. *Mitt. Schweiz. Ent. Ges.*, 60: 379–382.

KAZANCI, N. 1989. Second species of genus *Drunella* (Ephemeroptera; Ephemerellidae) *Drunella karia* n.sp. from Turkey. *Hydrobiologia* (in print).

STUDEMANN, D. & TOMKA, I. 1987. Contribution to the study of European Ephemerellidae (Ephemeroptera) I. Completion of description of three endemic Iberian species. *Mitt. Schweiz. Ent. Ges.* 60: 361–378.