

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 65 (1992)

Heft: 3-4

Artikel: Notes on host plant, larval features and life history of *Cephalcia hartigii*
(Bremi) (Hym., Pamphiliidae)

Autor: Battisti, A. / Cescatti, A.

DOI: <https://doi.org/10.5169/seals-402493>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Notes on host plant, larval features and life history of *Cephalcia hartigii* (BREMI) (Hym., Pamphiliidae)

A. BATTISTI & A. CESCATTI¹

Istituto di Entomologia agraria, Via Gradenigo 6, I-35131 Padova

Cephalcia hartigii is a rare species whose natural history and host plant were unknown. During a field study on the web-spinning sawflies in northern Italy, 16 adults (14 males and 2 females) of this species were collected with different kinds of traps. Most of them were found on *Abies alba*. The catch of 2 mature larvae falling to the ground under a fir and the following findings of 2 typical pamphiliid larval shelters, gave evidence that *A. alba* is a host plant of *C. hartigii*. Information is given on flight period, egg laying, larval feeding and development. The main morphological characters of the mature larva are briefly described. Most of the records of *C. hartigii* reported in literature match well with the distribution pattern of *A. alba* and *A. nordmanniana*. Considering the fact that some records are not clearly referable to fir stands, the possibility that *C. hartigii* may also live on other host plants cannot be excluded.

Keywords: Web-spinning sawfly, *Abies* spp., distribution.

INTRODUCTION

Among the European species of the *Cephalcia* PANZER, *C. hartigii* (BREMI) is considered a rare species, whose biology and host plant are unknown, as reported by BENEŠ (1976) in his revision of this genus. According to BENEŠ, its distribution pattern (central and southern Europe, North Africa and the Caucasus) and the fact that the holotype was collected on *Pinus sylvestris* L. (BREMI, 1849) suggests that the host plants are likely to be several species of *Pinus*.

Taxonomically, *C. hartigii* differs strongly from the other species of the genus, sharing morphological characters with some Nearctic species of *Cephalcia* and with the genus *Acantholyda* COSTA (BENEŠ, 1976; VAN ACHTERBERG & VAN AARTSEN, 1986). *C. hartigii* appears to be rather uniform in both morphological and colour features although its distribution pattern is discontinuous (BENEŠ, 1976).

During field work on the species of *Cephalcia* living on spruce (*Picea abies* KARST.) in a region of Northern Italy, we collected some specimens of *C. hartigii* in mixed stands of spruce and fir (*Abies alba* MILL.) (CESCATTI & BATTISTI, 1992). These records suggested further research into the biology and morphology of this interesting species.

STUDY SITE

The study site (Mt. Paganella and Mt. Fausior) is situated between the towns of Trento and Bolzano, on the right side of the river Adige and in front of the Brenta Dolomites. This area falls between 46°09' and 46°11' North and between 13°50' and 13°56' East. The altitude of the conifer stands considered here, ranged from 950 m and 1840 m (measured from sea level) and the main exposure is N-W.

¹ Authors' names are in alphabetical order.

the ground. The sticky boards were used as follows: 40 boards in 10 sites in 1989, 15 boards in one site in 1990 .

In addition, we used a trunk trap (designed and constructed by A. ZANOCCO for *Cephalcia* spp.), consisting of a chamber of plastic or wire 1.5 mm mesh screen, ending with a tube, in order to obtain living insects. These traps were used where *C. hartigii* had been collected by the sticky traps, for a total amount of 4 traps in 1990 and 14 traps in 1991.




Fig. 2: Female and male of *Cephalcia hartigii* (Photo P. PAOLUCCI).

In the site where the highest number of adults was collected, in the summer of 1991, a total amount of 60 m² of plastic net was placed under the crown of 19 firs. The nets were checked every 4 days, searching for mature larvae falling from the crowns, at the end of their feeding. The only tree under which we trapped larvae of pamphiliid sawflies was cut in the autumn, its crown analysed and every track of larval activity was collected. The number and the age of the needles eaten by the larvae were evaluated for each feeding. The weight of the needles was estimated using needles of corresponding age collected from the nearest twigs.

RESULTS AND DISCUSSION

Catches and host plant

A total of 16 adults (14 males and 2 females) was collected in the 3 years of the study (Tab. 1) (Fig. 2). The first catches (6 males in 1989) were performed inadvertently during a research on the *Cephalcia* spp. of *P. abies*, only in stands mixed with *A. alba*. All the individuals were found on sticky traps where adults of *Cephal-*

cia arvensis PANZER or *C. abietis* (LINNAEUS) or *C. alashanica* (GUSSAKOVSKIJ) had been caught and were thought to be still living.

The absence of *Pinus* species, assumed to be the host plants of this sawfly, and the fact that we failed to find it in pure stands of spruce, led us to suppose that *A. alba* could be a host plant of *C. hartigii*. Therefore, in the following years we

Tab. 1: Catches of *Cephalcia hartigii* adults and larvae in the study site.

DATE	MM	FF	LL	LOCALITY	NOTE
16 6 1989	1	0		Trentino, Fai, M. Paganella (1150 m)	on tree band on <i>Picea abies</i>
23 6 1989	3	0		Trentino, Andalo, M. Paganella (1385 m)	on tree band on <i>Picea abies</i>
1 7 1989	2	0		Trentino, Andalo, M. Paganella (1385 m)	on yellow board
1 6 1990	1	0		Trentino, Fai, M. Paganella (1150 m)	on transparent board
19 6 1990	2	0		Trentino, Spormaggiore, M. Fausior (1250 m)	on tree band on <i>Abies alba</i>
18 6 1991	4	1		Trentino, Spormaggiore, M. Fausior (1250 m)	on tree band on <i>Abies alba</i>
22 6 1991	1	0		Trentino, Spormaggiore, M. Fausior (1250 m)	on tree band on <i>Abies alba</i>
1 8 1991	0	1		Trentino, Spormaggiore, M. Fausior (1250 m)	on trunk trap on <i>Abies alba</i>
14 9 1991			1	Trentino, Spormaggiore, M. Fausior (1250 m)	on net collector
4 10 1991			1	Trentino, Spormaggiore, M. Fausior (1250 m)	on net collector
TOTAL	14	2	2		

concentrated on stands with a greater presence of this species, in the area of Mt. Fausior. Here, in 1990 and 1991, we collected 7 males and 2 females, while another male was collected on Mt. Paganella. The females were caught by tree bands and by a trunk trap, both placed on *A. alba*. Unfortunately, the female caught by the trunk trap died before the check and, therefore, it was impossible to undertake any rearing experiments. The finding of the females induced us to search for the larvae and we collected two larvae, with some exuviae and frass, of pamphiliids in the nets placed under a fir.

The successive findings of two typical pamphiliid larval shelters and feeding remains in the crown of this fir gave us evidence that *A. alba* is a host plant of *C. hartigii*.

Notes on the life history

The flight period of *C. hartigii* occurred mainly in the second half of June. The exceptional female, collected at the beginning of August, could be explained by the longevity of the females of the genus *Cephalcia* (PSCHORN-WALCHER, 1982). In fact, this female contained very few eggs in her ovaries, probably having already laid her eggs.

The finding of the remains of an egg, near the larval shelter, suggests that the egg was laid isolated on the upper side of an old needle (shade needle), in the low-medium part of the crown (Fig. 3). The egg was partially inserted into the needle toward the apex. Two mm under the egg, around the needle, was a ring of a dark sticky substance, similar to that found in *C. arvensis* (BATTISTI & DA ROS, 1992). The larval feeding and the size of the shelter are presented in Tab. 2 and Fig. 4. The




Fig. 3: A hatched egg of *Cephalcia hartigii* on a needle of *Abies alba* (Photo P. PAOLUCCI).




Fig. 4: A larval shelter of *Cephalcia hartigii* on a twig of *Abies alba* (Photo P. PAOLUCCI).

differences observed between the two individuals could be ascribed to the sex, males having one instar less than females in this genus. The larva feeds along the twig in front of and behind the shelter, leaving only those surrounding it. The shelter is connected to the needles by several silk threads. The larval development seems to last two months or more, the mature larvae falling to the ground in September and October.


Tab. 2: Larval feeding and shelter size of two individuals of *Cephalcia hartigii*.

	LARVA 1	LARVA 2
Feeding		
number of needles 2-year-old	0	10
number of needles 3-year-old	45	75
number of needles 4-year-old	87	73
number of needles 5-year-old	56	0
total number of needles	188	158
dry matter of needles (g)	1.251	1.059
Shelter		
diameter (mm)	3.85	3.31
length (mm)	24.6	20.1
Twig bearing the shelter		
diameter (mm)	3.61	2.62
age (years)	5	4

Description of the mature larva

The first finding of the mature larva of *C. hartigii* requires its morphological description for both identification needs and for taxonomic studies. According to the authors who have studied the morphology of *Cephalcia* larvae (LORENZ & KRAUS, 1957; EIDT, 1969; MARTINEK, 1988), the specific distinction of the larvae is possible only on the basis of some sclerotisation patterns of both head and abdomen, which become particularly evident in the mature larva.

The mature larva of *C. hartigii* is typical of *Cephalcia*, showing all the general features described by LORENZ & KRAUS (1957) and by EIDT (1969). Freshly-collected specimens are brown-green in colour. The sclerotisation patterns of both head and IX abdominal segment (ventral side) suggest some differences in relation to the other species described by the authors (Fig. 5). However, it was not possible to consider the intraspecific variation of *C. hartigii* and more material is needed in order to clarify this aspect.

Fig. 5: IX abdominal segment (ventral side) and head of a mature larva of *Cephalcia hartigii*.

Geographical distribution of *C. hartigii*

The distribution of *C. hartigii* is presented in Tab. 3 and Fig. 6. All of the known localities except three (n. 8, 9 and 14), match well with the distribution pattern of *A. alba* (SCHMUCKER, 1942; MEUSEL *et al.*, 1965) (Fig. 6). In particular, the holotype was collected in a site (Mandacheregg, Swiss Jura) where the fir is present (personal communication of Prof. E. OTT, Zurich). Furthermore, the records of Lombardy (Italy) and that of Cantal Lioran (France) fall within large, natural stands of this conifer. The French stands are also used for seed crop (CTGREF, 1976). The non-specific topographical reference of the other records do not allow a precise connection with stands of *A. alba*.

Tab. 3: Findings of *Cephalcia hartigii* from the literature.

n	DATE	MM	FF	LOCALITY	STATE	LEGIT
1	24 5 1844	(1)	1	Mandacheregg, Aargau	Switzerland	J. Bremi
2	10 6 1897	1	0	C. Moravia Milkov	Czechoslovakia	J. Slaviček
3	30 5 1900	0	1	Frimb. (Frymburk, S. Bohemia ?)	Czechoslovakia	P. Kubes
4	1 6 1914	2	0	C. Moravia Milkov	Czechoslovakia	J. Slaviček
5	26 5 1931	0	1	N.E. Moravia Libhošť'	Czechoslovakia	F. Gregor
6	12 6 1931	0	1	Lombardy, Valtellina, Gerola (1050)	Italy	G. Binaghi
7	26 4 1982		?	Lower Tatra, M. Dumbier (1400 m)	Czechoslovakia	A. Taeger
8	12 6 1984		?	W Caucasus, Teberda (1600 m)	Russia	J. Oehlke
9	?		1	Valley River Thana (Caucasus)	EX URSS	E. König
10	?	1	0	Vorarlberg Feldkirch	Austria	?
11	?		?	Thüringen, Ohrdruf	Germany	Schmiedeknecht
12	?		?	Thüringen, Blankenburg	Germany	Mengersen
13	?		?	Cantal Lioran	France	?
14	?		1	Tunis	Tunisia	?

n	NOTE	AUTHOR	PUBL.
1	a pair on <i>Pinus sylvestris</i> , only female caught (Holotype)	BREMI	1849
2	Coll. Moravian Museum, Brno	BENEŠ	1976
3	Coll. National Museum, Prague	BENEŠ	1976
4	Coll. Moravian Museum, Brno	BENEŠ	1976
5	Coll. National Museum, Prague	BENEŠ	1976
6	Coll. Doderò	PESARINI	1976
7	only seen, not caught	TAEGER, in litt.	1992
8	Coll. Taeger	TAEGER, in litt.	1992
9	Coll. DEI	BENEŠ	1976
10	Coll. Konow, DEI (Allotype)	ENSLIN	1917
11		SCHMIEDEKNECHT	1907
12		SCHMIEDEKNECHT	1907
13		PUTON	1883
14	Coll. Graeffe, Nat. Hist. Mus. Wien	BENEŠ	1976

One of the three records (n. 8) outside the range of *A. alba* falls within the range of *Abies nordmanniana* (STEV.) SPACH (Caucasus); the presence of mixed

stands of *A. nordmanniana* and *Picea orientalis* (L.) LINK, in the region of Teberda, is reported by SCHMIDT-VOGT (1977). Probably the other record from the Caucasus (n. 9) falls within the range of *A. nordmanniana* too, being the river Thana (= Tana) a tributary of the river Kura, not far from Tbilisi (BENEŠ, in litt. 25/2/92), but we cannot find any geographical reference to this valley.




Fig. 6: Distribution of *Abies alba* according to SCHMUCKER (1942) and MEUSEL *et al.* (1965) and findings of *Cephalcia hartigii*. Numbers refer to Tab. 3 and the n. 15 shows the site of the new records.

The third record (n. 14) is attributed to “Tunis” without other information. The nearest stands of fir (*Abies numidica* DE LANNOY) are localised on the Tell Atlas (Mt. Babor, 1500/1600-2000 m a.s.l., about 1000 ha) in the coastal part of Algeria (MAYER, 1965). The lack of more precise information about this record suggests that we consider its geographic origin doubtful. *C. hartigii* is reported also for Yugoslavia by BENEŠ (1976) and VIITASAARI (1982), but we could not find any more detailed reference about this record.

To our knowledge, *C. hartigii* seems to be the only species of *Cephalcia* living on *Abies* sp. in Europe; it is interesting to note that in North America there is only one species (*C. distincta* (MACGILLIVRAY)) living on *Abies balsamea* (L.) MILL. and, second in order, on *Tsuga canadensis* (L.) CARR. (EIDT, 1969).

Considering the fact that some records are not clearly referable to fir stands, we cannot exclude the possibility that *C. hartigii* may also live on other host plants. Further research should be done to clarify this aspect of the biology of *C. hartigii*.

ACKNOWLEDGEMENTS

The authors thank: Prof. L. MASUTTI for the useful suggestions and for the determination of the insect material; Dr. K. BENEŠ and Dr. A. TAEGER for the original information given about some records of *C. hartigii*. The work benefits from comments by: Dr. K. BENEŠ, Prof. O. EICHHORN, Prof. L. MASUTTI, Prof. H. PSCHORN-WALCHER, Dr. N. SPRINGATE, Dr. A. TAEGER. The authors are also grateful to Dr. N. SPRINGATE for the revision of the English and to D. PÖRNBACHER for the German summary. The work was supported by a C.N.R. grant N. 90.02685.

ZUSAMMENFASSUNG

Zur Kenntnis der Wirtspflanze, der Larvenmerkmale und der Biologie von *Cephalcia hartigii* (BREMI) (Hym., Pamphiliidae).- *Cephalcia hartigii* ist eine seltene Species, von der in Mittel- und Südeuropa, Nordafrika und vom Kaukasus berichtet wird. Ihre Biologie und Wirtspflanzen waren unbekannt. Während einer Felduntersuchung der Gespinstblattwespen in Norditalien wurden 16 Imagines (14 männlich und 2 weiblich) dieser Species mit verschiedenen Fallentypen gesammelt. Die meisten von diesen wurden an *Abies alba* gefunden. Das Ergreifen von 2 erwachsenen Larven, die auf den Boden unter einer Tanne gefallen waren, und die spätere Auffindung von 2 typischen Larvenspinnstücken hat uns gezeigt, daß *Abies alba* eine Wirtspflanze von *C. hartigii* ist. Es werden Ergebnisse über die Flugperiode, die Eiablage, den Larvenfraß und die Larvenentwicklung dargestellt. Die morphologische Hauptcharakteristik der erwachsenen Larve wird kurz beschrieben. Viele der bibliographischen Hinweise von *C. hartigii* stimmen gut mit der Verbreitung von *A. alba* und *A. nordmanniana* überein. Da mehrere der Hinweise nicht genau Tannenbeständen zuzuschreiben sind, ist die Möglichkeit nicht auszuschließen, daß *C. hartigii* auch an anderen Wirtspflanzen lebt.

REFERENCES

- ACHTERBERG, C. VAN & AARTENS, B. VAN 1986. The European Pamphiliidae (Hymenoptera: Symphyta) with special reference to the Netherlands. *Zool. Verh.* 234: 3-98.
- BATTISTI, A. & DA ROS, N. 1992. Accounts on the bio-ecology of *Trichogramma cephalciae* HOCHMUT & MARTINEK in N-E Italy (Hym., Trichogrammatidae). *Insect Parasitoids*. 4th Eur. Workshop *Redia* 74(3): 179-184.
- BENEŠ, K. 1976. Revision of the European species of the genus *Cephalcia* PANZER, 1805 (Hym. Pamphiliidae). *Studie ČSAV* 3: 1-67.
- BREMI J. 1849. Beschreibung einiger Hymenopteren, die ich für noch unbeschriebene und unpublizierte halte. *Stettin. ent. Ztg.* 10: 92-96.
- CESCATTI, A. & BATTISTI, A. 1992. Distribution and ecology of *Lymantria monacha* L. and *Cephalcia* spp. in non-outbreak areas of Trentino (N-Italy). *Anz. Schädlingskde., Pflanzenschutz, Umweltschutz* 65: 92-99.
- CTGREF 1976. Semences forestières. Les régions de provenance de sapin pectiné (*Abies alba* MILL.). *Note technique, Centre technique du génie rural, des eaux et des forêts (Min. Agric. Franç.)* 31:22.
- EIDT, D.C. 1969. The life histories, distribution, and immature forms of the North American sawflies of the genus *Cephalcia* (Hymenoptera: Pamphiliidae). *Mem. Ent. Soc. Can.* 59: 3-56.
- ENSLIN, E. 1918. Die Tenthredinoidea Mitteleuropas. VII. *Dt. ent. Z.* 1917: 663-673.
- LORENZ, H. & KRAUS, M. 1957. *Die Larvalsystematik der Blattwespen*. AkademieVerlag, Berlin, 339 pp.
- MARTINEK, V. 1988. Rozlišování eonymf a pronymf ploskohřbetek. *Lesnická práce* 67: 415-418.
- MAYER, H. 1965. Marokkanische und numidische Tanne im nordafrikanischen Atlas. *Cbl. ges. Forstwesen.* 84: 309-324.
- MAYER, H. & HOFMANN, A. 1969. *Tannenreiche Wälder am Südfall der mittleren Ostalpen*. BLV Verlagsgesellschaft, 259 pp.
- MEUSEL, H., JÄGER, E. & WEINHERT, E. 1965. *Vergleichende Chorologie der Zentraleuropäischen Flora*. Fischer Verlag, Jena, Karten 258 pp., Text 583 pp.
- PESARINI, C. & PESARINI, F. 1976. Materiali per un catalogo degli Imenotteri Sinfiti Italiani. I. Famiglia Pamphiliidae (Hymenoptera). *Boll. Soc. Ent. It.* 108: 53-66.
- PSCHORN-WALCHER, H. 1982. Symphyta. In SCHWENKE, W. (ed) *Die Forstschädlinge Europas*. Bd. IV. pp. 4-234. Paul Parey, Hamburg und Berlin.
- PUTON, A. 1883. Supplément à la liste des Tenthredides de France. *Revue Ent.* 2: 15-16.
- SCHMIDT-VOGT, H. 1977. *Die Fichte*. Band I. Parey, Hamburg und Berlin, XVIII+647 pp.
- SCHMIEDEKNECHT, O. 1907. *Hymenopteren Mitteleuropas*. Fischer Verlag, Jena, VII+804 pp.

SCHMUCKER, T. 1942. *Silvae orbis*. C.I.S. Berlin, pp. X+156 (Karten 250).

VIITASAARI, N. 1982. Sahapistiäiset 2. Xyeloidea ja Megalodontoidea. *Report Univ. Helsinki/Dep. Agric. For. Zool.* 5: 1-72.

WALTER, H. & LIETH, H. 1960. *Klimadiagramm - Weltatlas*. Fischer Verlag, Jena.

(received May 4, 1992; accepted September 2, 1992)