**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft =

Bulletin de la Société Entomologique Suisse = Journal of the Swiss

**Entomological Society** 

Herausgeber: Schweizerische Entomologische Gesellschaft

**Band:** 67 (1994)

**Heft:** 1-2

**Artikel:** Cytotoxic assay of 2,4-dichlorophenoxyacetic acid (2,4-D) by mitotic

index profiles in Culex pipiens fatigans

Autor: Ahmad, Waseem / Ali, M. Niamat

**DOI:** https://doi.org/10.5169/seals-402552

# Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 02.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Cytotoxic assay of 2,4–dichlorophenoxyacetic acid (2,4–D) by mitotic index profiles in *Culex pipiens fatigans*

# Waseem Ahmad & M. Niamat Ali

Section of Genetics, Chemical Mutagenesis Laboratory; Department of Zoology, Aligarh Muslim University, Aligarh, U.P., 202 002, India

Neuroblast cells obtained from the brain tissues of larval stages of *Culex pipiens fatigans* were subjected to mitotic index analysis as part of an elaborate study of 2,4-dichlorophenoxyacetic acid (2,4–D) induced genotoxicity. Multiple applications in varying concentrations and durations were maintained for three consecutive generations. Mitotic activity was seriously hampered in all the experiments. A steady state of depression in dividing cells was observed in all generations, more conspicuously in F<sub>1</sub>. The cell growth was inhibited and larval developmental time in treated series was affected and significantly delayed. 2,4-D was found to be extremely cyto-toxic in this organism.

Keywords: Cytotoxic assay, 2,4-dichlorophenoxyacetic acid (2,4-D), Culex pipiens fatigans.

#### INTRODUCTION

Since the late 1940's, 2,4-dichlorophenoxyacetic acid (2,4–D) has been widely used for its dual properties in plants. At higher concentrations it is used as a plant killer (Allard *et al.*, 1946; Hamner *et al.*, 1946; Moore, 1974) and at lower concentrations it is known to promote growth in a manner apparently identical with that of indole-acetic acid (Hsueh & Lou, 1947; Moore, 1974). The importance of 2,4–D for agriculture and forestry resulted in increased numbers of reports on its mode of action (Bamberger, 1970; Ashton & Crafts, 1973; Loos, 1975; Wegler & Eue, 1977).

Recently 2,4–D has been so widely used that similar studies have been extended to animals as well.

Jenssen & Remberg (1976) reported the broad cytogenetic effects in mouse blood. In-vivo tests performed by Yourchenko (1977) confirmed these findings. Cytogenetic effects have also been studied in SV–40 transformed human fibroblasts (Ahmed *et al.* 1977). However, all the reports are not positive. 2,4–D was not significantly able to influence DNA synthesis in mouse testes (Seiler, 1978). A less clear relationship is obtained with 2,4–D in animals.

Derivatives of phenoxyacetic acid, notably pentachlorophenol (PCP), were discovered subsequently to have different properties. These include seriously hampering the cell growth and inhibition of DNA, RNA and ribosome synthesis (Sikka & Sharma, 1976; Chand, 1980; Ehrlich *et al.*, 1987; De Marini *et al.*, 1990).

By that reckoning the toxicological evaluation of 2,4–D was not so high. A perusal of available literature indicated that barring reports of Mohandas & Grant (1972) and Bayliss (1973) – who studied 2,4–D induced mitotic abnormalities in *Drosophila* – not many studies have been undertaken relating to mitotic index, which is a convenient method of assessing the mutagenicity in any organism, especially in insects. Occasional studies by Vogel & Chandler (1974), Magnusson *et al.* (1977) and Rasmusson (1977) were not conclusive.

An elaborate mutagenicity testing programme in filarial vector *Culex pipiens* fatigans was therefore undertaken. The present report is a part of it and deals specifically with the effects of 2,4–D on the mitotic activity in neuroblast cells and its overall affect on growth and development. The results are unique and assume a significant importance in the absence of any real answer to the problems arising out of indiscriminate use of 2,4–D from a genetic point of view.

## MATERIALS AND METHODS

Recently, indiscriminate use of 2,4–D is being debated. For these experiments, a border line dose, 100 parts per million (ppm) in between the herbicidal and hormonal properties, was selected. Lesser doses of 75, 50, 25 and 1 ppm were also tried. For various concentrations a stock solution of 1000 ppm in 2% alcoholic distilled water made on a volumetric basis was used and administered through whole body immersion of 4<sup>th</sup> instar larvae in 2 and 4 hrs exposures series. The experiment was initiated by taking 1000 larvae of uniform size and age in each concentration and duration. Consequently 100 larvae were sacrificed for brain squashes and the rest were inbred to obtain the next generation. Three subsequent exposures of 2,4–D were given in as many generations.

Serial exposure as adopted above is in contrast to single application so often used by the majority of authors (Green et al., 1973; Khalatkar & Bhargava, 1982; Kumari & Vaidyanath, 1989). This ensured extensive information about the nature and extent of damage to neuroblast cells, and facilitated reliable comparative studies in and among the progenies. Strict controls were maintained simultaneously in 2% alcoholic water. The experiments were replicated at least 3 times under identical conditions. Other details relating to feeding and rearing were similar to those described earlier (Ahmad et al., 1983).

The cytological preparation were made using acetic acid-dissociated-air drying techniques (Crozier, 1968). Dividing cells from over a total of 20,000 were thoroughly observed. Mitotic index (MI) was computed by the formula

$$MI = \frac{\text{Total no. of dividing cells}}{\text{Total no. of cells observed}} \times 100$$

In another study undertaken to observe the effects of 2,4–D on acquatic development, all the details pertaining to preparation, concentration and treatment of 2,4–D were uniform, though the stage of treatment was different – 1<sup>st</sup> instar in this case. A total of 10 batches each of 100 larvae were exposed and a separate control series was also maintained for comparison. Total larval duration and time to pupation, its range and mortality were carefully recorded at regular intervals for the entire aquatic duration.

# RESULTS AND DISCUSSION

Profile of mitotic indices, inhibitory effect of 2,4–D on dividing cells and delayed larval development have been shown in Tab. 1 and Figs. 1 and 2, respectively.

As recorded earlier most of the given concentrations of 2,4–D register serious consequences to the dividing cells from one generation to another, indicating a cumulative effect, consequently as a result of constant pressure of multiple exposure throughout. A steady rise in the frequency of injured cells with respect to chro-

mosomes is observed: There is corresponding increase in breaks, acentrics, dicentrics, gaps, ploidy and other aberrations (ALI & AHMAD, 1994a, 1994b). Changes studied at morphological level are equally significant; some of them already discovered, register increased frequencies while others were new (ALI & AHMAD, 1994c). These studies substantiated the cumulative effects. Incidentally the initial chemical damage induced by 2,4–D is subsequently manifested at the level of chromosome, some of which can be related to morphological mutations.

Conversely, changes in mitotic profiles are more due to higher concentrations rather than cumulative pressure. Incidentally serious impairment of divisional stages and low mitotic rates leave a reasonably high number of cells with impaired and injured chromosomes. Comparable effects achieved with higher concentration were mimicked by lower concentrations in a later generation (Fig. 1). Depression in the




Fig. 1. Effect of various concentrations of 2,4–dichlorophenoxyacetic acid on mitotic activity in *Culex pipiens fatigans*. Exposure for each concentration was given for three successive generations. The decreasing profiles of the Mitotic Index are evident. – Symbols:  $\bullet = F_1$ ;  $O = F_2$ ;  $O = F_3$ ; full lines: 2hrs, hatched lines: 4 hrs exposure; C = control.

rate of division in exposed forms significantly increased the larval duration and further induced distorted growth in few cases. Time to pupation as a result of slow growth was increased by as much as 50 hrs in some replicates over and above the normal 183 hours. This has been depicted in Fig. 2.

A stable state of sterility induced by higher concentrations beyond 50 ppm is another striking effect of 2,4–D.

Inconclusive results indicate this phenomenon to be due to an impairment of spermatogenesis. Meiotic tissues are therefore under similar investigations. In the light of the above indications and also from the reports on pentachlorophenol (Sikka & Sharma, 1976), the potential of male sterility in genetic control of insect vectors of disease and pest management is fairly promising.




Fig. 2. Effect of various concentrations of 2,4-dichlorophenoxyacetic acid on larval development in *Culex pipiens fatigans*. Exposure for each concentration was given for three successive generations. There is a pronounced delayed development. – Symbols:  $\bullet = F_1$ ;  $O = F_2$ ;  $O = F_3$ ; full lines: 2hrs, hatched lines: 4 hrs exposure; C = control.

Tab. 1. Mitotic index (MI) of 2,4-D treated neuroblast cells.

Generation Concentration Duration	Cells screened	Cells in dividing stage	$MI \pm S.D.$
P1: Control	1130	985	$87.15 \pm 2.83$
F1: $C_1^2$	943	766	$81.21 \pm 3.46$
$C_1^4$	1234	982	$79.56 \pm 3.85$
$C_{25}^{2}$	756	610	$80.63 \pm 2.93$
C <sub>25</sub> <sup>4</sup>	1039	94	$76.35 \pm 4.62$
$C_{50}^{2}$	847	622	$73.43 \pm 4.08$
C <sub>50</sub> <sup>4</sup>	524	373	$71.01 \pm 2.76$
$C_{75}^{2}$	752	381	$50.59 \pm 2.30$
C <sub>75</sub> <sup>4</sup>	1665	836	$50.17 \pm 3.92$
$C_{100}^{2}$	425	199	$46.53 \pm 2.25$
$C_{100}^{-4}$	621	258	$41.49 \pm 4.00$
F2: C <sub>1</sub> <sup>2</sup>	1132	889	$78.51 \pm 3.64$
$C_1^4$	1367	1023	$74.80 \pm 2.04$
$C_{25}^{2}$	692	493	$71.94 \pm 2.25$
$C_{25}^{-4}$	963	657	$70.27 \pm 3.85$
$C_{50}^{2}$	715	426	$64.19 \pm 3.25$
$C_{50}^{4}/C_{75}/C_{100}$	_*	_*	_*
F3: C <sub>1</sub> <sup>2</sup>	620	456	$73.38 \pm 3.25$
$C_1^4$	992	724	$72.80 \pm 3.85$
$C_{25}^{2}$	559	387	$69.47 \pm 4.49$
$C_{25}^{4}$	963	657	$68.17 \pm 4.85$
$C_{50}^{2}$	715	426	$58.83 \pm 4.75$
$C_{50}^{4} / C_{75} / C_{100}$	_*	_*	_*

C-subscripts refer to concentrations in 1, 25, 50, 75 & 100 ppm; superscripts refer to 2 or 4 hours treatment; \* = sterility induced in these concentrations.

The low solubility of 2,4–D, as with most acids, makes it apparently less toxic on a molar basis. In the majority of mammals an accumulation of phenoxy acid is unlikely because of quick clearance (Seiler, 1978). This could not be confirmed in mosquitoes by any such report. It might, however, be possible that repeated exposures may have created a steady state of concentration capable of causing serious inhibition to mitotic cycle, inflicting injuries to chromosomes and delaying the development of various acquatic stages.

There are similar reports on 2,4–D in plant tissues – a comparison, however, had a limited validity (Melchers & Bergmann, 1959; Bayliss, 1973). In general it has been observed that phenoxy acids do influence the genetic material or the

mitotic cycle or both, but owing to the physiological activity of these substances on the plant cells, its significance for the present observations cannot be clearly assessed.

#### **ACKNOWLEDGEMENTS**

This study is supported by research grant No. F 3-206/90 (SR-II) by University Grants Commission, New Delhi, India. Chairman, Department of Zoology, is gratefully acknowledged for providing facilities.

#### **REFERENCES**

- Ahmad, W., A. Ara & U.M. Adhami, 1983. Genetics of "distorted" (d) in *Culex pipiens fatigans. Mitt. Schweiz. Entomol. Ges.* 56: 271-273.
- Ahmed, F.E., N.J. Lewis & R.W. Hart, 1977. Pesticide induced ouabain resistant mutants in Chinese hamster V79 cells. *Chem.-Biol. Interaction* 19: 369-374.
- ALI, M.N. & W. AHMAD, 1994a. Genotoxicity of 2,4-dichlorophenoxyacetic acid (2,4-D) on filarial vector *Culex pipiens fatigans*. *J. envir. Biol.* (In press).
- ALI, M.N. & W. Ahmad, 1994b. Mutagenic activity of 2,4-dichlorophenoxyacetic acid (2,4-D) in the neuroblast micronucleus assay in *Culex pipiens fatigans*. *Pollution Res.* 13 No.2 (In press).
- ALI, M.N. & W. AHMAD, 1994c. Mutagenic assessment of 2,4–D in *Culex pipiens fatigans*: Inheritance, Frequency distribution and test statistics (ø) of *crs* mutants. *Pollution Res*. (Communicated).
- ALLARD, R.W., H.R. DEROSE & C.P. SWANSON, 1946. Some effects of plant growth regulators on seed germination and seedling development. *Botan. Gaz.* 107: 575-583.
- Ashton, F.M. & A.S. Crafts, 1973. *Mode of Action of Herbicides*. John Wiley and Sons, New York, pp. 266-288.
- BAMBERGER, E.S., 1970. The effect of plant growth regulators on DNA. *Phytochemistry 10*: 957-966. BAYLISS, W.M., 1973. Origin of chromosome number variation in cultures plant cells. *Nature 246*: 529-530.
- CHAND, S., 1980. Effects of herbicide pentachlorophenol on mitosis and DNA synthesis in *Nigella sativa* L. *Acta Bot. Ind.* 8: 219-223.
- Crozier, R.H., 1968. An acetic acid dissociation, air-drying technique for insect chromosomes, with aceto-lactic orcein staining, *Stain Tech.* 43: 171-173.
- DE MARINI, D.M., H.G. BROOKS & D.G. PARKES, Jr., 1990. Induction of prophage lambda by chlorophenols. *Environ. Mol. Mutagen.* 15: 1-9.
- EHRLICH, W., M. MANGIR & E.R. LOCHMANN, 1987. The effect of pentachlorophenol and its metabolite tetrachloro-hydroquinone on RNA, protein, and ribosome synthesis in *Saccharomyces* cells. *Ecotoxicol. Environ. Safety 13:* 7-12.
- Green, S., F.M. Sauro & M.S. Legator, 1973. Cytogenetic effects of hycanthone in the rat. *Mutation Res.*, 17: 239-244.
- HAMNER, C.L., J.E. MOULTON & H.B. TUKEY, 1946. Effect of treating soil and seeds with 2,4–D on germination and development of seedlings. *Botan. Gaz. 107:* 352-361.
- HSUEH, Y.L. & C.H. Lou, 1947. Effect of 2,4-D on seed germination and respiration, *Science 105:* 283-285.
- JENSSEN, D. & L. REMBERG, 1976. Distribution and cytogenetic test of 2,4–D and 2,4,5–T phenoxyacetic acids in mouse blood tissues. *Chem.-Biol. Interactions* 14: 291-299.
- KHALATKAR, A.S. & Y.R. BHARGAVA, 1982. 2,4-Dichlorophenoxyacetic acid. A new environmental mutagen. *Mutation Res.* 103: 111-114.
- Kumari, T.S. & K. Vaidyanath, 1989. Testing of genotoxic effects of 2,4-dichlorophenoxyacetic acid (2,4-D) using multiple genetic assay systems of plants. *Mutation Res.* 226: 236-238.
- Loos, M.A., 1975. Phenoxyalkanoic acids. In: P.C. Kearney & D.D. Kaufman (Eds). Herbicides. Chemistry, Degradation and Mode of Action. 2nd ed., Marcel Dekker, New York, pp. 1-128.
- Magnusson, J., C. Ramel & A. Erikson, 1977. Mutagenic effects of chlorinated phenoxy acids in *Drosophila melanogaster. Hereditas* 87: 121-123.
- Melchers, M. & L. Bergmann, 1959. Ber. dt. bot. Ges. 71: 459.
- MOHANDAS, T. & W.F. Grant, 1972. Cytogenetic effects of 2.4–D and amitrol in relation to nuclear volume and DNA content in some higher plants. *Can. J. Genet. Cytol.* 14: 773-783.
- Moore, T.C., 1974. Effects of certain synthetic plant growth regulators on the development of selected species. In: Research Experiences in Plant Physiology A Laboratory Manual. Springer, New York, pp. 307-323.

- RASMUSSON, B., 1977. Mutagenicity tests of 2,4–D and 2,4,5–T in genetically stable and unstable strains of D. melanogaster. Conference on "Chlorinated phenoxyacids and heir dioxins, Mode of action, health risks and environmental effects". The Royal Swedish Academic Sciences, Stockholm, Feb. 6-10.
- SEILER, J.P., 1978. The genetic toxicology of phenoxy acids other than 2,4,5–T. *Mutation Res.* 55: 197-220.
- Sikka, K. & A.K. Sharma, 1976. The effect of some herbicides on plant chromosomes. *Proc. Ind. Natl. Sci. Acad. Part B* 42: 299- 307.
- Vogel, E. & J.L.R. Chandler, 1974: Mutagenicity testing of cyclamate and some pesticides in *Drosophila melanogaster*. Experientia 30: 621-623.
- Wegler, R. & L. Eue, 1977. Herbizide. In: R. Wegler (Ed.), *Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel*, Vol. 5. Springer, Berlin, Heidelberg, New York.
- YOURCHENKO, V.V., 1977. (Cytogenetic investigation of mutagenic properties, common to the repellents dimethylphthalate and N,N-diethylamide of phenoxyacetic acid.) *Pharmacol. Toksikol.* 1977: 454-457 (Russian).

(received August 8, 1993; accepted after revision March 4, 1994)