

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 69 (1996)

Heft: 3-4

Artikel: Description of the larva of Eucallia boussingaultii (Guérin, 1843)
(Coleoptera : Cicindelidae : Cicindelini)

Autor: Arndt, Erik / Cassola, Fabio / Putchkov, Alexander V.

DOI: <https://doi.org/10.5169/seals-402639>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Description of the larva of *Eucallia boussingaultii* (GUÉRIN, 1843)
(Coleoptera: Cicindelidae: Cicindelini)

ERIK ARNDT¹, FABIO CASSOLA² & ALEXANDER V. PUTCHKOV³

¹ Universität Leipzig, Institut für Zoologie, Talstr. 33, D-04103 Leipzig, Germany

² Via F. Tomassucci 12/20, I-00144 Roma, Italy

³ Academy of Sciences, Institute of Zoology, B. Hmelnitskogo 15, 252601 Kiev-30, Ukraine

The last instar larva of *Eucallia boussingaultii* (GUÉRIN, 1843) is described and figured in detail. Larval morphological characters confirm the transfer of the genus *Eucallia* from the tribe Megacephalini to the Cicindelini (subtribe Iresina). The T-shaped gular suture, the pronotum with the distinct semi-circular impressions on the disk, the spines of the first labial palpomere, and the 3 sclerotized hooks on the abdominal tergite V are typical characters of Iresina including *Eucallia*.

Keywords: Cicindelidae, *Eucallia*, larva, taxonomy.

INTRODUCTION

The cicindelid genus *Eucallia* GUÉRIN comprising only one species, *E. boussingaultii*, is distributed from the province of Santander del Norte in northeastern Colombia across Ecuador to the province of Cajamarca in northern Peru. The species inhabits humid mountain forests; adults stay along forest roads on moss covered stones (RODRIGUEZ *et al.*, 1994). Larvae are found on red clay banks with lichens.

Originally, *Eucallia* was placed in the tribe Megacephalini near the genus *Oxycheila* DEJEAN (HORN, 1926). RIVALIER (1971) transferred the genus to the Cicindelini (subtribe Iresina) because of adult characters. Since both tribes are also clearly characterized by larval characters, larval morphology could confirm the present taxonomic position. Moreover, recent studies of the larval morphology strongly suggest that the genus *Oxycheila* should also be transferred to the Cicindelini (Iresina) (PUTCHKOV & ARNDT, in press).

Larval characters of *Eucallia* were already mentioned by GUÉRIN-MÉNEVILLE (1843, 1844) and CHAPUIS & CANDEZE (1853). However, these papers contain only few and superficial details. The larva of *Eucallia* is compared with those of *Cicindela campestris* and *C. hybrida*. Characters which clarify the taxonomical position are not mentioned.

Here, we want to describe the *Eucallia* larva in detail and to discuss the taxonomic position of the genus.

MATERIALS & METHODS

The description is based on the following larval material (all specimens last instar): 4 specimens Ecuador, Loja Prov., Cordillera Cordóncillo, 11 km South of Saraguro, 3130 m; collected by C. YOUNG, R. DAVIDSON, and J. RAWLINS on red clay banks with lichens on October 27, 1987. 22 specimens from the same region, Mamanuca, 2650 m, collected by G. ONORE on December 29, 1994. The larvae were col-

lected together with adults. The habitat near Saraguro was unusual because there were hundreds of horizontal burrows in vertical or slightly inclined red clay cliffs that varied from 3 to 10 metres in height, mostly covered with lichens (R. D. DAVIDSON, pers. commun.). Only 6 adults of *E. boussingaultii* but no other cicindelid species were found in the area, so doubtless the larvae belong to this species.

Larvae of 25 further Cicindelidae genera (including all tribes) were available for comparison (PUTCHKOV & ARNDT, 1994). The larvae of *E. boussingaultii* are deposited at the Carnegie Museum of Natural History, Pittsburgh, in the collection of Pontificia Universidad Católica of Quito and (single specimens) in the collections of the authors.

The nomenclature follows RIVALIER (1971); terms of morphology and chaetotaxy follow KNISLEY & PEARSON (1984).

DESCRIPTION OF THE LAST INSTAR LARVA

Measurements: Head width 2.92–3.33 (av. 3.20) mm; frontal width 1.74–2.04 (av. 1.92) mm; pronotal width 3.25–3.74 (av. 3.52) mm; pronotal length 2.03–2.33 (av. 2.19) mm, $n = 15$.

Coloration: Head blackish brown, pronotum dark brown with discal area darker than sides, rest of sclerites paler brown.




Fig. 1. *Eucallia boussingaultii*: Prementum and labium, ventral aspect; vs – ventral double sclerite.

Head: Six stemmata of characteristic size and position (see KNISLEY & PEARSON, 1984). Small tubercles, each with 2 long setae, are near the large stemmata I and II. The rest of the setae without or with indistinct tubercles; most setae of the head capsule thin, not flattened; some lateral setae near the eyes slightly flattened. Coronal suture very short, indistinct. Ridge on the posterodorsal part of the parietale weakly separated from the slightly U-shaped ridge of the posterior part of the frontale. Nasale produced, wide, anterior margin nearly straight, terminating in 2 large teeth at the sides. Antenna 4-segmented; antennomeres I–III of nearly the same length, last antennomere 1.4 times shorter and thinner than the other segments; antennomere I with 4–5 setae, 3 of them subapical on outer margin, antennomere II with 4 setae, antennomere III with 2 large setae; sensorial appendage of antennomere III lacking, replaced by a group of very small setae. Maxilla with 2-segmented cardo, dorsal part of cardo slender and ring-shaped, ventral part triangular with 1 seta, stipes slender with 2–3 spikes and a small group of short setae mediobasally; setal group gMX on inner side of stipes situated in a less sclerotized field with 24–28 setae in two rows, setae of the inner row long, those of the lateral row short. Lacinia lacking, between stipes and maxillary palpus a sclerotized segment with 3 setae present; palpus and galea of about equal length or palpus slightly shorter, first segments of palpomere and galea fused, galeomere I with 3 typical setae on inner margin, galeomere II with 4 strong bristles. Hypopharynx of typical structure, large, overlapping the prementum, dorsally densely covered with small hairs, laterally and ventrally with numerous long setae. Labium (Fig. 1) with a longish oval double sclerite on the ventral side between prementum and labial palpomere I; labial palpomere I with 4–5 strong setae and 3–4 spines apically, palpomere II with 1 strong bristle ventrally near middle.

Thorax: Pronotum (Fig. 2) with distinct relief-like structures, anteriorly with distinct semicircular impressions; anterolateral angles sharp, triangular; 16–24 setae





Fig. 2. Pronotum.



3

Fig. 3. Sclerites of abdominal segment III, right side; cl – coxal lobe (= laterosternite), ep – epipleuron, hy – hypopleuron, st – sternites, te – tergite.

on each pronotal half, 6–10 setae along median suture. Ridges of anterolateral angles weakly defined. Legs 5-segmented of typical cicindeloid shape, each with two separated claws, the anterior one longer.

Abdomen: Abdominal tergites I–IV with about 7 long and 6 very short and thin setae. Hypopleuron consisting of 1 large and 1 small sclerite (Fig. 3). Abdominal tergite V with anterior part separated, caudolateral and caudal parts fused; anterior part with 8–10 long setae, lateral part with 4–6 setae, the first of which inserts on the outer margin of the lateral hook, caudal part of tergite with 12–15 strong setae; 3 pairs of hooks present, median and inner hook comparatively straight and slender, each with 1–2 setae medially, median hook slightly curved outside, inner hook with 2 setae in its middle part (Fig. 4). Tergite IX with 7 pairs of long setae, pygopod conical, with 5–7 pairs of setae dorsally and dorsolaterally, apical margin of pygopod with 16–20 setae, 12 of them visible from dorsal side, ventral part without setae. Sternite IX with 6, rarely 7 long setae on posterior margin.

DISCUSSION

Larval characters clearly show that the genus *Eucallia* is a representative of the Cicindelini. Typical character states of Cicindelini larvae are the T-shaped anterior end of the gular suture, the pronotum with the distinct semicircular impressions on disk, the presence of spines on the first labial palpomere, and the shape of the hooks of the abdominal segment V (PUTCHKOV & ARNDT, 1994).

In the tribe Cicindelini, the genus *Eucallia* belongs to the Iresina group as proposed by RIVALIER (1971), because of the 3 pairs of hooks on the abdominal tergite V. The larva is similar to that of genus *Euprosopus* DEJEAN, which is distinguished by the large number of flattened setae on the pronotum, fewer setae on the pygopod and abdominal tergites, and a longer coronal suture. The larva of *Eucallia* is distinguished from that of *Oxycheila* DEJEAN (see ARNDT & PUTCHKOV, in press) by the lack of a dorsoapical spine on the first freestanding maxillary palpomere (palpomere I according to VAN EMDEN (1935) who considered the first segment as pal-




Fig. 4. Tergites and hooks of abdominal segment V, left side; ih – inner hook, mh – median hook, oh – outer hook.

pifer) and large numbers of stout setae on the pygopod. Furthermore, the third instar larva of *Oxycheila* has 3–6 bristles on the median hook, contrary to *Eucallia* with only two bristles on each hook. However, the 2nd instar larva of *Oxycheila* is very similar to *Eucallia* besides the very small spine on the maxillary palpomere. Larvae of *Eucallia* and *Oxycheila* share the form of the ridges on the posterior part of the head, and the very short coronal suture. In our opinion, these character states confirm the close relationship of these genera, and both should be included in the subtribe Iresina.

ACKNOWLEDGEMENTS

The authors are indebted to R. D. DAVIDSON (Carnegie Museum of Natural History, Pittsburgh, Pennsylvania) and G. ONORE (Pontificia Universidad Católica, Quito, Ecuador) for loan and gift of the valuable larval material. We also thank M. L. LUFT (Newcastle, UK) and D. L. PEARSON (Tempe, Arizona) for proof-reading the manuscript.

ZUSAMMENFASSUNG

Die Larve von *Eucallia boussingaultii* (GUÉRIN, 1843) wird beschrieben und abgebildet. Larvalmorphologische Merkmale bestätigen die Umstellung der Gattung *Eucallia* von den Megacephalini in die Cicindelini (Subtribus Iresina). Die T-förmige Gularsutur, halbkreisförmige Eindrücke am Pronotum, Stacheln am ersten Labialpalpenglied und drei Paare sklerotisierter Haken am Abdominaltergit V sind typische Merkmale der Iresina (einschließlich *Eucallia*).

REFERENCES

- CHAPUIS, M.F. & CANDEZE, M.E. 1853. Catalogue des Larves des Coléoptères, connues jusqu'à ce jour avec la description de plusieurs espèces nouvelles. *Mem. Soc. R. Scienc. Liège* 8: 347–654.
- VAN EMDEN, F.I. 1935. Die Larven der Cicindelinae. 1. Einleitendes und Alocosternale Phyle. *Tijdschr. Ent.* 78: 134–183.
- GUÉRIN-MÉNEVILLE, M.F.E. 1843. Insectes nouveaux observés sur les plateaux des Cordillères et dans les vallées chaudes de la Nouvelle-Grenade, décrits par M. GUÉRIN-MÉNEVILLE; avec des notes relatives à leur mœurs, à leur distribution géographique etc., par GOUDOT, M.J.: Travaux inédits. *Rev. Zool.* 6: 1–22.
- GUÉRIN-MÉNEVILLE, M.F.E. 1844. G. *Eucallia* GUÉRIN. *Mag. Zool.* 2, pl. 144: 1–2.
- HORN, W. 1926. Carabidae: Cicindelinae. In: JUNK, W., *Coleopterorum Catalogus* vol. 86, pp. 1–345.
- KNISLEY, C.B. & PEARSON, D.L. 1984. Biosystematics of larval tiger beetles of the Sulphur Springs Valley, Arizona. Descriptions of species and a review of larval characters for *Cicindela* (Coleoptera: Cicindelidae). *Trans. Amer. Entom. Soc.* 110: 465–551.
- PUTCHKOV, A.V. & ARNDT, E. 1994. Preliminary list and key of known tiger beetle larvae (Coleoptera, Cicindelidae) of the world. *Mitt. Schweiz. Ent. Ges.* 67: 411–420.
- PUTCHKOV, A.V. & ARNDT, E. in press. Description of the larvae of *Oxycheila* DEJEAN, 1825 (Coleoptera, Cicindelidae) and the taxonomical state of the genus. *Beitr. Ent.*
- RIVALIER, E. 1971. Remarques sur la tribu des Cicindelini (Col. Cicindelidae) et sa subdivision en sous-tribus. *Nouv. Rev. Ent.* 1: 135–143.
- RODRIGUEZ, J.P., JOLY, L.J. & PEARSON, D.L. 1994. Los Escarabajos tigre (Coleoptera: Cicindelidae) de Venezuela: Su identificación, distribución e historia natural. *Bolm. Entomol. Venez. N. S.* 9: 55–120.

(received January 11, 1996; accepted after minor changes February 20, 1996)