

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 79 (2006)

Heft: 1-2

Artikel: Additions to the Phycitinae (Lepidoptera, Pyralidae) of the Galapagos Islands, Ecuador, with description of a new species of Caudellia Dyar

Autor: Landry, Bernard / Neunzig, H.H.

DOI: <https://doi.org/10.5169/seals-402906>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Additions to the Phycitinae (Lepidoptera, Pyralidae) of the Galapagos Islands, Ecuador, with description of a new species of *Caudellia* Dyar

BERNARD LANDRY^{1,3} & H.H. NEUNZIG²

¹ Muséum d'histoire naturelle, C. P. 6434, CH-1211, Geneva, Switzerland.

E-mail: bernard.landry@mhn.ville-ge.ch

² Department of Entomology, North Carolina State University, Raleigh, North Carolina, 27695-7613,
U.S.A. E-mail: h.neunzig@gte.net

³ Corresponding author

The new species *Caudellia galapagensis* (Pyralidae, Phycitinae) is described from the Galapagos Islands (Santa Cruz Island). A habitus figure of the holotype and line drawings of the male and female genitalia are included. *Unadilla maturella* (Zeller) is reported for the first time from the archipelago (Santa Cruz Island). New distribution records for 6 other species are also mentioned. The Galapagos fauna of Phycitinae now stands at 16 species although one, known from a single female here illustrated, hasn't yet been identified.

Key words: Lepidoptera, Pyralidae, Phycitinae, *Unadilla*, *Caudellia*, new species, Galapagos Islands, new distribution records

INTRODUCTION

Since the publication of Landry & Neunzig (1997) much effort has been made to document the lepidopteran fauna of the Galapagos Islands, notably by Lazaro Roque-Albelo and his colleagues at the Charles Darwin Research Station, Santa Cruz Island, Galapagos. These efforts and three 1-month collecting trips in the Galapagos by BL since 2002 have resulted in the new records of Phycitinae presented below, including the discovery of a new species of the genus *Caudellia* Dyar, described here.

The first Galapagos occurrence of *Cadra cautella* (Walker) and several new island records of Phycitinae were documented by Causton *et al.* (2006). In addition, 15 new host plant records for Galapagos Phycitinae will be presented elsewhere (Roque-Albelo *et al.*, unpublished). Tab. 1 summarizes the distribution of all Phycitinae in the Galapagos.

The following acronyms are used: BL for Bernard Landry; BMNH for The Natural History Museum (London, England); CDRS for Charles Darwin Research Station; HHN for Herb H. Neunzig; MHNG for Muséum d'histoire naturelle de Genève, Geneva, Switzerland; and RMCA for Royal Museum for Central Africa, Tervuren, Belgium.

Tab. 1. Status and distribution of the Phycitinae in the Galapagos Islands with new records in bold. N: native, E: endemic; I: introduced. Islands: Bal: Baltra, Esp: Española, Fer: Fernandina, Flo: Floreana, Gen: Genovesa, Isa: Isabela, Mar: Marchena, Pon: Pinzon, Pta: Pinta; Rab: Rabida, SCl: San Cristobal, SCz: Santa Cruz, Sey: Seymour Norte, SFe: Santa Fe, Sgo: Santiago, Wol: Wolf.

*The island records for this species in Landry & Neunzig (1997) were partly wrong.

Species	Status	Islands														
		Bal	Esp	Fer	Flo	Gen	Isa	Mar	Pon	Pta	Rab	SCl	SCz	Sey	SFe	Sgo
<i>Ancylostomia stercorea</i> (Zeller)	I			X	X	X			X		X				X	
<i>Cadra cautella</i> (Walker)	I													X		
<i>Caudellia galapagensis</i> n. sp.	E													X		
<i>Coptarthria dasypyga</i> (Zeller)	N				X		X			X			X			X
<i>Etiella zinckenella</i> (Treitschke)	I			X		X	X			X	X	X	X	X	X	X
<i>Fulrada carpasella</i> (Schaus)	E			X		X		X		X	X	X	X	X	X	X
<i>Fundella argentina</i> Dyar	I			X	X		X	X				X	X	X	X	
<i>Hypsipyla grandella</i> (Zeller)	I											X	X			
<i>Macrorrhinia pinta</i> * Land. & Neun.	E			X	X	X		X		X	X	X	X	X	X	X
<i>Nicetiodes apianellus</i> Schaus	E		X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Oryctometopia fossulatella</i> Rag.	N	X	X	X	X		X			X	X	X	X	X		X
<i>Shafferissa pumila</i> Land. & Neun.	E	X	X	X	X	X	X	X	X	X	X	X		X	X	
<i>S. galapagoensis</i> Land. & Neun.	E			X	X		X		X	X	X	X			X	X
<i>Tota galdinella</i> Schaus	E	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Unadilla maturella</i> (Zeller)	I												X			
Genus sp.	?					X										

DIAGNOSIS AND DESCRIPTION OF NEW SPECIES

Caudellia galapagensis n. sp.


(Figs 1, 4–6)

Material examined. Holotype: ♂. 1- «GALAPAGOS ISLANDS:/ Santa Cruz (Indefatigable)./ Feb. 1970/ R. Perry & Tj. de Vries/ B.M. 1970-172/ Ref. No. L.133». 2- «B.M. Pyralidae/ Genitalia slide/ ♂ No. 21117» [BMNH].

Paratypes: 4 ♂♂, 3 ♀♀ from Santa Cruz Island, Galapagos. 1 ♂ with same collection information as holotype, without abdomen [BMNH]; 2 ♀♀, Station Darwin, (lumière), X/XI-1964, J. & N. Leleup [MHNG, RMCA]; 1 ♀♀ (genitalia slide 6433 HHN), Barranco, E[stacion]. C[ientifica]. C[harles]. D[arwin].., En M[ercury]. V[apor]. L[ight].., 16 IX 1999, L. Roque [CDRS]; 2 ♂♂ (one with genitalia slide 6432 HHN), C[harles]. D[arwin]. R[esearch]. S[tation].., Barranco, M.V.L., 17 XI 1991, L. Roque [CDRS]; 1 ♂, Barranco, CDRS, 23 X 2001, UVL, L. Roque [CDRS].

Etymology. The name *galapagensis* is a combination of the type locality (Galapagos) and the Latin *-ensis*, a suffix denoting place or locality.

Diagnosis. The pair of large, crescentic lobes located at the apex of the gnathe (Fig. 4) are diagnostic for the species. In addition to this unique characteristic, *C. galapagensis* differs from other species of *Caudellia* in having the forewing mostly pale brown or pale reddish brown, rather than mostly dark brown or black. Also, the shaft of the antenna of *C. galapagensis* is more extensively covered with scales than in other species of the genus.



Figs 1–3. Galapagos Phycitinae: 1, *Caudellia galapagosensis*, holotype ♂; 2, *Unadilla maturella*, ♀; 3, Unidentified Phycitinae species.

Description. Forewing length 7.0–9.0 mm. Head: frons white, lightly washed with pale brown; scales of frons projecting anteromesially; vertex white, lightly washed with pale brown; labial palpus outwardly pale brown, lightly dusted with white, oblique, reaching slightly above vertex, third segment shorter than second segment; maxillary palpus outwardly pale brown, lightly dusted with white, short-scaled; antenna of male broader throughout than in female, with very shallow, basal sinus; antenna of female without sinus; antenna of both sexes with every other band of scales on shaft completely encircling shaft. Thorax dorsum mostly pale brown, mesially brownish white. Forewing mostly pale brown to pale reddish brown; with short, white patch at base; antemedial line white, oblique; postmedial line white, weakly developed medially; large white and pale brown patch on anterior half of wing between antemedial and postmedial lines and small patch of white scales following postmedial line; discal spots pale brown, indistinct, visible because of slight contrast between spots and large white and pale brown patch between antemedial and postmedial lines; male with costal fold, and scale tuft at basal half of wing. Hindwing mostly brownish white, brown near margins. Male genitalia (Figs 4, 5): uncus broad, rounded distally; gnathos with distal process developed as two large, crescentic lobes; transtilla strongly fused to gnathos; juxta a slender, curved band; valva well developed, broadened distally with slight costal prominence at about two-thirds distance from base of valva, and with rounded, more strongly developed costal projection apically; aedeagus simple; vesica unarmed; vinculum about as long as greatest width. Female genitalia (Fig. 6): ostium bursae sclerotized, strongly wrinkled with posteriorly flared lamella postvaginalis; ductus bursae sclerotized for about $\frac{1}{2}$ its length; corpus bursae elongate-oval with signum developed as a few, oval sclerotized discs; ductus bursae attached to corpus bursae adjacent to signum.

Remarks. *Caudellia* is a small New World group of eight species that are chiefly found in the Neotropics (Schaffer 1995; Neunzig 1990, 1996; Neunzig & Dow 1993). The genus has been treated in detail by Heinrich (1956) and Neunzig (1990). The male genitalia are unusual in that the basal elements of the transtilla are, with a few exceptions, very broad and fused to the basal parts of the gnathos. The apical part of the gnathos has either a short medial hook or a pair of enlarged lateral processes. Females have the ductus bursae heavily sclerotized for a third or more of its length from the ostium bursae, and have the ductus seminalis joined to the corpus bursae close to the signum of the corpus bursae.


OTHER NEW RECORDS

Unadilla maturella (Zeller)

(Fig. 2)

This is a new species record for the Galapagos. Only two specimens have been collected so far, both on Santa Cruz Island: 1 ♂, low agriculture zone, GPS: S 00° 42.132', W 90° 19.156', 13.iii.2004, ultra-violet light (B. Landry, P. Schmitz); 1 ♀, casa L. Roque-Albelo & V. Cruz, GPS: 137 m elev., S 00° 42.595', W 90° 19.196', 27.ii.2005, ultra-violet light (B. Landry).

The female is represented on Fig. 2. The male collected has more contrasting black forewing markings in the form of a pair of short longitudinal dashes at $\frac{1}{3}$ wing length, along midline and below, another short dash along midline at about $\frac{2}{3}$ wing



Figs 4–6. Genitalia of *Caudellia galapagensis*. 4, ♂, aedeagus omitted; 5, aedeagus; 6, ♀.

length, and a narrow line on the costa from the base to about $\frac{1}{3}$ wing length. The species was treated in detail and illustrated by Heinrich (1956) and Neunzig (1997).

The species is found in the southern United States (Florida), Mexico, the Caribbean, and Central and northern South America. Larvae have been reared from flowers and seeds of *Pluchea odorata* Cassini and *Melanthera parvifolia* Small (Asteraceae). Neither of these species occur in the Galapagos (Chris Buddenhagen, CDRS botanist, pers. comm.) so *U. maturella* must develop on other Asteraceae, for which the Galapagos are rich in endemics.

Genus, sp.

(Fig. 3)

One ♀ of this unidentified species was taken at light at Cabo Douglas on the coast of Fernandina on 15 February 2005 (B. Landry, P. Schmitz). Its wingspan is 13 mm. A male of this species would be necessary for identification.

ACKNOWLEDGMENTS

We are thankful to the authorities of the Charles Darwin Research Station and Galapagos National Park, as well as to Lazaro Roque-Albelo for providing permits, logistical support, hospitality, and partial financing to BL during expeditions to the Galapagos in 2002, 2003, and 2004. We also thank L. Roque-Albelo for allowing us to publish some of the new distribution records and for the loan of CDRS specimens. We further thank Kevin Tuck (BMNH) and J. De Prins (RMAC) for their loans of specimens in their care. The companionship and help in the field of Novarino Castillo, José Loaiza, L. Roque-Albelo, and Patrick Schmitz was greatly appreciated. The 2005 expedition to Fernandina Island was made possible by funds awarded by the «Fondation Schmidheiny» to P. Schmitz. We are also thankful for the comments of the reviewers.

REFERENCES

- Causton, C.E., Peck, S.B., Sinclair, B.J., Roque-Albelo, L., Hodgson, C.J. & Landry, B. 2006. Alien Insects: Threats and Implications for Conservation of Galapagos Islands. Annals of the Entomological Society of America 99: 121–143.
- Heinrich, C. 1956. American moths of the subfamily Phycitinae. Bulletin of the United States National Museum 207: i–viii, 1–581.
- Landry, B. & Neunzig, H.H. 1997. A review of the Phycitinae of the Galapagos Islands (Lepidoptera: Pyralidae). Entomologica scandinavica 28: 493–508.
- Neunzig, H.H. 1990. Pyraloidea, Pyralidae (part). In: Dominick, R.B., et al. (eds), The Moths of America north of Mexico, Fasc. 15.3. The Wedge Entomological Research Foundation, Washington, 165 pp.
- Neunzig, H.H. 1996. New species of Phycitinae (Lepidoptera: Pyralidae) from the Dominican Republic. Proceedings of the entomological Society of Washington 98: 774–801.
- Neunzig, H.H. 1997. Pyraloidea, Pyralidae (part). In: Dominick, R.B. et al. (eds), The moths of America North of Mexico, Fasc. 15.4. The Wedge Entomological Research Foundation, Washington, 157 pp.
- Neunzig, H.H. & Dow, L.C. 1993. The Phycitinae of Belize (Lepidoptera: Pyralidae). North Carolina Agricultural Research Service Technical Bulletin 304: 1–133.
- Shaffer, J.C. 1995. Phycitinae. pp. 93–105. In: Heppner, J.B. (ed.) Atlas of Neotropical Lepidoptera, Volume 3. Association for Tropical Lepidoptera and Scientific Publishers, Gainesville, iv + 243 pp.

(received October 31, 2005; accepted January 16, 2006)