

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 79 (2006)

Heft: 3-4

Artikel: Faunistically remarkable spiders (Arachnida : Araneae) of the timberline
in the Swiss Central Alps (Alp Flix, Grisons)

Autor: Frick, Holger / Hänggi, Ambros / Kropf, Christian

DOI: <https://doi.org/10.5169/seals-402918>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Faunistically remarkable spiders (Arachnida: Araneae) of the timberline in the Swiss Central Alps (Alp Flix, Grisons)

HOLGER FRICK^{1,2}, AMBROS HÄNGGI³, CHRISTIAN KROPF^{2,4},
WOLFGANG NENTWIG¹ & ANGELO BOLZERN³

¹ Zoological Institute, University of Bern, Baltzerstrasse 6, CH-3012 Bern

² Natural History Museum Bern, Department of Invertebrates, Bernastrasse 15, CH-3005 Bern

³ Natural History Museum Basel, Abteilung Biowissenschaften, Augustinergasse 2, CH-4001 Basel

⁴ Correspondence to Christian Kropf: christian.kropf@nmbe.unibe.ch

Spiders of the timberline on Alp Flix (Switzerland, Canton Grisons, 1960 m) were collected between June 2002 and May 2004 around and on stand-alone Norway spruce trees (*Picea abies*) by means of pitfall traps, funnel traps, frame and litter sampling, branch electrocutors, restricted canopy fogging, beating, sweep netting and hand sampling. 93 species were recorded, four of them endemic to the Alps and another six to the European mountain system. Five species are new to the Canton Grisons: *Evansia merens*, *Meioneta innotabilis*, *Moebelia penicillata*, *Obscuriphantes obscurus* and *Agroeca proxima*.

Most species can be classified as forest or open-land dwellers that contribute equally to species diversity, only few are generalists. 13 % of the species (27 % of the individuals) are winter-active. It is shown that stand-alone trees at the alpine timberline offer a broad range of habitats for a high number of spider species. This should be considered in future conservation and research efforts. Phenologies of certain species are included.

Keywords: Spiders, Araneae, faunistics, phenology, timberline, new records, Grisons, Central Alps, Switzerland.

INTRODUCTION

Faunistic data of diverse groups with differentiated habitat preferences of species represent one of the basic tools for many disciplines including conservation biology. Due to their often strict dependency on certain microclimatic factors and therefore sensitivity to environmental changes (Bauchhens 1990; Wise 1993; Foelix 1996; Samu *et al.* 1999), species associations of spiders (Araneae) have a high indicator value, reflecting the status and the «health» of a habitat (Clausen 1986; Hänggi 1998; Thaler 1998). This makes them ideal subjects of study for many conservation and ecological problems.

The Alp Flix in the Canton Grisons is a hotspot of biodiversity in Switzerland (Hänggi & Müller 2001). In spite of this, only one preliminary study has been devoted to the spider fauna at this location (Hasselmann 2000; Hänggi & Kropf 2001). It has been previously shown that the diversity of spiders at the timberline exceeds that of the surroundings (Thaler 1989; Zingerle 1999a, b). The spider associations of different areas around single trees of the timberline and those of different strata, however, have not yet been studied. The timberline on Alp Flix with its mosaic patterns of open areas and stand-alone trees is, therefore, of particular faunistic and ecological interest.



Fig. 1: Map of Switzerland with study site (modified after <http://www.lib.utexas.edu/maps/switzerland.html>; Courtesy of the University of Texas Libraries, The University of Texas at Austin).

Two ecological studies devoted to the influence of stand-alone spruce trees on the small scale distribution of spiders at the timberline on Alp Flix (results of the diploma theses of A. Bolzern and H. Frick) provided a number of faunistically remarkable spider species that are reported here. The ecological data are published elsewhere (Bolzern & Hänggi 2005, Bolzern *et al.* 2005, Frick *et al.* 2007).

METHODS

Study site

Alp Flix (GPS coordinates: 769 400 / 154 350; WGS84 coordinates: N: 46° 31' 8.41", E: 9° 38' 47.12") is situated in the Swiss Central Alps in the Canton Gri-

sons and belongs to the village of Sur (Fig. 1). The exposed terrace on the south-west of Alp Flix is approximately 1000 m wide and ranges from 1950 m a.s.l. up to 2100 m. It is bordered in the north-east by steep mountains with an elevation of over 3000 m, and in the south-west by the valley of the river Julia. The geology includes granite, gneiss, quartzite, chalk and serpentine formations. These originated from moraine material from the nearby Err massive.

The study site is situated about 550 m north-east of the hamlet Salategnas at the lower edge of the terrace, between 1950 m and 1960 m above sea level. The south western border is defined by the anthropogenically affected forest edge where occasional cattle-grazing occurs throughout the vegetation period. The site is characterised by numerous stand-alone Norway spruce trees (*Picea abies*). In the north-east, the biotope changes to dwarf shrub heath, dominated by *Juniperus communis* and *Rhododendron ferrugineum*.

The climate on Alp Flix is defined as inneralpine-continental (Hänggi & Müller 2001) with a snow-free period from May to the end of October. The temperature extremes in summer range from 0°C at night to 30°C at daytime and in winter from -20°C to ca. +15°C (Fig. 2). The snow cover ranges from a few centimetres to more than one meter.

Study period

The studies occurred from June 2002 to May 2004. Funnel traps (FUN) and nearby pitfall traps (PTF) were used from June 2002 to June 2003 (leg. A. H.), and pitfall traps (PTB) from May 2003 to May 2004 (coll. H. F.). The same timetable applied for the frame (FRS) and litter sampling (LIT) and the sweep netting (SWE), conducted during the snow free time of 2003 (coll. H. F.). During that time, spiders were collected by using branch eclectors (BRE), beatings (BEA) and restricted canopy fogging (RCF) (coll. A. B.). Hand samples (coll. H. F., A. B.) were also taken sporadically between May 2003 and May 2004 at the timberline (HST) and at nearby sites (HSS).

SAMPLING METHODS FOR SPIDERS ON TREES

Branch eclectors (BRE): The branch eclector traps are essentially constructed after Behre (1989) and Mühlenberg (1993). The traps were filled with formaldehyde (4 %); a tenside (liquid soap, non-perfumed) was added. Two eclectors were positioned on each of six Norway spruce trees between 110 cm and 430 cm above the ground. All branches sampled had a diameter of roughly 5 cm.

Beating (BEA): 84 branches were beaten 20 to 25 times at a distance between 80 cm and 500 cm from the ground. Each branch was sampled only once. The spiders were collected on a white umbrella (Ø: 85 cm) and transferred into ethanol 75 %.

Restricted Canopy Fogging (RCF): 36 branches were separately enclosed by robust transparent sealed plastic bags (height 150 cm; width 100 cm; volume approx. 400 l) at a height of up to 240 cm above ground. The bags were filled with CO₂. 20 minutes later, the branch was shaken and the bag removed. Each branch was sampled only once.

Further information on these methods is given in Bolzern & Hänggi (2005).




Fig. 2: Weather data of Radons (1870 m), opposite to Alp Flix on the other valley side. The data show average, minimum and maximum temperatures (in °C) and the total precipitation (bars, in l/m²) per week from May 2003 to May 2004.

Sampling methods for ground dwelling spiders

Funnel traps (FUN): Plastic bottles (ca. 0.75 l) were used with a funnel (Ø: 15 cm) fixed on the top. At the lower end of the funnel, a wire-cross prevented small vertebrates from falling into the trap. The traps were covered by a plastic plate at 10 cm above the ground (Obrist & Duelli, 1996). Two funnel traps (filled with 4 % formaldehyde with a detergent) were installed in opposite corners of a rectangle of 2 x 3 m together with two pitfall traps (PTF) in the other corners that were of the same type as the PTBs described below.

Pitfall traps (PTB) at sites near the branch ecollectors: White plastic cups with an upper diameter of 6.9 cm and a depth of 7.5 cm were filled with 4 % formaldehyde and 0.05 % SDS as tenside up to 3 cm under the rim. Quadrangular transparent plastic covers (15 x 15cm) were fixed at ca. 5 cm above the traps. Nine pitfall traps per tree were positioned around four trees of similar size (diameter ca. 5 m, height ca. 12 m) in three directions with three traps each. These trees were sampled with pitfall traps (PTB) and branch ecollectors (BRE) exclusively.

Frame (FRS) and litter samples (LIT): Frames (FRS): Four trees were investigated per month without any additional sampling (except LITs) in the same month. Three samples of 22 x 22 cm (5 cm depth) per tree were cut out of the soil. All samples were taken in three different directions at random places under the branches. Specimens were extracted by a MacFadyen high-gradient funnel (MacFadyen 1962). The litter samples (LIT) were taken in the same way under the same trees, but only the litter layer was collected.

Sweep netting (SWE): The vegetation under four trees was sampled every month by sweep netting, without any additional sampling in the same month. Sampling was done ca. 2 m away from the tree trunk and at the outer border of the branch cover.

Determination

Specimens were determined mostly after Nentwig et al. (2003), Roberts (1985, 1987, 1995), and Wiehle (1956, 1960). Other literature used: Bosmans & de Smet (1993); Braun (1965); Buchar (1981); Buchar & Thaler (1995); de Blauwe (1973); Denis (1965); Grimm (1985, 1986); Hänggi (1989); Heimer & Nentwig (1991); van Helsdingen *et al.* (1977, 2001); Jantscher (2001); Knoflach (1992, 1996); Kronstedt (1990); Levi (1977); Locket *et al.* (1974); Logunov (1996); Logunov & Kronstedt (2003); Lugetti & Tongiorgi (1965, 1969); Miller (1967); Millidge (1975, 1979); Ovtsharenko *et al.* (1992); Pesarini (1996, 2000); Relys & Weiss (1997); Saaristo (1971); Snazell (1983); Szita & Samu (2000); Thaler (1969, 1970, 1971, 1972a, b, 1976, 1978a, b, 1981, 1983, 1987); Thaler & Buchar (1993); Tongiorgi (1966a, b); Wiehle (1937); Wunderlich (1972, 1980, 1985, 1986); Zabka (1997).

The nomenclature follows Platnick (2005) but the families are ordered alphabetically. The material is stored in the Natural History Museums of Bern and Basel.

RESULTS AND DISCUSSION

List of species

93 species from 13 families were found (Tab. 1, see Annex), including one species, *Maro lehtineni* newly discovered for Switzerland (Bolzern *et al.* 2005) and five species, *Evansia merens*, *Meioneta innotabilis*, *Moebelia penicillata*, *Obscuriphantes obscurus* and *Agroeca proxima* new for the Canton Grisons. Four species are endemic to the Alps, i.e. *Caracladus avicula*, *Scotinotylus clavatus*, *Tenuiphantes jacksonoides* and *Walckenaeria languida*. Another six species are classified as endemic to the European mountain system: *Erigonella subelevata*, *Arctosa renidescens*, *Pardosa blanda*, *Talavera monticola*, *Robertus truncorum* and *Xysticus macedonicus*. See Tab. 1 for a detailed list of all species.

Comments on first records, endemic and rare species and their phenology

Araneidae

Araniella displicata (Hentz, 1847)

Determination: Roberts (1985).

Comments: Only known locality of this species in Grisons (Hänggi 2000). All records in June.

Gnaphosidae

Drassodes cupreus (Blackwall, 1834)

Determination: Roberts (1995), Grimm (1985).

Comments: According to Thaler (1981, 2004), most high-altitude records of *D. lapidosus* should refer to *D. cupreus* (Blackwall, 1834). The species status of *D. lapidosus* and *D. cupreus* is under debate (e.g. Grimm 1985). See the detailed discussion by Bolzern & Hänggi (2006). Specimens were found between mid May and July.

Micaria aenea Thorell, 1874

Determination: Wunderlich (1980), Miller (1967).

Habitat: Most observations occurred in the subalpine belt between 1500 and 2000 m; in dry and sunny habitats; clearings, meadows, birch fen woods, coniferous forests, timberline and dwarf shrub heath (Hänggi *et al.* 1995; Muster 2001; Thaler 2004).

Distribution: Holarctic, eurosibirian (Thaler 2004); after Thaler (1997a; 1998; 2004) arctoalpine to boreomontane distribution.

Comments: Rarely found in Switzerland (Maurer & Hänggi 1990). The highest activity was observed in May and June (Fig. 3a).

Linyphiidae

Caracladus avicula (L. Koch, 1869)

Determination: Thaler (1969, 1972b).

Habitat: Subalpine belt; from subalpine coniferous forest up to the timberline (~1000–2000 m) (Thaler, 1999); epigeal (Maurer & Hänggi 1990); in forests and meadows (Hänggi *et al.* 1995; Muster 2001).

Distribution: Endemic to the Alps (Muster 2001; Thaler 1999).

Comments: Usually found only in small numbers. Diplochronous after Thaler




Fig. 3a-h: Phenologies of selected species for males (black) and females (grey) separately. The graphics are based on the data of Tab. 1 excluding the hand samplings. The sampling data were fit into the sampling periods of the pitfall traps (PTB). Abbreviations: May: 15. 05. 2003–14. 06. 2003; Jun: 14. 06. 2003–07. 07. 2003; Jul: 07. 07. 2003–04. 08. 2003; Aug: 04. 08. 2003–03. 09. 2003; Sep: 03. 09. 2003–28. 09. 2003; Oct: 28. 09. 2003–28. 10. 2003 and Win: 28. 10. 2003–24. 05. 2004.

(1999); however, main activity should be in lower altitudes between the end of March and mid May, in higher altitudes between June and September (Thaler 1972b). Muster (2001) found most of the specimens in June and July at similar sites. Here, the highest activity was observed between the beginning of November and mid June (Fig. 3b). So, the phenology of this species should be studied in more detail – on Alp Flix no diplochrony is detectable.

Evansia merens O. P.-Cambridge, 1900

Determination: Wiehle (1960).

Habitat: Myrmecophilous (Thaler *et al.* 1990); nidicolous in ant nests; mostly associated with ants from the *Formica fusca* group; sometimes under stones in dwarf shrub heath between 1900 and 2200 m (Thaler 1999).

Distribution: West-palaeartic, possibly extra-mediterranean (Thaler 1999).

Comments: First record for Grisons, second record for Switzerland after Schenkel's (1936) finding. We found two specimens in May at a site with a high abundance of *Formica* sp. Adults found during the whole year; most individuals found in autumn (Wiehle 1960).

Macrargus carpenteri O. P.-Cambridge, 1900

Determination: Wiehle (1956).

Habitat: Habitat preferences unclear; seems to be xerophilous (Thaler 1983). Thaler (1995b) found the species at the timberline and in the lower alpine zone (in dwarf shrubs and grass meadow); epigeal (Maurer & Hänggi 1990; Hänggi *et al.* 1995).

Distribution: Central Europe and West Siberia; the Alps mark the southern distribution border (Thaler 1995b).

Comments: *M. carpenteri* seems to be mature and active in winter (Thaler 1995b); we found males and females between the beginning of October and May exclusively (Fig. 3c). Our record is the second in Switzerland (Maurer & Hänggi 1990). A check of the material of *M. rufus* (Wider, 1834), a species that had formerly been confused with *M. carpenteri*, in the museums of Bern and Basel, revealed no misidentifications.

Mansuphantes pseudoarciger (Wunderlich, 1985)

Determination: Wunderlich (1985), Heimer & Nentwig (1991).

Habitat: montane; epigeal; in moderate humid moss of a *P. abies* forest. (Maurer & Hänggi 1990).

Distribution: Western Alps; France and Switzerland (Heimer & Nentwig 1991).

Comments: second observation in Switzerland, after Müller's (1985) finding; first record for Grisons. Adults were found nearly exclusively between November and early June (Fig. 3d). 70 % of all specimens in pitfall traps were caught in a range of about 1 m around tree trunks of *P. abies*.

Meioneta innotabilis O. P.-Cambridge, 1900

Determination: Roberts (1987).

Habitat: On bark of spruce (Muster 2001); from low altitudes up to the timberline (Thaler 1995b), also in cultivated spruce forests (Hänggi *et al.* 1995).

Distribution: Europe (Muster 2001).

Comments: One male between mid May and early June in branch eclector.




Fig. 4: Prosoma of male *Metopobactrus schenkeli* Thaler, 1976 from Alp Flix. Drawing by A. Bolzern.

First recorded observation for Grisons and first record since Schenkel (1923). He found males in June in a vegetable field close to Basel. Findings by Lessert (1907, 1910) were under tree bark of spruce trees in June in Bad Ragaz (St. Gallen) and Bex (Vaud).

Metopobactrus schenkeli Thaler, 1976

Determination: Thaler (1976), Hänggi (1989), Nentwig *et al.* (2003).

Habitat: In leaf litter of dwarf shrub heath and meadows (Maurer & Hänggi 1990).

Distribution: In the Swiss and Italian Alps, including northern Italy (Hänggi 1989; Platnick 2005) mostly in higher regions up to the nival zone (Maurer & Hänggi 1990). Findings in and around Venice (Thaler 1976; Hansen 1995) suggest that this species is also present in lower altitudes. Thaler (1976, 1978a) discusses the southern distribution of *M. schenkeli* and vicariance with *M. prominulus* (O. P.-Cambridge, 1872).

Comments: We found two males in funnel traps (FUN) in early July at the forest edge and one male in a pitfall trap (PTB) under *P. abies* in winter. The males of *M. schenkeli* differ from those of *M. prominulus* by the elevated head region (Fig. 4) and from those of *M. nadigi* Thaler, 1976 and *M. ascitus* (Kulczyński, 1894) by the shape of the tibia of the pedipalp and position of the trichobothria on metatarsi I and IV (position in our specimens: 0.8 on metatarsus I and between 0.8 and 0.82 on metatarsus IV).

Our record is from the drainage area of the river Rhine and is together with findings near Davos (Maurer & Walter 1984) the most northern one. So, *M. schenkeli* is more widely distributed than expected and could occur in sympatry with *M. prominulus*.

Mughiphantes cornutus Schenkel, 1927

Determination: Thaler (1972), Heimer & Nentwig (1991).

Habitat: Mainly distributed in subalpine forest between 1500 and 2000 m (Thaler 1995b); only a few specimens found in higher subalpine belt in the Alps (Thaler 1972a); epigeal; in forest litter (Maurer & Hänggi 1990).

Distribution: Boreo-montane/boreo-alpine and also in South Siberia (Thaler 1995b).

Comments: The main activity was observed between May and August, covering nearly the whole snow-free period (Fig. 3e). After Thaler (1972a) the species should be diplochronous. 85 % of all observed specimens from pitfall traps were caught directly at the tree trunk or in a range of about 1 m around the tree trunk of *P. abies*.

Panamomops tauricornis (Simon, 1981)

Determination: Wiehle (1960).

Distribution: Previously considered to be endemic to the Alps but recently also recorded from North Asia (Eskov 1994; Thaler 1999).

Comments: Rare species, diplochronous (Thaler 1995a). We found adult specimens all through the year with the main activity between mid May and end of July (Fig. 3f).

Porrhomma cf. campbelli O. P.-Cambridge, 1900

Determination: Thaler (1991), Roberts (1987).

Comments: Confusion with *P. pallidum* (22 males, 15 females in our sample) possible. Determination of one intermediate female uncertain. If this species is correctly identified, it would be the second recorded observation in Grisons (Maurer & Hänggi 1990).

Scotinotylus clavatus (Schenkel, 1927)

Determination: Thaler (1970).

Distribution: Endemic to the Alps with Nearctic twin species *Scotinotylus sacer* (Crosby, 1929) (Thaler 1999).

Comments: Diplochronous after Thaler (1999); females long-living and overwintering (Thaler 1995a); adults were observed in all months, males had the highest activity between July and September, diplochrony was not detectable. Females were equally common between September and mid June (Fig. 3g). Most specimens close to the tree trunks of *P. abies*.

Stemonyphantes conspersus (L. Koch, 1879)

Determination: Thaler (1983), Roberts (1987).

Comments: Second recorded observation in Switzerland and Grisons, first one by Maurer & Hänggi (1989); mainly active in winter (Muster 2001); most specimens were found between October and early June (Fig. 3h).

Tenuiphantes jacksonoides (van Helsdingen, 1977)

Determination: van Helsdingen *et al.* (1977).

Habitat: Subalpine spruce forest up to grass heath, 1500–2500 m (Thaler 1995b); more in open vegetation than in forests (van Helsdingen *et al.* 1977); epigeal (Maurer & Hänggi 1990);

Distribution: Narrow endemic of the Eastern Alps (Thaler 1995b; Muster 2001).

Comments: Second recorded observation in Grisons, first after van Helsdingen *et al.* (1977), third record for Switzerland. Adults are found from June to Sep-




Fig. 5: Male palp of *Tenuiphantes jacksonoides* (van Helsdingen, 1979); prolateral view. Drawing by H. Frick.

tember (van Helsdingen *et al.* 1977). Our specimen is slightly different from the drawing of van Helsdingen *et al.* (1977); it lacks one process on the inner part of the lamella (Fig. 5). Apart from this, the male palp fits well to *T. jacksonoides*.

Lycosidae

Pardosa riparia (C. L. Koch, 1833)

Determination: Tongiorgi (1966a), Tongiorgi (1966b).

Habitat: In light forests of pine, spruce, and larch, with high grass below, up to 2000 m (Thaler 1995a; 1997a); epigeal; mostly humid meadows in higher altitudes (Maurer & Hänggi 1990; Hänggi *et al.* 1995); mountain pastures (Muster 2001).

Distribution: Palaearctic (Maurer & Hänggi 1990).

Comments: The eudominant species at the timberline of Alp Flix. The maximum activity occurred between mid May and July (Fig. 3h). Females displayed high variability in colour and shape of the epigyne. However, species specific traits according to Tongiorgi (1966a; b) were clearly visible.

Thomisidae

Xysticus audax (Schrank, 1803) / *X. macedonicus* Šilhavý, 1944

Determination: Jantscher (2001).

Habitat: *X. audax*: euryzonic forest species; dry and wet meadows; epigeal and arboricolous up to the dwarf shrub heath to ca. 2100 m, sometimes in open sites and above the timberline (Maurer & Hänggi 1990; Muster 2001; Thaler 1997b; 2004).

X. macedonicus: Euryzonic up to 2400 m, in the Alps up to 2160 m, thermophilous, open-land, alpine grass, debris, dry forests, mixed deciduous forests, epigeal (Kropf & Horak 1996; Thaler 1997b; Muster 2000).

Distribution: *X. audax* in Europe, Siberia, Asia Minor, Japan, Sachalin (Maurer & Hänggi 1990); *X. macedonicus* is endemic to the European mountain system (Muster 2001).

Comments: The females of *X. audax* can be easily confused with *X. macedonicus* and *X. cristatus* (Clerck, 1757). Several females appeared to be intermediate between *X. audax* and *X. macedonicus* and were counted as *X. audax*; however, one female clearly matched the drawings of *X. macedonicus* (Jantscher 2001). Males are mostly distinct (Jantscher 2001). Still, it seems that even after the elaborate work by Jantscher (2001) problems to distinguish these two species remain. *X. audax* and *X. macedonicus* occur syntopically but with different microhabitat preferences (Hänggi 2003; Muster 2000). Second record from Switzerland, first one also on Alp Flix (Hänggi 2003).

CONCLUSION

The timberline on Alp Flix has been shown to harbour a rich community of spider species with diverse ecological preferences, alpine-endemic species and faunistic rarities. Presumably, this is also true for other invertebrates. The spectrum of species covers only few generalists. There are equal numbers of forest and open-land species; together they comprise roughly 90 % of the species. 3 species can be classified as myrmecophilous. $\frac{2}{3}$ of the species are active during the vegetation period, 12 species (13 %) are winter-active and remarkably represent 27 % of the total number of individuals (the remaining species are active the whole year or are badly known in this respect).

24 % of the species and 76 % of the individuals were collected by pitfall traps exclusively (Tab. 1). This is most probably due to the fact that pitfall trapping was used more intensively than the other methods. So, the assessed fauna is clearly biased by the sampling method. Future studies and conservation efforts should pay special attention to the timberline - a major part of the invertebrate diversity of the region could possibly be maintained in this relatively small area.

ACKNOWLEDGEMENTS

We are deeply indebted to all the people who helped us during our field work, especially to Victoria Spinaz (Sur, CH) for her support on the alp. Special thanks go to the «Schatzinsel Alp Flix» foundation and to Dr. Jürg Paul Müller (Chur, CH) who made available accommodation at the study site. The late Prof. Dr. Konrad Thaler (Innsbruck, A) kindly checked the determination of *Scotargus pilosus*, *Metopobactrus schenkeli* and *Tenuiphantes jacksonoides* which is greatly acknowledged. We are grateful to Theo Blick (Hummelthal, D) for important comments on an earlier version of the manuscript. For technical help, material and fruitful discussions we thank Dr. Denise Wyniger (Luzern, CH), Roland Mühlethaler (Basel, CH), Prof. Dr. Jürg Zettel (Bern, CH) and Alain Jacob (Bern, CH). We are also grateful to Gianni Savoldelli (Randons, CH) for the weather data and Halinka Lamparski (Perth, AUS) and Tristan Derham (Perth, AUS) for correcting the English of an earlier version of the manuscript. H.F. cordially thanks his parents for continuous help and support.

ZUSAMMENFASSUNG

Von Juni 2002 bis Mai 2004 wurden an der Waldgrenze auf Alp Flix (Schweiz, Graubünden, 1960 m) Spinnen um und auf einzeln stehenden Fichten (*Picea abies*) gesammelt. Dabei kamen verschiedene Methoden wie Barberfallen, Trichterfallen, Stech- und Gesiebeproben, Asteklektoren, CO₂-

Begasungen, Klopfschirme, Netzfänge und Handaufsammlungen zum Einsatz. 93 Arten wurden erfasst, darunter vier Endemiten der Alpen und sechs des Europäischen Gebirgssystems. Fünf Arten waren neu für den Kanton Graubünden: *Evansia merens*, *Meioneta innotabilis*, *Moebelia penicillata*, *Obscuriphantes obscurus* und *Agroeca proxima*.

Abgesehen von ein paar Generalisten konnten die meisten Arten entweder als Offenland- oder Waldarten klassifiziert werden. Beide Typen waren etwa gleich häufig. 13% der Arten (27% der Individuen) sind winteraktiv. Es konnte gezeigt werden, dass einzeln stehende Bäume der alpinen Übergangszone ein breites Spektrum von Habitaten für eine hohe Anzahl Spinnenarten bieten. Dies sollte in zukünftigen Schutz- und Forschungsprojekten berücksichtigt werden. Für einige Arten wurden Phänologien erstellt.

REFERENCES

- Bauchhenss, E. 1990. Mitteleuropäische Xerotherm-Standorte und ihre epigäische Spinnenfauna – eine autökologische Betrachtung. — Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg (NF) 31/32: 153–162.
- Behre, G. F. 1989. Freilandökologische Methoden zur Erfassung der Entomofauna (Weiter- und Neuentwicklung von Geräten). — Jahresberichte des Naturwissenschaftlichen Vereins in Wuppertal 42, 238–242.
- de Blauwe, R. 1973. Révision de la famille des Agelenidae (Araneae) de la région méditerranéenne. — Bulletin de l'Institut Royal des Sciences Naturelles de Belgique 49: 1–111.
- Bolzern, A. & Hänggi, A. 2005. Spinnenfänge (Arachnida, Araneae) auf subalpinen Fichten der Alp Flix (GR, Schweiz) – ein Methodenvergleich. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 78: 1–10.
- Bolzern, A., Hänggi, A., Kropf, C. & Frick, H. 2005. Erstnachweis von *Maro lehtineni* Saaristo 1971 für die Schweiz. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 78: 165–172.
- Bolzern, A. & Hänggi, A. 2006. *Drassodes lapidosus* und *Drassodes cupreus* (Araneae: Gnaphosidae) – eine unendliche Geschichte. — Arachnologische Mitteilungen 31: 16–22.
- Bosmans, R. & Smet, K. de 1993. Le genre *Walckenaeria* Blackwall en Afrique du Nord (Araneae, Linyphiidae). Etudes sur les Linyphiidae nord-africaines, I. — Revue Arachnologique 10 (2): 21–51.
- Braun, R. 1965. Beitrag zu einer Revision der paläarktischen Arten der *Philodromus aureolus*-Gruppe (Arach., Araneae). I. Morphologisch-systematischer Teil. — Senckenbergiana biologica 46: 369–428.
- Buchar, J. 1981. Zur Lycosiden-Fauna von Tirol (Araneae, Lycosidae). — Vestník Československe Spolecnosti Zoologické 45: 4–13.
- Buchar, J. & Thaler, K. 1995. Die Wolfspinnen von Österreich 2: Gattungen *Arctosa*, *Tricca*, *Trochosa* (Arachnida, Araneida: Lycosidae). Faunistisch-tiergeographische Übersicht. — Carinthia II. 185 (105): 481–498.
- Clausen, I.H.S. 1986. The use of spiders as ecological indicators. — Bulletin of the British arachnological Society 7: 83–86.
- Denis, J. 1965. Notes sur les Erigonides. XXX. Le genre *Minicia* Thorell. — Bulletin de la Société d'Histoire Naturelle de Toulouse 100: 181–205.
- Escov, K.Y. 1994. Catalogue of the linyphiid spiders of northern Asia (Arachnida, Araneae, Linyphiidae). — Pensoft Publishers, Sofia and Moscow, 144 pp.
- Foelix, R.F. 1996. Biology of spiders. — Oxford University Press. New York. pp. 1–330.
- Frick, H., Nentwig, W. & Kropf, C. 2007. Influence of stand-alone trees on epigeic spiders (Araneae) at the alpine timberline. — Annales Zoologici Fennici 44: In press.
- Grimm, U. 1985. Die Gnaphosidae Mitteleuropas (Arachnida, Araneae). — Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg (NF) 26: 1–318.
- Grimm, U. 1986. Die Clubionidae Mitteleuropas. Corinninae und Liocraninae (Arachnida, Araneae). — Abhandlungen des Naturwissenschaftlichen Vereins in Hamburg (NF) 27: 1–91.
- Hänggi, A. 1989. Beiträge zur Kenntnis der Spinnenfauna des Kt. Tessin II – Bemerkenswerte Spinnenfunde aus Magerwiesen der Montanstufe. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 62: 167–174.
- Hänggi, A. 1998. Bewertung mit Indikatoren versus Erfassung des gesamten Artenspektrums - ein Konfliktfall? — Laufener Seminarbeiträge 8: 33–42.
- Hänggi, A. 2003. Nachträge zum «Katalog der schweizerischen Spinnen» - 3. Neunachweise von 1999 bis 2002 und Nachweise synanthroper Spinnen. — Arachnologische Mitteilungen 26: 36–49.
- Hänggi, A., Stöckli, E. & Nentwig, W. 1995. Lebensräume Mitteleuropäischer Spinnen. Charakterisierung der Lebensräume der häufigsten Spinnenarten Mitteleuropas und der mit diesen vergesellschafteten Arten. — Miscellanea Faunistica Helvetiae 4: 1–460.

- Hänggi, A. & Kropf, C. 2001. Erstnachweis der Zwergspinne *Micrargus alpinus* für die Schweiz – Mit Bemerkungen zur Bedeutung von Museumssammlungen und den Grenzen der Aussagekraft von Literaturangaben. — Jahresbericht der Naturforschenden Gesellschaft Graubünden 110: 45–49.
- Hänggi, A. & Müller, J.P. 2001. Eine 24-Stunden Aktion zur Erfassung der Biodiversität auf der Alp Flix (Grisons): Methoden und Resultate. — Jahresbericht der Naturforschenden Gesellschaft Graubünden 110: 5–36.
- Hansen, H. 1995. Über die Arachniden-Fauna von urbanen Lebensräumen in Venedig - III. Die epigäischen Spinnen eines Stadtparkes (Arachnida: Araneae). — Bollettino del Museo Civico di Storia Naturale di Venezia 44: 7–36.
- Hasselmann, A. 2000. Alp Flix - Das Ergebnis. — In: Supplement zum 2. Geotag der Artenvielfalt, GEO 9, pp. 7–23. Gruner und Jahr, Hamburg.
- Heimer, S. & Nentwig, W. 1991. Spinnen Mitteleuropas. — Paul Parey Verlag, Berlin und Hamburg. pp. 1–543.
- van Helsdingen, P.J., Thaler, K. & Deltshv, C. 1977. The *tenuis* group of *Lepthyphantes* Menge (Araneae, Linyphiidae). — Tijdschrift voor Entomologie 23: 1–54.
- van Helsdingen, P.J., Thaler, K. & Deltshv, C. 2001. The European species of *Bolyphantes* with an attempt of a phylogenetic analysis (Araneae Linyphiidae). — Memorie della Società entomologica italiana 80: 3–35.
- Jantscher, E. 2001. Diagnostic characters of *Xysticus cristatus*, *X. audax* and *X. macedonicus* (Araneae: Thomisidae). — Bulletin of the British arachnological Society 12 (1): 17–25.
- Knoflach, B. 1992. Neue *Robertus*-Funde in den Alpen: *R. mediterraneus* Eskov und *Robertus* sp. (Arachnida, Aranei: Theridiidae). — Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck 79: 161–171.
- Knoflach, B. 1996. Die Arten der *Steatoda phalerata*-Gruppe in Europa (Arachnida: Araneae, Theridiidae). — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 69: 377–404.
- Kronstedt, T. 1990. Separation of two species standing as *Alopecosa aculeata* (Clerck) by morphological, behavioural and ecological characters, with remarks on related species in the *pulverulenta* group (Araneae, Lycosidae). — Zoologica Scripta 19(2): 203–225.
- Kropf, C. & Horak, P. 1996. Die Spinnen der Steiermark (Arachnida, Araneae). — Mitteilungen des Naturwissenschaftlichen Vereines für Steiermark. Sonderheft: 5–112.
- Lessert, R. de 1907. Notes arachnologiques. — Revue Suisse de Zoologie 15: 93–128.
- Lessert, R. de 1910. Catalogue des invertébrés de la Suisse. Fascicule 3, Araignées. — Musée d'histoire naturelle de Genève, 635 pp.
- Levi, H.W. 1977. The orb-weaver genera *Metepeira*, *Kaira* and *Aculepeira* in America North of Mexico (Araneae: Araneidae). — Bulletin of the Museum of Comparative Zoology 148 (5): 185–238.
- Locket, G. H., Millidge, A. F. & Merrett, P. 1974. British spiders vol. 3. — The Ray Society, London, 314 pp.
- Logunov, D. V. 1996. A critical review of the spider genera *Apollophanes* O. P.-Cambridge, 1898 and *Thanatus* C. L. Koch, 1837 in North Asia (Araneae, Philodromidae). — Revue Arachnologique 11: 133–202.
- Logunov, D. V. & Kronstedt, T. 2003. A review of the genus *Talavera* Peckham and Peckham, 1909 (Araneae, Salticidae). — Journal of Natural History 37: 1091–1154.
- Lugetti, G. & Tongiorgi P. 1965. Revisione delle specie italiane dei generi *Artosa* C. L. Koch e *Tricca* Simon con note su una *Acantholycosa* delle Alpi Giulie (Aran. Lycosidae). — Redia 49: 165–228.
- Lugetti, G. & Tongiorgi, P. 1969. Ricerche sul genere *Alopecosa* Simon (Araneae - Lycosidae). — Atti della società Toscana di Science naturali Memorie 76 (B): 1–100.
- MacFadyen, W.A., 1962. Soil arthropod sampling. — Advances in Ecological Research 1: 1–34.
- Maurer, R. & Hänggi, A. 1989. Für die Schweiz neue und bemerkenswerte Spinnen (Araneae) III. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 62: 175–182.
- Maurer, R. & Hänggi, A. 1990. Katalog der schweizerischen Spinnen. — Documenta Faunistica Helvetiae 12.
- Maurer, R. & Walter, J.E. 1984. Für die Schweiz neue und bemerkenswerte Spinnen (Araneae) II. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 57: 65–73.
- Miller, F. 1967. Studien über Kopulationsorgane von *Zelotes*, *Micaria*, *Robertus* und *Dipoena*. — Prirodovedne Prace Ustavu Ceskoslovenske Akademie ved v Brne (N.S.) 1 (7): 253–296.
- Millidge, A. F. 1975. Re-examination of the erigonine spiders «*Micrargus herbigradus*» and «*Pocadicnemus pumila*» (Araneae, Linyphiidae). — Bulletin of the British arachnological Society 3: 145–155.

- Millidge, A. F. 1979. Some erigonine spiders from southern Europe. — *Bulletin of the British arachnological Society* 4: 316–328.
- Mühlenberg, M. 1993. *Freilandökologie*. 3. Auflage. — Quelle & Meyer (UTB), Heidelberg and Wiesbaden, 512 pp.
- Müller, H.-G. 1985. Die moosbewohnenden Araneen eines Fichtenwaldes am Col de la Forclaz - ein Beitrag zur schweizerischen Spinnenfauna (Arachnida: Araneae). — *Entomologische Zeitschrift* 95: 268–272.
- Muster, C. 2000. Weitere für Deutschland neue Spinnentiere aus dem bayerischen Alpenraum (Araneae: Linyphiidae, Agelenidae, Thomisidae, Salticidae). — *Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck* 87: 209–219.
- Muster, C. 2001. Biogeographie von Spinnentieren der mittleren Nordalpen (Arachnida: Araneae, Opiliones, Pseudoscorpiones). — *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg* 39: 5–196.
- Nentwig, W., Hänggi, A., Kropf, C. & Blick, T. 2003. Central European Spiders. An internet identification key. V. 08.12.2003. Online at: <http://www.araneae.unibe.ch>
- Obrist, M. K. & Duelli, P. 1996. Trapping efficiency of funnel- and cup-traps for epigeal arthropods. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 69: 361–369.
- Ovtsharenko, V.I., Platnick, N.I. & Song, D.X. 1992. A review of the North Asian ground spiders of the genus *Gnaphosa* (Araneae, Gnaphosidae). — *Bulletin of the American Museum of Natural History* 212: 1–88.
- Pesarini, C. 1996. Note su alcuni Erigonidae italiani, con descrizione di una nuova specie (Araneae). — *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano* 135: 413–429.
- Pesarini, C. 2000. Contributo alla conoscenza della fauna araneologica italiana (Araneae). — *Memorie della Società Entomologica Italiana* 78: 379–393.
- Platnick, N. I. 2005. The world spider catalog. V. 5.5. American Museum of Natural History. Online at: <http://research.amnh.org/entomology/spiders/catalog/index.html>.
- Relys, V. & Weiss, I. 1997. *Micrargus alpinus* sp. n., eine weitere Art der *M. herbigradus*-Gruppe aus Österreich (Arachnida: Araneae: Linyphiidae). — *Revue Suisse de Zoologie* 104: 491–501.
- Roberts, M. J. 1985. The spiders of Great Britain and Ireland. Volume 1: Atypidae to Theridiosomatidae. — Harley Books, Colchester, 229 pp.
- Roberts, M. J. 1987. The spiders of Great Britain and Ireland. Volume 2: Linyphiidae and checklist. — Harley Books, Colchester, 204 pp.
- Roberts, M. J. 1995. Spiders of Britain and Northern Europe. — Harper Collins Publishers, London, 383 pp.
- Saaristo, M. I. 1971. Revision of the genus *Maro* O. P.-Cambridge (Araneae, Linyphiidae). — *Annales Zoologici Fennici* 8: 463–482.
- Samu, F., Sunderland, K.D. & Szinetar, C. 1999: Scale-dependant dispersal and distribution patterns of spiders in agricultural systems: a review. — *Journal of Arachnology* 27: 325–332.
- Schenkel, E. 1923. Beitrag zur Spinnenkunde. — *Verhandlungen der Naturforschenden Gesellschaft in Basel* 34: 78–127.
- Schenkel, E. 1936. Kleine Beiträge zur Spinnenkunde. — *Revue Suisse de Zoologie* 43: 307–333.
- Snazell, R. 1983. On two spiders recently recorded from Britain. — *Bulletin of the British arachnological Society* 6: 93–98.
- Szita, E. & Samu, F. 2000. Taxonomical review of *Thanatus* species (Philodromidae, Araneae) of Hungary. — *Acta Zoologica Academiae Scientiarum Hungaricae* 46: 155–179.
- Thaler, K. 1969. Über einige wenig bekannte Zwergspinnen aus Tirol (Arachn., Araneae, Erigonidae). — *Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck* 57: 195–219.
- Thaler, K. 1970. Über einige wenig bekannte Zwergspinnen aus den Alpen (Arachn., Araneae, Erigonidae). — *Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck* 58: 255–276.
- Thaler, K. 1971. Über drei wenig bekannte hochalpine Zwergspinnen (Arachn., Aranei, Erigonidae). — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 44: 309–322.
- Thaler, K. 1972a. Über vier wenig bekannte *Leptyphantès*-Arten der Alpen (Arachnida, Aranei, Linyphiidae). — *Archives des Sciences* 25 (3): 289–308.
- Thaler, K. 1972b. Über einige wenig bekannte Zwergspinnen aus den Alpen, II (Arachnida: Aranei, Erigonidae). — *Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck* 59: 29–50.
- Thaler, K. 1976. Über wenig bekannte Zwergspinnen aus den Alpen, IV (Arachnida, Aranei, Erigonidae). — *Archives des Sciences* 29: 227–246.
- Thaler, K. 1978a. Über wenig bekannte Zwergspinnen aus den Alpen, V (Arachnida: Aranei, Erigonidae). — *Beiträge zur Entomologie* 28: 183–200.

- Thaler, K. 1978b. Die Gattung *Cryphoeca* in den Alpen (Arachnida, Aranei, Agelenidae). — Zoologischer Anzeiger 200: 334–346.
- Thaler, K. 1981. Bemerkenswerte Spinnenfunde in Nordtirol (Österreich) (Arachnida: Aranei). — Veröffentlichungen des Tiroler Landesmuseums Ferdinandeum 61: 105–150.
- Thaler, K. 1983. Bemerkenswerte Spinnenfunde in Nordtirol (Österreich) und Nachbarländern: Deckennetzspinnen, Linyphiidae (Arachnida: Aranei). — Veröffentlichungen des Tiroler Landesmuseums Ferdinandeum 63: 135–167.
- Thaler, K. 1987. Über einige Linyphiidae aus Kashmir (Arachnida: Araneae). — Courier Forschungs-institut Senckenberg 93: 33–42.
- Thaler, K. 1989. Epigäische Spinnen und Weberknechte (Arachnida, Aranei, Opiliones) im Bereich des Höhentransektes Glocknerstrasse - Südabschnitt (Kärnten, Österreich). — Veröffentlichungen des österreichischen MAB-Hochgebirgsprogrammes Hohe Tauern 13: 201–215.
- Thaler, K. 1995a. Oekologische Untersuchungen im Unterengadin 15. Lieferung D11. Spinnen (Araneida) mit Anhang über Weberknechte (Opiliones). — Ergebnisse der wissenschaftlichen Untersuchungen im Schweizerischen Nationalpark. 12: D473–D538.
- Thaler, K. 1995b. Beiträge zur Spinnenfauna von Nordtirol. 5. Linyphiidae 1. Linyphiinae (sensu Wiehle) (Arachnida: Araneida). — Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck. 82: 153–190.
- Thaler, K. 1997a. Beiträge zur Spinnenfauna von Nordtirol – 3: «Lycosaeformia» (Agelenidae, Hahnidae, Argyronetidae, Pisauridae, Oxyopidae, Lycosidae) und Gnaphosidae (Arachnida: Araneae). — Veröffentlichungen des Tiroler Landesmuseums Ferdinandeum 75/76: 97–146.
- Thaler, K. 1997b. Beiträge zur Spinnenfauna von Nordtirol – 4. *Dionycha* (Anyphaenidae, Clubionidae, Heteropodidae, Liocranidae, Philodromidae, Salticidae, Thomisidae, Zoridae). — Veröffentlichungen des Tiroler Landesmuseums Ferdinandeum. 77: 233–285.
- Thaler, K. 1998. Die Spinnen von Nordtirol (Arachnida, Araneae): Faunistische Synopsis. — Veröffentlichungen des Tiroler Landesmuseums Ferdinandeum 78: 37–58.
- Thaler, K. 1999. Beiträge zur Spinnenfauna von Nordtirol - 6. Linyphiidae 2: Erigoninae (sensu Wiehle) (Arachnida: Araneae). — Veröffentlichungen des Tiroler Landesmuseums Ferdinandeum 79: 215–264.
- Thaler, K. 2004. Zur Faunistik der Spinnen (Araneae) von Österreich: Gnaphosidae, Thomisidae (*Dionycha* pro parte). — Linzer biologische Beiträge 36 (1): 417–484.
- Thaler, K., Kofler, A. & Meyer, E. 1990. Fragmenta Faunistica Tirolensia - IX (Arachnida: Aranei, Opiliones; Myriapoda: Chilopoda, Diplopoda: Glomerida; Insecta: Dermaptera, Coleoptera: Staphylinidae). — Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck. 77: 225–243.
- Thaler, K. & Buchar, J. 1993. Eine verkannte Art der Gattung *Lepthyphantes* in Zentraleuropa: *L. tripartitus* Miller & Svaton (Araneida: Linyphiidae). — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 66: 149–158.
- Tongiorgi, P. 1966a. Italian wolf spiders of the genus *Pardosa* (Araneae: Lycosidae). — Bulletin of the Museum of Comparative Zoology 134: 275–334.
- Tongiorgi, P. 1966b. Wolf spiders of the *Pardosa monticola* group (Araneae, Lycosidae). — Bulletin of the Museum of Comparative Zoology 134: 335–359.
- Wiehle, H. 1937. Spinnentiere oder Arachnoidea, VIII. 26: Familie Theridiidae oder Haubennetzspinnen (Kugelspinnen). — Die Tierwelt Deutschlands und der angrenzenden Meeresteile 33, Gustav Fischer Verlag, Jena, pp.119–222.
- Wiehle, H. 1956. Spinnentiere oder Arachnoidea, X. 28: Familie Linyphiidae-Baldachinspinnen. — Die Tierwelt Deutschlands und der angrenzenden Meeresteile 44, Gustav Fischer Verlag, Jena, 337 pp.
- Wiehle, H. 1960. Spinnentiere oder Arachnoidea, XI. Micryphantidae-Zwergspinnen. — Die Tierwelt Deutschlands und der angrenzenden Meeresteile 47. — Gustav Fischer Verlag, Jena, 620 pp.
- Wise, D. 1993. Spiders in ecological webs. — Cambridge University Press, Cambridge; p.328.
- Wunderlich, J. 1972. Zur Kenntnis der Gattung *Walckenaeria* Blackwall, 1833 unter besonderer Berücksichtigung der europäischen Subgenera und Arten (Arachnida: Araneae: Linyphiidae). — Zoologische Beiträge 18: 371–472.
- Wunderlich, J. 1980. Revision der europäischen Arten der Gattung *Micaria* Westring, 1851, mit Anmerkungen zu den übrigen paläarktischen Arten (Arachnida: Araneida: Gnaphosidae). — Zoologische Beiträge 25: 233–341.
- Wunderlich, J. 1985. *Lepthyphantes pseudoarciger* n. sp. und verwandte Arten der *Lepthyphantes pallidus*-Gruppe. - Senckenbergiana biologica 66: 115–118.
- Wunderlich, J. 1986. Spinnenfauna gestern und heute - Fossile Spinnen in Bernstein und ihre heute lebenden Verwandten. — Erich Bauer Verlag bei Quelle & Meyer, Wiesbaden, 283 pp.

- Zabka, M. 1997. Salticidae – Pajaki skaczace (Arachnida: Araneae) (Salticidae of Poland). — *Fauna Polski (Fauna Poloniae)* 19: 1–189.
- Zingerle, V. 1999a. Epigäische Spinnen und Weberknechte im Naturpark Sextner Dolomiten und am Sellajoch (Südtirol, Italien (Araneae, Opiliones). — *Berichte des Naturwissenschaftlich-medizinischen Vereins in Innsbruck* 86: 165–200.
- Zingerle, V. 1999b. Spider and harvestman communities along a glaciation transect in the Italian Dolomites. — *Journal of Arachnology* 27: 222–228.

(received September 14, 2005; accepted June 1, 2006)

Annex:

Tab. 1: List of all species found from June 2002 to May 2004 on Alp Flix (Grisons, Switzerland) with a variety of methods. Abbreviations: BRE: branch eclector; BEA: beating; RCF: restricted canopy fogging; FUN: funnel trap; PTF: pitfall traps near the funnel traps; PTB: pitfall traps under trees with branch eclectors; FRS: frame samples; LIT: litter samples; SWE: sweep nettings; HST: hand samples at the timberline; HSS: hand samples at nearby sites.

taxa	methods on branches			methods on the ground						hand sampling		m/f	total
	BRE	BEA	RCF	FUN	PTF	PTB	FRS	LIT	SWE	HST	HSS		
Amaurobiidae													
<i>Coelotes terrestris</i> (Wider, 1834)						1/0						1/0	1
Araneidae													
<i>Aculepeira ceropegia</i> (Walckenaer, 1802)										0/2	0/1	0/3	3
<i>Araneus diadematus</i> Clerck, 1757										0/2		0/2	2
<i>Araneus quadratus</i> Clerck, 1757										1/6	1/0	2/6	8
<i>Araniella displicata</i> (Hentz, 1847)			0/2							1/0		1/2	3
<i>Parazygiella montana</i> (C.L. Koch, 1834)	0/1										0/1	0/2	2
Dictynidae													
<i>Dictyna arundinacea</i> (Linnaeus, 1758)		3/3	0/5							2/5		5/13	18
<i>Mastigusa arietina</i> (Thorell, 1871)						0/3						0/3	3
Gnaphosidae													
<i>Drassodes cupreus</i> (Blackwall, 1834)				3/0		6/0						9/0	9
<i>Drassodes pubescens</i> (Thorell, 1856)				3/0		7/4						10/4	14
<i>Gnaphosa leporina</i> (L. Koch, 1866)				1/0	1/1	7/3						9/4	13
<i>Haplodrassus signifer</i> (C.L. Koch, 1839)				16/11	4/0	29/29						49/40	89
<i>Micaria aenea</i> Thorell, 1871				12/5	10/1	36/14						58/20	78
<i>Micaria pulicaria</i> (Sundevall, 1831)						0/2				1/0		1/2	3
<i>Zelotes talpinus</i> (L. Koch, 1872)				1/0		1/0				0/1		2/1	3
Hahniidae													
<i>Cryphoea silvicola</i> (C.L. Koch, 1834)	15/7	0/4	0/1			141/59	0/2	0/2	1/1			157/76	233
Linyphiidae													
<i>Agyphantes expunctus</i> (O. P.-Cambridge, 1875)	55/12	4/72	5/37			28/60			2/2	2/13	0/2	96/198	294
<i>Agyreta cauta</i> (O. P.-Cambridge, 1902)				4/0	2/0	32/11						38/11	49
<i>Anguliphantes monticola</i> (Kulczyński, 1881)						8/1						8/1	9
<i>Bolephthyphantes index</i> (Thorell, 1856)						2/5						2/5	7
<i>Bolyphantes alticeps</i> (Sundevall, 1833)				1/1	4/0	7/18			3/1			15/20	35
<i>Bolyphantes luteolus</i> (Blackwall, 1833)	0/1			5/3	11/4	113/89	0/1		1/5			130/103	233
<i>Caracladus avicula</i> (L. Koch, 1869)	0/1			11/10	5/8	251/219	10/34	0/7	0/1			277/280	557

taxa	methods on branches			methods on the ground						hand sampling		m/f	total
	BRE	BEA	RCF	FUN	PTF	PTB	FRS	LIT	SWE	HST	HSS		
<i>Centromerus arcanus</i> (O. P.-Cambridge, 1873)						3/0						3/0	3
<i>Centromerus pabulator</i> (O. P.-Cambridge, 1875)				4/0	11/0	75/31	1/4					91/35	126
<i>Ceratinella brevis</i> (Wider, 1834)						1/0						1/0	1
<i>Erigone atra</i> Blackwall, 1833											1/0	1/0	1
<i>Erigonella subelevata</i> (L. Koch, 1869)						9/6						9/6	15
<i>Evansia merens</i> O. P.-Cambridge, 1900						0/2						0/2	2
<i>Gonatium rubens</i> (Blackwall, 1833)				0/1		0/8	0/1					0/10	10
<i>Improphantes nitidus</i> (Thorell, 1875)						136/122	0/1	0/1				136/124	260
<i>Lepthyphantes nodifer</i> Simon, 1884						0/1						0/1	1
<i>Macrargus carpenteri</i> (O. P.-Cambridge, 1894)				1/0	5/0	98/12						104/12	116
<i>Mansuphantes pseudoarciger</i> (Wunderlich, 1985)						24/4						24/4	28
<i>Maro lehtineni</i> Saaristo, 1971				1/0		7/1						8/1	9
<i>Meioneta innotabilis</i> (O. P.-Cambridge, 1863)	1/0											1/0	1
<i>Meioneta rurestris</i> (C.L. Koch, 1836)				0/1								0/1	1
<i>Metopobactrus schenkeli</i> Thaler, 1976				2/0		1/0						3/0	3
<i>Micrargus alpinus</i> Relys & Weiss, 1997						3/2						3/2	5
<i>Minicia marginella</i> (Wider, 1834)							0/1					0/1	1
<i>Moebelia penicillata</i> (Westring, 1851)	0/1											0/1	1
<i>Mughiphantes cornutus</i> (Schenkel, 1927)						26/25						26/25	51
<i>Mughiphantes mughi</i> (Fickert, 1875)	3/1	0/3	1/3			5/4			3/2	7/8		19/21	40
<i>Obscuriphantes obscurus</i> (Blackwall, 1841)	1/0					1/0			0/1			2/1	3
<i>Panamomops tauricornis</i> (Simon, 1881)						46/37	4/34	6/22				56/93	149
<i>Pelecopsis elongata</i> (Wider, 1834)				0/1		208/150	2/5	3/4				213/160	373
<i>Pelecopsis radiculicola</i> (L. Koch, 1872)				1/3	1/2	6/1	6/9					14/15	29
<i>Pityohyphantes phrygianus</i> (C.L. Koch, 1836)	0/2	0/10	0/4			2/1				1/13	0/1	3/31	34
<i>Porrhomma cf. campbelli</i> F.O. P.-Cambridge, 1894						0/4						0/4	4
<i>Porrhomma pallidum</i> Jackson, 1913						22/15						22/15	37
<i>Scotargus pilosus</i> Simon, 1913				1/2		5/2						6/4	10
<i>Scotinotylus alpigena</i> (L. Koch, 1869)						536/92	8/19	1/1	0/1			545/113	658
<i>Scotinotylus clavatus</i> (Schenkel, 1927)						69/20	1/20	0/3				70/43	113

taxa	methods on branches			methods on the ground						hand sampling		m/f	total	
	BRE	BEA	RCF	FUN	PTF	PTB	FRS	LIT	SWE	HST	HSS			
<i>Pityohyphantes phrygianus</i> (C.L. Koch, 1836)	0/2	0/10	0/4			2/1					1/13	0/1	3/31	34
<i>Porrhomma cf. campbelli</i> F.O. P.-Cambridge, 1894						0/4							0/4	4
<i>Porrhomma pallidum</i> Jackson, 1913						22/15							22/15	37
<i>Scotargus pilosus</i> Simon, 1913				1/2		5/2							6/4	10
<i>Scotinotylus alpigena</i> (L. Koch, 1869)						536/92	8/19	1/1	0/1				545/113	658
<i>Scotinotylus clavatus</i> (Schenkel, 1927)						69/20	1/20	0/3					70/43	113
<i>Stemonyphantes conspersus</i> (L. Koch, 1879)	14/2	2/3	0/2	1/0		8/5							25/12	37
<i>Tapinocyba affinis</i> Lessert, 1907				1/0	0/1	177/46	30/116	2/6					210/169	379
<i>Tenuiphantes cristatus</i> (Menge, 1866)						2/5							2/5	7
<i>Tenuiphantes jacksonoides</i> (van Helsdingen, 1977)						1/0							1/0	1
<i>Tenuiphantes menzei</i> (Kulczyński, 1887)				1/6	2/0	18/15			0/1				21/22	43
<i>Tenuiphantes tenebricola</i> (Wider, 1834)						4/2			1/0				5/2	7
<i>Thyreostenius biovatus</i> (O. P.-Cambridge, 1875)				0/1		0/9							0/10	10
<i>Walckenaeria antica</i> (Wider, 1834)				1/0		15/2							16/2	18
<i>Walckenaeria languida</i> (Simon, 1914)						0/1							0/1	1
<i>Walckenaeria mitrata</i> (Menge, 1868)						0/1							0/1	1
<i>Walckenaeria monoceros</i> (Wider, 1834)						1/0							1/0	1
Liocranidae														
<i>Agroeca proxima</i> (O. P.-Cambridge, 1871)				2/0		4/0							6/0	6
Lycosidae														
<i>Alopecosa accentuata</i> (Latreille, 1817)				2/0	1/0								3/0	3
<i>Alopecosa pulverulenta</i> (Clerck, 1757)				17/4	9/2	40/16				2/0			68/22	90
<i>Alopecosa taeniata</i> (C.L. Koch, 1835)				119/17	24/6	303/43		0/1	0/1	3/2			449/70	519
<i>Arctosa renidescens</i> Buchar and Thaler, 1995				11/1	14/4	28/17							53/22	75
<i>Pardosa blanda</i> (C.L. Koch, 1833)				49/9	7/1	41/27				2/9	1/1		100/47	147
<i>Pardosa ferruginea</i> (L. Koch, 1870)				0/1									0/1	1
<i>Pardosa mixta</i> (Kulczyński, 1887)										3/19	0/1		3/20	23
<i>Pardosa riparia</i> (C.L. Koch, 1833)				257/95	104/35	1835/510				36/70	2/1		2234/711	2945
<i>Pirata piraticus</i> (Clerck, 1757)												0/1	0/1	1
<i>Trochosa terricola</i> Thorell, 1856						1/0							1/0	1

taxa	methods on branches			methods on the ground						hand sampling		m/f	total
	BRE	BEA	RCF	FUN	PTF	PTB	FRS	LIT	SWE	HST	HSS		
Philodromidae													
<i>Philodromus cespitum</i> (Walckenaer, 1802)	4/0	0/2										4/2	6
<i>Philodromus collinus</i> C.L. Koch, 1835						1/0						1/0	1
<i>Philodromus vagulus</i> Simon, 1875	2/2					0/1					0/1	2/4	6
<i>Thanatus formicinus</i> (Clerck, 1757)				0/1	2/0	18/1				0/1		20/3	23
Salticidae													
<i>Evarcha arcuata</i> (Clerck, 1757)											1/0	1/0	1
<i>Salticus scenius</i> (Clerck, 1757)											2/2	2/2	4
<i>Talavera monticola</i> (Kulczyński, 1884)							0/1					0/1	1
Sparassidae													
<i>Micrommata virescens</i> (Clerck, 1757)	2/0		0/1			1/1					1/0	4/2	6
Theridiidae													
<i>Achaearanea ohlerti</i> (Thorell, 1870)			0/2							0/2		0/4	4
<i>Robertus truncorum</i> (L. Koch, 1872)				1/0	1/0	20/12	0/10	1/0				23/22	45
<i>Steatoda phalerata</i> (Panzer, 1801)						1/0						1/0	1
<i>Theridion impressum</i> L. Koch, 1881		0/1	3/1							3/17	0/2	6/21	27
Thomisidae													
<i>Ozyptila atomaria</i> (Panzer, 1801)				1/0								1/0	1
<i>Xysticus audax</i> (Schrank, 1803)	3/1	1/0		2/1	1/0	12/3			0/1	0/1		19/7	26
<i>Xysticus bifasciatus</i> C.L. Koch, 1837											1/0	1/0	1
<i>Xysticus gallicus</i> Simon, 1875						1/0					0/1	1/1	2
<i>Xysticus luctuosus</i> (Blackwall, 1836)				5/0		8/0						13/0	13
<i>Xysticus macedonicus</i> äïlhavy, 1944			0/1									0/1	1
number of individuals	m/f	100/31	10/100	9/57	537/174	219/65	4492/1774	62/258	13/47	11/17	64/171	10/15	5527/2709
total		131	110	66	711	284	6266	320	60	28	235	25	8236
number of species		15	10	10	37	21	69	15	10	12	19	17	93
number of exclusive species per method		2	0	1	3	0	22	2	0	0	1	3	34