

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 81 (2008)

Heft: 3-4

Artikel: Revision of the Nearctic *Drosophila pinicola* species group (Diptera,
Drosophilidae)

Autor: Bächli, Gerhard / Vilela, Carlos R.

DOI: <https://doi.org/10.5169/seals-402973>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Revision of the Nearctic *Drosophila pinicola* species group (Diptera, Drosophilidae)

GERHARD BÄCHLI¹ & CARLOS R. VILELA²

¹ Zoologisches Museum, Universität Zürich, Winterthurerstrasse 190, CH-8057, Zürich, Switzerland.
E-mail: baechli@zm.uzh.ch

² Departamento de Genética e Biologia Evolutiva, Instituto de Biociências, Universidade de São Paulo, Caixa Postal 11461, São Paulo - SP, 05422-970, Brazil. E-mail: crvilela@ib.usp.br

Drosophila flavopinicola Wheeler, 1954 and *D. pinicola* Sturtevant, 1942, belonging to the small Nearctic *pinicola* group, are redescribed based both on types, whenever they were available, and ordinary specimens deposited in The American Museum of Natural History (New York), The National Museum of Natural History (Washington, D.C.), and in the Zoologisches Museum der Universität Zürich. *Drosophila luiserrai* Vilela and Bächli, 2002, a species previously not assigned to any group, is considered a junior synonym of *D. flavopinicola* Wheeler, 1954. Photomicrographs and line drawings of male terminalia are also provided.

Keywords. Western USA, Southwestern Canada, fleshy fungi, male terminalia, new synonym, *Sequoiadendron giganteum*.

INTRODUCTION

In September 2001, we had the opportunity to analyze two conspecific male specimens of a *Drosophila* species from the northwestern United States. They had been sent to the senior author by Prof. Luis Serra, from the Universidad de Barcelona, Spain, who made the collection at Bellingham, Washington, USA, in September 1995. Our attempt to identify the flies using the key to North American species of *Drosophila* published by Patterson (1943) was completely unsuccessful. Being unable to ascribe the two male specimens to any of the described species included in that key, and believing they belonged to a species new to science we decided to describe them as *Drosophila luiserrai* Vilela and Bächli, 2002, honoring the collector.

Early in 2004, while revising the literature regarding published drawings of the male terminalia of species of *Drosophila* for one of the last versions of a manuscript of a book on the Scandinavian Drosophilidae (Bächli *et al.* 2004), we realized that we had overlooked a paper by Malogolowkin (1953) who had depicted the male terminalia somewhat similar to those we had published for *Drosophila luiserrai*. According to the figure caption, Malogolowkin's (1953: 246, figs 1 and 2) drawings depicted *Drosophila pinicola* Sturtevant, 1942, intriguingly a species included in Patterson's key (*op. cit.*) we had previously checked.

Then, we surveyed the literature regarding keys to and descriptions of the species of *Drosophila* included in the *pinicola* group. Surprisingly we found a key to the *Drosophila* species of the United States in another overlooked publication aimed at teaching practical experiments in Genetics with *Drosophila* (Strickberger 1962:

117), and therefore not expected to have taxonomy information. In the couplet of the *pinicola* group, the two males we described as *Drosophila luisserrai* run to *Drosophila flavopinicola* Wheeler, 1954, a species obviously absent from Patterson's 1943 key as it was described later.

The only two species of *Drosophila* currently ascribed to the *pinicola* group, namely *D. pinicola* and *D. flavopinicola*, occur allopatrically (Spieth & Heed 1975) in the extreme western United States; the latter species is also found in southwestern Canada (Wheeler 1981a). The first one, as implied by its epithet *pinicola* (Latin *pinus* = pine + *cola* = dweller), lives in close association with conifers and is probably a sap-feeder (Sturtevant 1942: 42), as it has been observed on moist spots on the trunks of *Sequoia gigantea* [currently *Sequoiadendron giganteum* (Lindl.)]. Additionally, according to a personal communication of Theodosius Dobzhansky to Spieth & Heed (1975: 288), it has been reared from field-collected mushrooms. As originally suspected by Wheeler (1954: 48), *D. flavopinicola* has also been bred from several fungal species (Spieth & Heed 1975: 288), in particular those of the genus *Amanita*.

The present revision was designed to clarify both the identities of the two species of *Drosophila* currently ascribed to the *pinicola* group and their relation with *Drosophila luisserrai*.

MATERIAL AND METHODS

The 55 specimens analyzed in the present revision are housed in the following institutions: American Museum of Natural History, New York (AMNH), National Museum of Natural History, Washington, D.C. (NMNH), Zoologisches Museum der Universität Zürich, Zurich (ZMUZ).

Label data attached to each specimen are cited in full with a backslash indicating a label change. Our own notes or interpretations are included in brackets (also in other items throughout the text). Where appropriate, one or two of the following additional labels were added to every analyzed specimen: «*Drosophila pinicola* Sturtevant, 1942 Bächli & Vilela det. 2007», «*Drosophila flavopinicola* Wheeler, 1954 Bächli & Vilela det. 2007», «terminalia illustrated and photographed». Only a sample of the examined material was used for measurements.

Preparations of microscope slides were made following Wheeler & Kambyzellis (1966) and Kaneshiro (1969). The abdominal sclerites, including the disarticulated terminalia, are preserved in microvials filled with glycerin and attached by the stopper to the pin of the respective specimen. Refer to Bächli *et al.* (2004) for further details.

Male terminalia were drawn using a camera lucida (1.8x) attachment on a compound microscope under a 20x or 25x or 40x objective. They were all photographed with a photomicroscope under a 6.3x objective.

Whenever in the same plate, all figures were drawn to the same scale and all photomicrographs were taken and enlarged to the same magnification, unless otherwise indicated.

For measurements and indices see Vilela & Bächli (1990), for morphological terminology see Vilela & Bächli (2000) and Bächli *et al.* (2004).

pinicola




Fig. 1. *Drosophila pinicola* Sturtevant from Santa Anita Canyon, California, USA, July 1949 (stock), external male terminalia. A, epandrium, cerci, surstyli, and decasternum, oblique posterior view. B, surstyli and decasternum, posterior view.

TAXONOMY

***Drosophila pinicola* species group**

pinicola species group Sturtevant 1942: 30 [diagnosis]; Freire-Maia & Pavan 1950: 22 [Portuguese diagnosis]; Patterson & Stone 1952: 22 [diagnosis, ecology, distribution]; Okada 1958: 267 [description]; Hardy 1965: 26 [ecology]; Spieth & Heed 1975: 287 [taxa included]; Throckmorton 1975: 449, 461 [phylogeny]; Hardy & Kaneshiro 1981: 333 [phylogeny]; Carson & Yoon 1982: 330 [phylogeny]; Beverley & Wilson 1984: 9 [phylogeny]; Stacey et al. 1986: 526 [phylogeny]; Ashburner et al. 2005: 1129, 1194 [phylogeny].

Diagnosis. Thorax ground color yellowish or brownish; scutum with a faint median stripe; acrostichals in six rows; anterior scutellars parallel; both or only the posterior crossveins clouded; inner subventral region of cerci conspicuously membranous; surstyli longer than wide, dorsally membranous and pleated; hypandrium wider than long.

Taxa included (2). *Drosophila flavopinicola* Wheeler, 1954; *D. pinicola* Sturtevant, 1942.

Distribution. According to Spieth & Heed (1975: 288), both species occur allopatrically in the extreme western USA, and *D. flavopinicola* also reaches southwestern Canada. However, based on the specimens' labels, we found out that at least in one of the collection sites located in the state of Washington (10 mi N. of Raymond) both species were collected together in the same patch of forest. Refer to the list of material examined under each binomial for further data.

Ecology. Both species live in cool and moist areas, are fungivorous during their larval stage, and females need temperatures below 18 °C to produce mature eggs (Spieth & Heed 1975: 290, 292). Whether the adults of both species are obligatory sap-feeders remains an open question.

Comments. Although Spieth & Heed (1975: 290) stated that the male genitalia of both species are essentially identical except for size, we figured out that, in addition to the differences in adult body size and color (*D. flavopinicola* is larger and the wings are partially brownish; *D. pinicola* is smaller and the wings are generally hyaline), the detailed structures of some of the male terminalia sclerites, such as surstyli, decasternum, gonopods, and aedeagus, are the most reliable way to tell them apart.

***Drosophila flavopinicola* Wheeler, 1954**

(Figs 2D, 4B, E–G, J)

Drosophila flavopinicola Wheeler, 1954: 47, 48 [description]. Clayton & Ward 1954: 99, 101 [karyotypes]; Wheeler 1959: 190 [catalog]; Wheeler 1960: 136 [supplementary description]; Strickberger 1962: 117 [key]; Wheeler 1965: 764 [Nearctic catalog]; Spieth 1974: 146 [ecology]; Clayton & Wheeler 1975: 481 [karyotypes]; Spieth & Heed 1975: 288 [distribution, breeding sites], 289–294 [feeding sites and behavior, life cycle, sexual behavior, relationships]; Kaneshiro *et al.* 1977: 85 [ecology]; Lakovaara *et al.* 1977: 323 [polymorphism]; Hardy & Kaneshiro 1981: 333 [phylogeny], Wheeler 1981a: 41 [World catalog]; Wheeler 1981b: 116 [distribution]; Carson & Yoon 1982: 330 [phylogeny]; Shorrocks 1982: 409 [ecology]; Beverley & Wilson 1982: 256, 262 [phylogeny]; Courtney *et al.* 1990: 241 [ecology]; Pascual *et al.* 1993: 219, 225 [distribution]; Brake & Bächli 2008: 55 [World catalogue].

Drosophila luisserrai Vilela and Bächli, 2002: 211 [description]; Brake & Bächli 2008: 65 [World catalogue]. **New Synonymy.**




Fig. 2. Male terminalia of species of *Drosophila* of the *pinicola* group: A–C, *D. pinicola*, two specimens (B and C, same specimen) from Santa Anita Canyon, California, USA, July 1949 (stock). D, *D. flavopinicola*, holotype. A and B, hypandrium and gonopods, ventral and left lateral view respectively. C and D, decasternum, dorsal view. The letters «a» indicate the membranous articulations between decasternum and surstyli.

Diagnosis. Generally yellowish flies; scutum with a faint median stripe; pleura brownish-yellow, with diffuse darkenings along the upper margin and through kat-episternum; tergites with diffuse, usually medially interrupted brown marginal bands; wing with brown shadows along both main crossveins and a distinctly brownish costal margin; surstylus with ca. 10 cone-shaped prensisetae; anterior region of decasternum with a large, open V-shaped membranous area; posterior inner margin


pinicola

Fig. 3. *Drosophila pinicola* Sturtevant from Santa Anita Canyon, California, USA, July 1949 (two specimens [A–C, one; D, E, another] from the same stock). A–E, aedeagus+aedeagal apodeme and paraphyses, several aspects from dorsal through ventral.

of gonopod not projected backwards; aedeagus pointed at tip in dorsal view, bearing a pair of short dorsal expansions projected frontwards; paraphysis microtrichose in the inner wall.

Material examined (10 ♂♂, 16 ♀♀). Holotype ♂ (dissected, deposited in the NMNH), labelled: «Prarie [sic] Creek S.P. [Prairie Creek Redwoods State Park],

Calif. July 1951 \ MR Wheeler WBHeed \ 2179.4 yellow pinicola \ HOLOTYPE [red label] \ *Drosophila flavopinicola* Wheeler». Paratypes (6 ♂♂, 14 ♀♀, AMNH); 6 ♀♀: same labels as holotype [except the last two; with yellow paratype labels]; 3 ♂♂, 6 ♀♀: «Rogue River NE Gold Beach Oregon \ MR Wheeler HB Heed June [striketrough] 1951 JULY \ PARATYPE [yellow label]»; 3 ♂♂, 1 ♀: «Olympic Natl. For. nr Sequim, Wash. Aug 3, 1951 \ PARATYPE [yellow label]»; 1 ♀: «WASHINGTON: 10 mi. N. Raymond 2186.5 VII/31–VIII/1/51 W.B.Heed/M.R.Wheeler \ PARATYPE [yellow label]». Five ordinary specimens deposited in the AMNH (3 ♂♂, 2 ♀♀); 1 ♂, 2 ♀♀: «Carmel Calif. \ A. Sokoloff July 1958»; 1 ♂: «OREGON: Lake Creek, 40 mi. W. N.W. Eugene VII29–VII30\51 2184.10 W.B. Heed/M.R. Wheeler»; 1 ♂, dissected: «Prairie [sic] Creek S.P. [Prairie Creek Redwoods State Park], Calif. July 1951 \ MR Wheeler WBHeed \ 2179.4 flavicola [sic, manuscript name]».

Type locality. Prairie Creek Redwood State Park, near Crescent City, California, USA.

Redescription (n = 7). ♂.

Head. Frons brownish-yellow, darker brownish towards vertex, frontal length 0.38 (0.32–0.42) mm; frontal index = 1.00 (0.96–1.09), top to bottom width ratio = 1.32 (1.21–1.41). Frontal triangle blackish-brown, grayish microtrichose, laterally slightly convex, about 58–76 % of frontal length; ocellar triangle blackish-brown, prominent, about 40–56 % of frontal length. Orbital plates blackish-brown, grayish microtrichose, apically diverging from eye margin, about 72–87 % of frontal length. Orbital setae black, almost in a line, distance of or3 to or1 = 86–112 % of or3 to vtm, or1 / or3 ratio = 0.89 (0.79–1.07), or2 / or1 ratio = 0.53 (0.40–0.64), postvertical setae = 60 (47–72) %, ocellar setae = 68 (50–82) % of frontal length; vibrissal index = 0.40 (0.38–0.43). Face brownish-yellow. Carina nose-like, high, longitudinally slightly carved. Cheek pale yellow, narrowly brownish along the lower genal margin, index about 6 (4–9). Eye with short, dense pile, main axis oblique, index = 1.14 (1.00–1.25). Antennae brownish, length to width ratio of flagellomere 1 = 1.45 (1.29–1.60). Arista with 3 (–4) dorsal, (1–) 2 ventral and about 6 small inner branches, plus terminal fork. Proboscis and palpus yellowish.

Thorax length 1.30 (1.04–1.53) mm. Scutum yellowish, centrally slightly darker, forming a diffuse brownish stripe which is darker towards scutellum, 6–8 rows of acrostichal setae. h index = 1.72 (1.50–1.91). Transverse distance of dorsocentral setae 153–215 % of longitudinal distance; dc index = 0.67 (0.59–0.76). Scutellum basal-centrally darker brownish, laterally yellowish, distance between apical scutellar setae about 71–82 % of that between apical and basal one, scut index = 1.03 (0.72–1.14). Pleura yellowish, slightly darkened along upper margin, forming a stripe in some specimens, also katapisternum somewhat darkened, sterno index = 0.54 (0.48–0.59), median katapisternal seta about 41 (38–44) % of the anterior one. Halter yellow. Legs brownish-yellow, preapical seta on tibia 3, apical seta on tibia 2.

Wing distinctly darkened along costal margin, reaching close to the tip of R_{4+5} , veins R_{4+5} and M apically slightly convergent, in some specimens with darkened tips, both main crossveins more or less darkened, length 3.22 (2.45–3.68) mm, length to width ratio = 2.22 (2.09–2.33). Indices: C = 3.64 (3.24–4.05), ac = 2.48 (2.11–3.00), hb = 0.41 (0.33–0.44), 4C = 0.67 (0.59–0.75), 4v = 1.58 (1.44–1.67), 5x = 1.34 (1.20–1.44), M = 0.49 (0.40–0.55), prox. x = 0.53 (0.46–0.62).

Abdomen predominantly yellowish, with more or less diffuse brownish marginal bands which are medially interrupted on tergites 2 and 3, broadened, reaching almost the anterior margins on tergites 4 to 6; ventral margins of all tergites dark.

♂ terminalia (n = 2) (Figs 2D, 4B, E–G, J). See Vilela & Bächli (2002) for additional detailed illustrations (p. 212 [fig. 1]) and description (p. 213) of the male terminalia sclerites of its junior synonym *Drosophila luisserrai*.

♀ (n = 5)

Measurements: Frontal length 0.42 (0.39–0.44) mm; frontal index = 0.89 (0.85–0.93), top to bottom width ratio = 1.27 (1.04–1.37). Frontal triangle about 64–72 % of frontal length; ocellar triangle about 40–50 % of frontal length. Orbital plates about 74–85 % of frontal length. Distance of or3 to or1 = 67–90 % of or3 to vtm, or1 / or3 ratio = 0.82 (0.76–0.88), or2 / or1 ratio = 0.65 (0.50–0.73), post-vertical setae = 72 (64–76) %, ocellar setae = 84 (76–96) % of frontal length; vibrissal index = 0.42 (0.37–0.50). Cheek index about 5 (3–7). Eye index = 1.16 (1.10–1.24). Thorax length 1.57 (1.49–1.62) mm. h index = 1.81 (1.62–2.10). Transverse distance of dorsocentral setae 167–208 % of longitudinal distance; dc index = 0.65 (0.64–0.67). Distance between apical scutellar setae about 67–81 % of that between apical and basal one; scut index = 1.09 (1.03–1.23), sterno index = 0.57 (0.50–0.68), median katapisternal seta about 31 (29–33) % of anterior one. Wing length 3.76 (3.60–3.92) mm, length to width ratio = 2.32 (2.20–2.49). Indices: C = 3.78 (3.41–4.11), ac = 2.14 (1.90–2.33), hb = 0.46 (0.42–0.50), 4C = 0.65 (0.59–0.69), 4v = 1.55 (1.35–1.75), 5x = 1.14 (1.00–1.27), M = 0.42 (0.35–0.50), prox. x = 0.55 (0.50–0.61).

Distribution. Extreme western USA (Washington, Oregon and California) and southwestern Canada (British Columbia) (Wheeler 1965: 764).

Ecology. Fungivorous at larval stage. It has been reared from several fleshy fungi, especially of the genus *Amanita* (Spieth & Heed 1975: 288). Adult feeding site unknown.

Comments. In the original description, Wheeler (1954: 48) stated that 23 specimens were selected as paratypes. However we have received only 19 specimens labeled as such from the AMNH. We considered the last specimen cited in the item «material examined» as an aberrant individual of *Drosophila flavopinicola* as regards the internal male terminalia (aedeagus much shorter, although *D. flavopinicola*-like, Fig. 4E). It bears the same labels as the type series specimens, except for the manuscript name *flavicola* and the absence of a paratype label. We suppose it has been excluded from the type series by Wheeler because of its being an aberrant specimen. It was received from the AMNH loan together with the specimens identified as *Drosophila pinicola*, in spite of being abnormal as regards the body color (remarkably lighter).

***Drosophila pinicola* Sturtevant, 1942**

(Figs 1, 2A–C, 3, 4A, C, D, H, I)

Drosophila pinicola Sturtevant, 1942: 40–42 [description]. Patterson 1943: 49 [key], 97–98 [redescription], plate V [female imago illustration]; Patterson & Wagner 1943: 224–226 [distribution]; Dobzhansky 1948: 160 [distribution]; Patterson & Wheeler 1949: 225 [catalog]; Carson 1951: 92 [distribution, ecology]; da Cunha *et al.* 1951: 99, 100 [ecology]; Patterson & Stone 1952: 83–85, 437, 504 [phylogeny]; de Castro 1953: 365 [description], 368 [phylogeny]; Malogolowkin 1953: 246 [male terminalia]; Spiess 1953: 113 [distribution]; Carson *et al.* 1956: 541




Fig. 4. Photomicrographs of male terminalia of *Drosophila* species of the *pinicola* group. *D. pinicola* Sturtevant : A, C, D, H, and I (two specimens [A, C, and I, one specimen; D and H, another specimen] from the same stock from Santa Anita Canyon, California, USA, July 1949). *D. flavopinicola* Wheeler: B, E–G, and J (B, F, and G, holotype; E, aberrant specimen from Prairie Creek Redwoods State Park, California, USA, July 1951; J, paratype of its junior synonym *Drosophila luisserrai* Vilela & Bächli). A, and B, posterior view; C–E, and G, dorsal view; F, H–J, left lateral view.

[distribution, ecology]; Cooper & Dobzhansky 1956: 528, 529 [distribution], 532 [sex ratio]; Dobzhansky *et al.* 1956: 545–549 [distribution, ecology]; Phaff *et al.* 1956: 535–538 [distribution, ecology]; Wheeler 1959: 195 [catalog]; Miller *et al.* 1962: 2, 4 [distribution, ecology]; Strickberger 1962: 117 [key]; Dobzhansky, 1963: 336 [distribution]; Sokoloff 1964: 207, 212 [distribution, ecology]; Wheeler 1965: 765 [Nearctic catalog]; Strickberger & Wills 1966: 595 [distribution]; Takada 1966: 45 [male external terminalia]; Cole 1970: 410 [distribution]; Anderson *et al.* 1972: 101 [distribution]; Spieth 1974: 146 [ecology]; Clayton & Wheeler 1975: 481 [karyotype]; Spieth & Heed 1975: 288 [distribution, breeding sites], 289–294 [feeding sites, behavior, life cycle, sexual behavior, relationship]; Kaneshiro *et al.* 1977: 85 [ecology]; Lakovaara *et al.* 1977: 323 [polymorphism]; Moore *et al.* 1979: 163 [distribution]; Taylor & Condra 1979: 299 [distribution]; Marks *et al.* 1980: 142 [distribution]; Starmer 1981: 49 [ecology]; Wheeler 1981a: 47 [World catalog]; Wheeler 1981b: 116 [distribution]; Beverley & Wilson 1982: 256, 262 [phylogeny]; Bryant *et al.* 1982: 28 [distribution]; Shorrocks 1982: 401, 409 [ecology]; Klaczko *et al.* 1983: 412 [distribution, ecology]; Moore & Moore 1984: 739, 740 [distribution]; Beckenbach & Prevosti 1986: 14 [distribution]; Zelentsova *et al.* 1986: 474 [phylogeny]; Prevosti *et al.* 1987: 8, 14 [distribution]; Spieth 1987: 249, 252 [distribution, ecology]; Prevosti *et al.* 1989: 116, 117 [distribution]; Courtney *et al.* 1990: 241, 250, 262 [ecology]; DeSalle 1992: 914 [phylogeny]; Pascual *et al.* 1993: 220, 221, 225 [distribution]; Lachance *et al.* 1995: 485, 486, 489–493 [distribution, ecology]; DeSalle *et al.* 1997: 471 [phylogeny]; Remsen & DeSalle 1998: 227–230 [phylogeny]; Remsen & O'Grady 2002: 254, 257, 258, 261 [phylogeny]; Brake & Bächli 2008: 80 [World catalogue].

Diagnosis. Generally brownish-yellow flies; scutum with a faint median stripe; pleura yellowish-brown, with a diffuse stripe along the upper margin; tergites with diffuse, medially interrupted brown marginal bands; wing almost hyaline with diffuse brown shadows along the posterior crossveins; surstylus with ca. 13 cone-shaped prensisetae; anterior region of decasternum an open V-shaped, non membranous area; posterior inner margin of gonopod roundish, corrugated and projected backwards; aedeagus roundish at tip in dorsal view, dorsomedially bubble-shaped in lateral view, projected dorsad, and anteriorly wrinkled; paraphysis not microtrichose on the inner wall.

Material examined (10 ♂♂, 19 ♀♀; all ordinary specimens, two males dissected). AMNH: 2 ♀♀: «CALIFORNIA: Six Rivers Nat. Forest, 5 mi. W. Willow Creek VII/26–VII/27/51 2181.12 W.B.Heed/M.R.Wheeler»; 1 ♂: «WASHINGTON: 10 mi. N. Raymond VII/31–VIII/1/51 2186.10 W.B. Heed/M.R. Wheeler»; 1 ♂, 5 ♀♀: «CALIFORNIA: Inyo Nat. Forest, nr. Lone Pine VII/19–VII/20/51 2176.3 W.B. Heed/M.R. Wheeler»; 2 ♂♂ [both dissected], 1 ♀: «CALIFORNIA: Santa Anita Canyon VII/49 [plus Stock and male symbol in one male]»; 1 ♀: «CALIFORNIA: Rio Hondo X/29/49 [female symbol]»; 2 ♂♂, 2 ♀♀: «CALIFORNIA: Monrovia III/27/50»; 1 ♂, 6 ♀♀: «Carmel Calif. \ A. Sokoloff July 1958». Plus four ordinary specimens deposited in the NMNH: 3 ♂♂: «Kern. [sic] Cany. Calif. Apr. 34 \ A.H. Sturtevant Collection, 1970»; 1 ♀: «Keen [Hemet Valley; on the slopes of Mount San Jacinto], Calif. \ A.H. Sturtevant Collection, 1970»; ZMZ: 1 ♀: «Arcadia Cal. 10.4.49 M. Wheeler coll. \ D. pinicola do Frota [belonged to Dr. Frota-Pes-soa]».

Type locality. Slopes of Mt. [Mount] San Jacinto, [San] Andreas Canyon, California, USA.

Redescription (n = 7). ♂.

Head. Frons brownish-yellow, darker brownish towards vertex, frontal length 0.38 (0.34–0.44) mm; frontal index = 1.10 (0.92–1.33), top to bottom width ratio = 1.40 (1.24–1.67). Frontal triangle blackish-brown, grayish microtrichose, laterally almost straight, about 62–75 % of frontal length; ocellar triangle blackish-brown, prominent, about 38–45 % of frontal length. Orbital plates blackish-brown, grayish

microtrichose, apically slightly diverging from eye margin, about 62–87 % of frontal length. Some specimens with a very narrow, grayish margin along the eye. Orbital setae black, almost in a line, distance of or3 to or1 = 75–100 % of or3 to vtm, or1 / or3 ratio = 0.90 (0.75–1.00), or2 / or1 ratio = 0.45 (0.31–0.58), postvertical setae = 54 (45–60) %, ocellar setae = 76 (60–83) % of frontal length; vibrissal index = 0.44 (0.33–0.57). Face brownish-yellow. Carina nose-like, high, longitudinally slightly carved. Cheek yellowish, in some specimens narrowly brownish along the lower genal margin, index about 6 (4–9). Eye index = 1.21 (1.13–1.27). Pedicel yellowish. Flagellomere 1 brownish, length to width ratio = 1.60 (1.40–1.80). Arista with 3 dorsal, (1–) 2 ventral and about 6 small inner branches, plus terminal fork. Proboscis and palpus yellowish.

Thorax length 1.19 (0.98–1.34) mm. Scutum yellowish, in some specimens centrally slightly darker, forming a diffuse brownish stripe which is darker towards scutellum, 6 rows of acrostichal setae. h index = 2.09 (1.73–2.38). Transverse distance of dorsocentral setae 155–192 % of longitudinal distance; dc index = 0.73 (0.62–0.83). Scutellum basal-centrally darker brownish, laterally yellowish, distance between apical scutellar setae about 69–87 % of that between apical and basal one, scut index = 1.05 (0.97–1.13). Pleura yellowish, slightly darkened along upper margin, forming a stripe in some specimens, also katapisternum somewhat darkened, sterno index = 0.46 (0.42–0.48), median katapisternal seta about 54 (44–70) % of the anterior one. Halter yellow. Legs brownish-yellow, preapical seta on tibia 3, apical seta on tibia 2.

Wing almost hyaline, crossvein dM-Cu in some specimens slightly clouded and a diffuse shadow may occur also along the costal margin, veins R_{4+5} and M almost parallel, length 2.83 (2.66–3.01) mm, length to width ratio = 2.29 (2.23–2.38). Indices: C = 3.43 (3.06–3.86), ac = 2.36 (2.00–3.00), hb = 0.41 (0.38–0.44), 4C = 0.68 (0.61–0.85), 4v = 1.48 (1.35–1.85), 5x = 1.09 (1.00–1.25), M = 0.40 (0.36–0.50), prox. x = 0.56 (0.50–0.65).

Abdomen predominantly yellowish, with very diffuse darker marginal bands.

♂ terminalia (Figs 1A, B, 2A–C, 3, 4A, C, D, H, I). Epandrium posteriorly microtrichose, with ca. 7 median, and 2 upper setae; ventral lobe narrow, dorso-posteriorly microtrichose, with ca. 17 setae, partially covering surstylus. Cerci devoid of ventral lobe, mostly microtrichose, linked to epandrium by membranous tissue, conspicuously membranous on inner ventral region, except for the very ventral strip, which is sclerotized and bear a tuft of short setae (Figs 1A, 4A). Surstylus (Figs 1A, B, 4A) dorsally conspicuously membranous and pleated, not microtrichose, with a more or less straight row of 13 peg-like prenisetae, ca. 4 outer setae and 29 long setae. Decasternum as in Figs 1A, C, 2C, 4A. Hypandrium (Figs 2A, B, 4C, D, H, I) about half epandrium length, anterior margin concave and wide; posterior hypandrial process and dorsal arch absent; inner ventral margin of gonopod linked to paraphysis by membranous tissue, bearing one thin seta and a patch of microtrichia near the anterior inner margin; subdistal inner area and posterior inner margin slightly corrugated (Figs 2A, 4C, D). Aedeagus (Figs 3, 4C, D, H, I) short, roundish at tip in dorsal view, dorsomedially bubble-shaped in lateral view, projected dorsad, and anteriorly wrinkled. Aedeagal apodeme rod-shaped, as long as and posteriorly fused to aedeagus (Figs 3, 4B, D, H, I). Ventral rod dorsoventrally flattened, as long as paraphysis' ventral margin. Paraphysis triangle-shaped in profile, bearing a straight row of ca. 6 thin setulae in the inner wall (Figs 3D, E), but

devoid of microtrichia, linked to dorsodistal margin of aedeagal apodeme by membranous tissue (Fig. 3C).

♀ (n = 7)

Measurements: Frontal length 0.41 (0.39–0.42) mm; frontal index = 0.97 (0.88–1.04), top to bottom width ratio = 1.25 (1.19–1.39). Frontal triangle about 64–79 % of frontal length; ocellar triangle about 42–50 % of frontal length. Orbital plates about 79–87 % of frontal length. Distance of or3 to or1 = 78–112 % of or3 to vtm, or1 / or3 ratio = 0.75 (0.71–0.81), or2 / or1 ratio = 0.75 (0.62–0.85), postvertical setae = 53 (50–57) %, ocellar setae = 81 (76–88) % of frontal length; vibrissal index = 0.43 (0.36–0.53). Cheek index about 5 (5–7). Eye index = 1.17 (1.14–1.19). Thorax length 1.32 (1.29–1.36) mm. h index = 2.24 (2.00–2.83). Transverse distance of dorsocentral setae 153–192 % of longitudinal distance; dc index = 0.73 (0.67–0.81). Distance between apical scutellar setae about 59–86 % of that between apical and basal one; scut index = 0.99 (0.93–1.10), sterno index = 0.51 (0.46–0.55), median katepisternal seta about 47 (42–58) % of anterior one. Wing length 2.96 (2.80–3.08) mm, length to width ratio = 2.24 (2.15–2.34). Indices: C = 3.26 (3.17–3.35), ac = 2.32 (2.13–2.57), hb = 0.40 (0.35–0.44), 4C = 0.73 (0.68–0.78), 4v = 1.52 (1.40–1.61), 5x = 1.13 (1.00–1.22), M = 0.43 (0.40–0.44), prox. x = 0.59 (0.48–0.65).

Distribution. Extreme western USA (Washington and California).

Ecology. Bred from field-collected mushrooms (personal communication of T. Dobzhansky to Spieth and Heed, 1975: 288). Adults were observed on moist spots of Sequoia trunks and probably are sap-feeders (Sturtevant, 1942: 42).

Comments. The syntypes of *Drosophila pinicola* are supposed to be housed in the AMNH (Brake & Bächli, 2008: 80), however no specimens labelled as such were found either in the collections of the AMNH or in those of the NMNH. On the other hand, 8 out of 11 specimens of *Drosophila pinicola* we received as a loan from the NMNH are labelled «Keen, Calif. \ A.H. Sturtevant Collection, 1970». Considering that Keen is one of the three localities on the slopes of Mount San Jacinto (type locality), which in turn is part of the San Andreas Canyon (see fig. I of Wright *et al.* 1942: 364), there is a high probability that those 8 specimens from the Sturtevant Collection (now housed at the NMNH) belong to the type series of *Drosophila pinicola*. However, 7 out of those 8 specimens had no abdomen [probably eaten by dermestids, whose feces remained on the cardboard triangles of the double mountings] and their sexes could not be determined. The remaining one is a female in good conditions, and was analyzed and detailed under the item material examined.

ACKNOWLEDGMENTS

We are indebted to Pierre Brauchli for helping with digital image processing, to Dr. Wayne Mathis, National Museum of Natural History, and to Dr. David A. Grimaldi, American Museum of Natural History, for kindly loaning the specimens included in the present study.

REFERENCES

- Anderson, W.W., Dobzhansky, T. & Pavlovsky, O. 1972. A natural population of *Drosophila* transferred to a laboratory environment. — *Heredity* 28: 101–107.
- Ashburner, M., Golic, K.G. & Hawley, R.S. 2005. *Drosophila*. A laboratory handbook. 2. ed.. 1409 pp. — Cold Spring Harbor Laboratory Press.
- Bächli, G., Vilela, C.R., Andersson Escher, S. & Saura, A. 2004. The Drosophilidae (Diptera) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica*, vol. 39, 362 pp. — Brill, Leiden.

- Beckenbach, A.T. & Prevosti, A. 1986. Colonization of North America by the European species, *Drosophila subobscura* and *D. ambigua*. — American Midland Naturalist 115: 10–18.
- Beverley, S.M. & Wilson, A.C. 1982. Molecular Evolution in *Drosophila* and Higher Diptera. I. Micro-Complement Fixation Studies of a Larval Hemolymph Protein. — Journal of molecular Evolution 18: 251–264.
- Beverley, S.M. & Wilson, A.C. 1984. Molecular Evolution in *Drosophila* and the Higher Diptera. II. A Time Scale for Fly Evolution. — Journal of molecular Evolution 21: 1–13.
- Brake, I. & Bächli, G. 2008. Drosophilidae (Diptera). — World Catalogue of Insects 9: 1–412.
- Bryant, S.H., Beckenbach, A.T. & Cobbs, G.A. 1982. «sex-ratio» trait, sex composition, and relative abundance in *Drosophila pseudoobscura*. — Evolution 36: 27–34.
- Carson, H.L. 1951. Breeding sites of *Drosophila pseudoobscura* and *Drosophila persimilis* in the transition zone of the Sierra Nevada. — Evolution 5: 91–96.
- Carson, H.L. & Yoon, J.S. 1982. Genetics and Evolution of Hawaiian *Drosophila*. In: Ashburner, M., Carson, H.L. & Thompson, J.N. (eds), The Genetics and Biology of *Drosophila*, Vol. 3b, pp. 297–344. — Academic Press, London.
- Carson, H.L., Knapp, E.P. & Phaff, H.J. 1956. Studies on the ecology of *Drosophila* in the Yosemite region of California. III. The yeast flora of the natural breeding sites of some species of *Drosophila*. — Ecology 37: 538–544.
- de Castro, L.E. 1953. Estudo comparativo das cerdas cuneiformes das pernas dos drosofilídeos (Diptera). — Revista brasileira de Biologia 13: 363–368.
- Clayton, F.E. & Ward, C.L. 1954. Chromosomal studies of several species of *Drosophilidae*. — The University of Texas Publication 5422: 98–105.
- Clayton, F.E. & Wheeler, M.R. 1975. A catalog of *Drosophila* metaphase chromosome configurations. In: King, R.C. (ed.), Handbook of Genetics, vol. 3, pp. 471–512. — Plenum Press.
- Cole, F.R. 1970. The Flies of Western North America. — University of California Press.
- Cooper, D.M. & Dobzhansky, T. 1956. Studies on the Ecology of *Drosophila* in the Yosemite region of California. I. The occurrence of species of *Drosophila* in different life zones and at different seasons. — Ecology 37: 526–533.
- Courtney, S.P., Kibota, T.T. & Singleton, T.A. 1990. Ecology of Mushroom-feeding Drosophilidae. — Advances in Ecological Research 20: 225–274.
- da Cunha, A.B., Dobzhansky, T. & Sokoloff, A. 1951. On food preferences of sympatric species of *Drosophila*. — Evolution 5: 97–101.
- DeSalle, R. 1992. The Origin and Possible Time of Divergence of the Hawaiian Drosophilidae: Evidence from DNA Sequences. — Molecular Biology and Evolution 9: 905–916.
- DeSalle, R., Brower, A.V.Z., Baker, R. & Remsen, J. 1997. A Hierarchical View of the Hawaiian Drosophilidae (Diptera). — Pacific Science 51: 462–474.
- Dobzhansky, T. 1948. Genetics of natural populations. XVI. Altitudinal and seasonal changes produced by natural selection in certain populations of *Drosophila pseudoobscura* and *Drosophila persimilis*. — Genetics 33: 158–176.
- Dobzhansky, T. 1963. Genetics of natural populations. XXXIII. A progress report on genetic changes in populations of *Drosophila pseudoobscura* and *Drosophila persimilis* in a locality in California. — Evolution 17: 333–339.
- Dobzhansky, T., Cooper, D.M., Phaff, H.J., Knapp, E.P. & Carson, H.L. 1956. Studies on the ecology of *Drosophila* in the Yosemite region of California. IV. Differential attraction of species of *Drosophila* to different species of yeasts. — Ecology 37: 544–550.
- Freire-Maia, N. & Pavan, C. 1950. Introdução ao estudo da Drosófila. — Cultus 1(5): 1–71.
- Hardy, D.E. 1965. Family Drosophilidae. Insects of Hawaii, vol. 12, Diptera: Cyclorrhapha II, Series Schizophora, Section Acalypterae I, 814 pp. — University of Hawaii Press, Honolulu.
- Hardy, D.E. & Kaneshiro, K.Y. 1981. Drosophilidae of Pacific Oceania. In: Ashburner, M., Carson, H.L. & Thompson, J.N. (eds), The Genetics and Biology of *Drosophila*, vol. 3a, pp. 309–347. — Academic Press.
- Kaneshiro, K.Y. 1969. A Study of the Relationships of Hawaiian *Drosophila* Species Based on External Male Genitalia. — University of Texas Publication 6918: 55–70.
- Kaneshiro, K.Y., Ohta, A.T. & Spieth, H.T. 1977. Mushrooms as bait for *Drosophila*. — *Drosophila* Information Service 52: 85.
- Klaczko, L.B., Powell, J.R. & Taylor, C.E. 1983. *Drosophila* baits and yeasts: species attracted. — Oecologia 59: 411–413.
- Lachance, M.-A., Gilbert, D.G. & Starmer, W.T. 1995. Yeast communities associated with *Drosophila* species and related flies in an eastern oak-pine forest: a comparison with western communities. — Journal of Industrial Microbiology 14: 484–494.
- Lakovaara, S., Saura, A. & Lankinen, P. 1977. Evolution at the alpha-GPDH locus in Drosophilidae. — Evolution 31: 319–330.

- Malogolowkin, C. 1953. Sobre a genitália dos Drosophilídeos. IV. A genitália masculina no subgênero «*Drosophila*» (Diptera, Drosophilidae). — Revista brasileira de Biologia 13: 245–264.
- Marks, R.W., Brittnacher, J.G., McDonald, J.F., Prout, T. & Ayala, F.J. 1980. Wineries, *Drosophila*, Alcohol, and Adh. — Oecologia 47: 141–144.
- Miller, M.W., Phaff, H.J. & Snyder, H.E. 1962. On the occurrence of various species of yeast in nature. — Mycopathologia et Mycologia Applicata 16: 1–18.
- Moore, J.A. & Moore, B.C. 1984. The *Drosophila* of southern California II. Isolations of populations in the Death Valley region. — American Naturalist 124: 738–744.
- Moore, J.A., Taylor, C.E. & Moore, B.C. 1979. The *Drosophila* of southern California. I. Colonization after a fire. — Evolution 33: 156–171.
- Okada, T. 1958. Comparative Morphology of the Drosophilid Flies. VIII. Ejaculatory Caeca of the Adult Flies. (in Japanese, with English summary). — Zoological Magazine, Tokyo 67: 264–271.
- Pascual, M., Ayala, F.J., Prevosti, A. & Serra, L. 1993. Colonization of North America by *Drosophila subobscura*. Ecological analysis of three communities of drosophilids in California. — Zeitschrift für zoologische Systematik und Evolutions-Forschung 31: 216–226.
- Patterson, J.T. 1943. The Drosophilidae of the Southwest. — The University of Texas Publication 4313: 7–216.
- Patterson, J.T. & Stone, W.S. 1952. Evolution in the Genus *Drosophila*. 610 pp. — Macmillan, New York.
- Patterson, J.T. & Wagner, R.P. 1943. Geographical distribution of species of the genus *Drosophila* in the United States and Mexico. — The University of Texas Publication 4313: 217–281.
- Patterson, J.T. & Wheeler, M.R. 1949. Catalogue of described species belonging to the genus *Drosophila*, with observations on their geographical distribution. — The University of Texas Publication 4920: 207–233.
- Phaff, H.J., Miller, M.W., Recca, J.A., Shifrine, M. & Mrak, E.M. 1956. Studies on the ecology of *Drosophila* in the Yosemite region of California. II. Yeasts found in the alimentary canal of *Drosophila*. — Ecology 37: 533–538.
- Prevosti, A., Serra, L., Monclus, M., Mestres, F., Latorre, A., Ribo, G. & Aguade, M. 1987. Colonization de America por *Drosophila subobscura*. — Evolucion Biologica, Bogota, 1: 1–24.
- Prevosti, A., Serra, L., Aguade, M., Ribo, G., Mestres, F., Balana, J. & Monclus, M. 1989. Colonization and Establishment of the Palearctic Species *Drosophila subobscura* in North and South America. In: Fontdevila, A. (ed.), Evolutionary Biology of Transient Unstable Populations, pp. 114–129. — Springer.
- Remsen, J. & DeSalle, R. 1998. Character Congruence of Multiple Data Partitions and the Origin of the Hawaiian Drosophilidae. — Molecular Phylogeny and Evolution 9: 225–235.
- Remsen, J. & O'Grady, P. 2002. Phylogeny of Drosophilinae (Diptera: Drosophilidae), with comments on combined analysis and character support. — Molecular Phylogeny and Evolution 24: 249–264.
- Shorrock, B. 1982. The breeding sites of temperate woodland *Drosophila*. In: Ashburner, M., Carson, H.L. & Thompson, J.N. (eds), The Genetics and Biology of *Drosophila*, vol. 3b, pp. 385–428. — Academic Press.
- Sokoloff, A. 1964. Studies on the Ecology of *Drosophila* in the Yosemite Region of California. V. A Preliminary Survey of Species Associated with *D. pseudoobscura* and *D. persimilis* at Slime Fluxes and Banana Traps. — Pan-Pacific Entomologist 40: 203–218.
- Spiess, E.B. 1953. *Drosophila* from Yosemite National Park, California. — *Drosophila* Information Service 27: 113.
- Spieth, H.T. 1974. Rearing method for some fungal feeding *Drosophila* species. — *Drosophila* Information Service 51: 146.
- Spieth, H.T. 1987. The *Drosophila* Fauna of a Native California Forest (Diptera: Drosophilidae). — Pan-Pacific Entomologist 63: 247–255.
- Spieth, H.T. & Heed, W.B. 1975. The *Drosophila pinicola* Species Group (Diptera: Drosophilidae). — Pan-Pacific Entomologist 51: 287–295.
- Stacey, S.N., Lansman, R.A., Brock, H.W. & Grigliatti, T.A. 1986. Distribution and Conservation of Mobile Elements in the Genus *Drosophila*. — Molecular Biology and Evolution 3(6): 522–534.
- Stарmer, W.T. 1981. A comparison of *Drosophila* habitats according to the physiological attributes of the associated yeast communities. — Evolution 35: 38–52.
- Strickberger, M.W. 1962. Key to United States Species of the Genus *Drosophila*. In: Experiments in Genetics with *Drosophila*, pp. 111–123. — Wiley.
- Strickberger, M.W. & Wills, C.J. 1966. Monthly frequency changes of *Drosophila pseudoobscura* third chromosome gene arrangements in a California locality. — Evolution 20: 592–602.

- Sturtevant, A.H. 1942. The classification of the genus *Drosophila*, with descriptions of nine new species. — The University of Texas Publication 4213: 5–51.
- Takada, H. 1966. Differentiation of the External Male Genitalia in the Drosophilidae. (in Japanese, with English summary). — Proceedings of the Kushiro Woman's College 1: 39–50, 131–132.
- Taylor, C.E. & Condra, C. 1979. Competitor diversity and chromosomal variation in *Drosophila pseudoobscura*. In: Grassle, J.F., Patil, G.P., Smith, W.K. & Taillie, C. (eds), Ecological Diversity in Theory and Practice. — Statistical Ecology Series, vol. 6, pp. 297–308.
- Throckmorton, L.H. 1975. The Phylogeny, Ecology, and Geography of *Drosophila*. In: King, R.C. (ed.), Handbook of Genetics, vol. 3, pp. 421–469. — Plenum Press.
- Vilela, C.R. & Bächli, G. 1990. Taxonomic studies on Neotropical species of seven genera of Drosophilidae. — Mitteilungen der Schweizerischen entomologischen Gesellschaft 63(Suppl.): 1–332.
- Vilela, C.R. & Bächli, G. 2000. Morphological and ecological notes on the two species of *Drosophila* belonging to the subgenus *Siphlodora* Patterson & Mainland, 1944 (Diptera, Drosophilidae). — Mitteilungen der Schweizerischen entomologischen Gesellschaft 73(1–2): 23–47.
- Vilela, C.R. & Bächli, G. 2002. Two new North American *Drosophila* species (Diptera: Drosophilidae). — Mitteilungen der Schweizerischen entomologischen Gesellschaft 75: 211–221.
- Wheeler, M.R. 1954. Taxonomic studies on American Drosophilidae. — The University of Texas Publication 5422: 47–64.
- Wheeler, M.R. 1959. A Nomenclatural Study of the Genus *Drosophila*. — The University of Texas Publication 5914: 181–205.
- Wheeler, M.R. 1960. Sternite modification in males of the Drosophilidae (Diptera). — Annals of the entomological Society of America 53: 133–137.
- Wheeler, M.R. 1965. Family Drosophilidae. In: Stone, A., Sabrosky, C.W., Wirth, W.W, Foote, R.H. & Coulson, J.R. (eds), A Catalog of the Diptera of America north of Mexico, pp. 760–771. — U. S. Department of Agriculture.
- Wheeler, M.R. 1981a. The Drosophilidae: A Taxonomic Overview. In: Ashburner, M., Carson, H.L. & Thompson, J.N. (eds), The Genetics and Biology of *Drosophila*, vol. 3a, pp. 1–97. — Academic Press.
- Wheeler, M.R. 1981b. Geographical survey of Drosophilidae: Nearctic species. In: Ashburner, M., Carson, H.L. & Thompson, J.N. (eds), The Genetics and Biology of *Drosophila*, vol. 3a, pp. 99–121. — Academic Press.
- Wheeler, M.R. & Kambysellis, M.P. 1966. Notes on the Drosophilidae (Diptera) of Samoa. — University of Texas Publication 6615: 533–565.
- Wright, S., Dobzhansky, T. & Hovanitz, W. 1942. Genetics of Natural Populations VII. The allelism of lethals in the third chromosome of *Drosophila pseudoobscura*. — Genetics 27: 363–394.
- Zelentsova, E.S., Vashakidze, R.P., Krayev, A.S. & Evgen'ev, M.B. 1986. Dispersed repeats in *Drosophila virilis*: Elements mobilized by interspecific hybridization. — Chromosoma 93: 469–476.

(received November 14, 2008; accepted November 16, 2008)