

<b>Zeitschrift:</b>	Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society
<b>Herausgeber:</b>	Schweizerische Entomologische Gesellschaft
<b>Band:</b>	82 (2009)
<b>Heft:</b>	3-4
<b>Artikel:</b>	On the identity of the olive psyllids in Iran (Hemiptera, Psylloidea)
<b>Autor:</b>	Asadi, Rahil / Talebi, Ali Ashgar / Burckhardt, Daniel
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-402988">https://doi.org/10.5169/seals-402988</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 31.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## On the identity of the olive psyllids in Iran (Hemiptera, Psylloidea)

RAHIL ASADI<sup>1,4</sup>, ALI ASHGAR TALEBI<sup>1</sup>, DANIEL BURCKHARDT<sup>2</sup>, JAFAR KHALGANI<sup>3</sup>, YAGHOUB FATHIPOUR<sup>1</sup> & SAEID MOHARRAMIPOUR<sup>1</sup>

<sup>1</sup> Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, P. O. Box: 1415-336, Tehran, Iran.

<sup>2</sup> Naturhistorisches Museum, Augustinergasse 2, CH-4001 Basel, Switzerland.

E-Mail: daniel.burckhardt@unibas.ch

<sup>3</sup> Agricultural Research, Education and Extension Organization, Ministry of Jihad-Keshavarzi, Keshavarz Boulevard, Tehran, Iran.

<sup>4</sup> E-Mail: rahil\_asadi@yahoo.com

Psyllids are probably the most serious pests of olive in Iran. Four names have been used for olive psyllids in Iran (*Euphyllura olivina*, *E. pakistanica*, *E. phillyreae* and *E. straminea*). During field surveys *E. straminea* was found in northern Iran (Provinces Gilan, Qazvin, Zanjan) and *E. pakistanica* in southern Iran (Province Fars). *E. olivina* and *E. phillyreae* do not occur in Iran. An identification table is provided for separating the morphologically similar *E. olivina*, *E. pakistanica* and *E. straminea*.

Keywords: Hemiptera, Psylloidea, *Euphyllura pakistanica*, *Euphyllura straminea*, olive pest, Iran, identification.

### INTRODUCTION

The olive (*Olea europaea* L.) is an economically important crop in Iran. In Fars (southern Iran), as well as Gilan, Qazvin and Zanjan (northern Iran) olive production is hampered by pests causing substantial losses in harvest. Olive psyllids are particularly harmful in the larval stages by removing plant sap and secreting honeydew. Honeydew serves as a medium for the growth of sooty mold which reduces photosynthesis (Zouiten & El Hadrami 2001). The infested trees are easily recognised by white flocculent waxy secretions which are excreted by the larvae (Abou-Kaf & Hamoudi 1999).

*Euphyllura olivina* (Costa) was recorded for the first time from Iran (southern provinces Fars and Kerman, northern province Gilan) by Farahbakhch & Moini (1975). Later Halperin *et al.* (1982) and Burckhardt & Lauterer (1993) added *E. straminea* Loginova. Gegechkori & Loginova (1990) mentioned *E. phillyreae* Foerster from Iran erroneously referring to Loginova (1972). According to Burckhardt & Lauterer (1993) the record of this species in Iran is doubtful. Noyes & Fallazadeh (2005) described an encyrtid parasitoid of *E. pakistanica* Loginova from the Fars province of Iran.

In summary, four names have been used for olive psyllids in Iran (*E. olivina*, *E. pakistanica*, *E. phillyreae* and *E. straminea*). Due to the continuous presence, high infestation and propagation rates resulting in fruit and bud loss, the psyllids are considered as the most important pests of olive trees in Iran. The objective of the present paper is to clarify the nomenclature and taxonomy of olive psyllids in Iran.

## MATERIAL AND METHODS

Surveys were conducted in olive orchards in 6 localities in southern Iran: province Fars (Fasa, Jahrom, Khafr, Sadra, Beiza and Shiraz) and 4 localities in northern Iran: province Gilan (Rudbar), province Qazvin (Qazvin) and province Zanjan (Zanjan, Tarom) (Tab. 1). Samples were collected in summer 2007 and spring 2008. The psyllids were identified by Daniel Burckhardt, Naturhistorisches Museum Basel, Switzerland (NHMB). For comparison material of *E. olivina* from France (NHMB) was studied and is illustrated below.

Tab. 1. Localities of surveyed olive orchards in southern (province Fars) and northern Iran (provinces Gilan, Qazvin and Zanjan) and sampled olive psyllid species (*Euphyllura*).

Province	locality	coordinates N/E in °	altitude in m above sea level	olive psyllid
Fars	Fasa	28.58/53.41	1370	<i>E. pakistanica</i>
Fars	Jahrom	29.92/53.33	1055	<i>E. pakistanica</i>
Fars	Khafr	29.03/53.60	1000	<i>E. pakistanica</i>
Fars	Shiraz	29.32/52.36	1550	<i>E. pakistanica</i>
Fars	Beiza	30.02/52.37	1598	<i>E. pakistanica</i>
Fars	Sadra	29.45/52.40	1580	<i>E. pakistanica</i>
Gilan	Rudbar	36.22/49.27	1230	<i>E. straminea</i>
Qazvin	Qazvin	36.15/50.30	1280	<i>E. straminea</i>
Zanjan	Zanjan	36.41/48.29	1350	<i>E. straminea</i>
Zanjan	Tarom	37.17/49.50	1250	<i>E. straminea</i>

## RESULTS

*Distribution*


All specimens collected in southern Iran (province Fars) belong to *E. pakistanica*, whereas those from northern Iran (provinces Gilan, Qazvin and Zanjan) are *E. straminea* (Tab. 1). Each sample contained larval stages as well as adult females and males. In the present study we did not find *E. olivina*.

*Identification*

*Euphyllura olivina*, *E. pakistanica* and *E. straminea* are morphologically similar. They can be identified by the pterostigma venation in the forewing and the male paramere shape (Tab. 2; Figs 1–6).

## CONCLUSIONS

Four names have been used for olive psyllids in Iran (*Euphyllura olivina*, *E. pakistanica*, *E. phillyreae* and *E. straminea*). The present study documents the occurrence of *E. pakistanica* in southern and *E. straminea* in northern Iran, respectively. *E. olivina* and *E. phillyreae* do not occur in Iran. *E. pakistanica* was reported from Pakistan, India, Syria and Iran (Loginova 1973; Thakur *et al.* 1997; Noyes & Fallazadeh 2005). In India, it is the most important pest of olive (Virender *et al.* 2007). *E. olivina*, *E. straminea* and *E. phillyreae* are known from the mediterranean



Figs 1–6. Morphological characters of three *Euphyllura* spp. — 1, 4, *E. olivina*; 2, 5, *E. pakistanica*; 3, 6, *E. straminea*. — 1–3, Forewing, arrow indicating apex of pterostigma; 4–6, paramere, lateral view.

region, the most important region of olive cultivation (Abou-Kaf & Hamoudi 1999; Zouiten & El Hadrami 2001; Cotes *et al.* 2007; Kumral *et al.* 2008).

Our findings will be useful in biological control and integrated pest management (IPM) of olive trees in Iran. In fact, the first and crucial step in any IPM programme is to accurately identify the pests (Adams & Clark 1996).

Tab. 2. Morphological characters separating *Euphyllura olivina*, *E. pakistanica* and *E. straminea*.

Morphological characters	<i>E. olivina</i>	<i>E. pakistanica</i>	<i>E. straminea</i>
<b>forewing</b>	Fig. 1	Fig. 2	Fig. 3
pterostigma length	long, see arrow	short, see arrow	long, see arrow
number of veins on pterostigma	many	few or absent	many
<b>paramere</b>	Fig. 4	Fig. 5	Fig. 6
apex	subacute	obliquely truncate	obliquely truncate
foremargin	angular	sinuous	broadly rounded, convex

## ACKNOWLEDGMENTS

We thank Igor Malenovský (Brno) and Denise Wyniger (Lucerne) for useful comments on an earlier manuscript version.

## REFERENCES

- Adams, R.G. & Clark, C.J. 1996. Northeast Sweet Corn Production and Integrated Pest Management Manual. — University of Connecticut Cooperative Extension. Storrs, CT, 120 pp.
- Abou-Kaf, N. & Hamoudi, O. 1999. Evaluation of damage caused by olive psylla *Euphyllura straminea* Loginova (Hom.: Aphalaridae) in Syria. — Arabian Journal of Plant Protection 17: 71–76.
- Burckhardt, D. & Lauterer, P. 1993. The jumping plant-lice of Iran (Homoptera, Psyllidae). — Revue suisse de Zoologie 100: 829–898.
- Cotes, B., Ruano, F., García, P., Pascual, F., Tinaut, A., Pepa, A. & Campos, M. 2007. Differences in insects within the olive orchard agroecosystem under integrated management regime in south Spain. — Bulletin OILB/SROP 30: 47.
- Farahbakhch, G. & Moini, M. 1975. Olive pests in Iran. — Plant Pests and Diseases Research Institute, National Agricultural Research Organisation, Ministry of Agriculture and Natural Resources, Tehran, Iran, 73 pp.
- Gegechkori, A.M. & Loginova, M.M. 1990. Psyllid (Homoptera, Psylloidea) SSSR (annotirovannyi spisok). — Metsniereba, Tbilisi, 164 pp. (in Russian).
- Halperin, J., Hodgkinson, I.D., Russell, L.M. & Berlinger, M.J. 1982. A contribution to the knowledge of the psyllids of Israel (Homoptera: Psylloidea). — Israel Journal of Entomology 16: 27–44.
- Kumral, N.A., Kovancı, B. & Akbudak, B. 2008. Using degree-day accumulations and host phenology for predicting larval emergence patterns of the olive psyllid, *Euphyllura phillyreae*. — Journal of Pest Science 81: 63–69.
- Loginova, M.M. 1972. On the fauna of Psylloidea (Homoptera) from Morocco. — Commentationes Biologicae 47, 37 pp.
- Loginova, M.M. 1973. Taxonomy of the tribe Euphyllurini (Psylloidea, Homoptera). — Zoolo-gicheskii Zhurnal 52(6): 853–869. (In Russian)
- Noyes, J.S. & Fallazadeh, M. 2005. *Psyllaephagus zdeneki* sp. nov. (Hymenoptera: Encyrtidae) from Iran, parasitoid of *Euphyllura pakistanica* (Hemiptera: Psyllidae). — Acta Societatis Zoologicae Bohemicae 69: 203–208.
- Thakur, J.R., Atwal, R.K. & Gupta, P.R. 1997. Insect pests of olive in Himachal Pradesh. Pest management and economic entomology. — Pest Management and Economic Zoology 5: 75–79.
- Virender, K., Malik, G.H., Uma, S. & Monobrullah, M. 2007. Incidence and management of olive psylla, *Euphyllura pakistanica*. — Indian Journal of Entomology 69: 331–340.
- Zouiten, N. & El Hadrami, I. 2001. La psylle de l'olivier: état des connaissances et perspectives de lutte. — Cahiers Agricultures 10(4): 225–232.

(received August 12, 2009; accepted September 11, 2009)