

Zeitschrift:	Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society
Herausgeber:	Schweizerische Entomologische Gesellschaft
Band:	82 (2009)
Heft:	3-4
Artikel:	Some taxonomical changes in the genus Rhaphium (Diptera, Dolichopodidae), with a key to the Rhaphium albifrons species group
Autor:	Naglis, Stefan
DOI:	https://doi.org/10.5169/seals-402989

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 31.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Some taxonomical changes in the genus *Rhaphium* (Diptera, Dolichopodidae), with a key to the *Rhaphium albifrons* species group.

STEFAN NAGLIS

Zoological Museum, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.

E-mail: s.naglis@bluewin.ch

The following taxonomical changes in the genus *Rhaphium* are proposed: *Rhaphium fluviatile* Vaillant syn. n. of *Rhaphium pectinatum* (Becker) nec *Rhaphium pectinatum* (Loew), *Rhaphium sagittiferum* Vaillant syn. n. of *Rhaphium albifrons* Zetterstedt, *Rhaphium montium* Vaillant syn. n. of *Rhaphium albomaculatum* (Becker), *Systemus viklundi* Grichanov comb. n., probably a synonym of *Systemus bipartitus* (Loew). A key to Palaearctic species of the *Rhaphium albifrons* species group based on male genitalia characters is provided.

Keywords: Dolichopodidae, *Rhaphium*, Palaearctic, synonyms, key.

INTRODUCTION

The genus *Rhaphium* Meigen, 1803 includes 185 species worldwide which occur in all biogeographical regions, except the Australasian and Oceanian regions, and is represented in the Palaearctic by 89 species (Yang *et al.* 2006). Pollet (2004) listed 49 species for Europe, but he did not include the three species described by Vaillant (1983). In Switzerland 20 species have been recorded so far (Naglis 2009).

SYSTEMATIC ACCOUNT

The Palaearctic species of the genus *Rhaphium* were revised by Negrobov (1979), including a key. Since then the following species have been described from Central and Western Europe:

- Rhaphium fluviatile* Vaillant, 1983: 272 (France)
- Rhaphium montium* Vaillant, 1983: 273 (France)
- Rhaphium sagittiferum* Vaillant, 1983: 273 (France)
- Rhaphium viklundi* Grichanov, 2004: 268 (Sweden)
- Rhaphium latimanum* Kahanpaa, 2007: 228 (Finland)

Because the publications of Negrobov (1979) and Vaillant (1983) include adequate drawings of the genitalia of the *Rhaphium* species, a study of the type material seems unnecessary. By comparing the characters, in particular those of the genitalia, the following decisions can be taken:

- (i) Negrobov (1979) transferred all species of *Xiphandrium* and *Porphyrops* to *Rhaphium* and synonymised *Xiphandrium pectinatum* (Becker, 1908) with *R. bre-*

vicorne Curtis, 1835. In his description of *R. fluviatile*, Vaillant (1983) noted the similarity with *R. brevicorne* but mentions differences in the shape of the surstyli, where the ventral projection is longer than the dorsal one, whereas in *R. brevicorne* the length of the ventral projection is shorter than the dorsal one. This is in fact the case in the figure in Negrobov (1979, fig. 1692). Vaillant's figure of the surstyli is identical with that of *Xiphandrium pectinatum* Becker, figured in Becker (1918). An additional character for separating *R. brevicorne* from *R. pectinatum* is the length of the first flagellomere which is about 3 times as long as its basal width in *R. pectinatum* and 5 times as long in *R. brevicorne*. Therefore I herewith remove *R. pectinatum* Becker from synonymy with *R. brevicorne*. Since the combination *Rhaphium pectinatum* is preoccupied by Loew, 1859, the actual name is that of its synonym *R. fluviatile*.

Rhaphium fluviatile Vaillant, 1983: 272
= *Rhaphium pectinatum* (Becker, 1908: 49), **syn. n.**
preocc. by *Rhaphium pectinatum* (Loew, 1859: 16)

(ii) Vaillant's (1983) description of *R. sagittiferum* includes figures of the antenna and male genitalia. Both the antenna and the hypopygium are identical with *R. albifrons* Zetterstedt as described and figured in Negrobov (1979). Therefore I herewith synonymise *R. sagittiferum* with *R. albifrons*.

Rhaphium albifrons Zetterstedt, 1843: 479
= *Rhaphium sagittiferum* Vaillant, 1983: 273, **syn. n.**

(iii) Vaillant's (1983) description and figures of *R. montium*, especially the male hypopygium is identical with the description and the figures of the cerci and the surstyli of *R. albomaculatum* Becker in Negrobov (1979, figs 1663–1665). Therefore I herewith synonymise *R. montium* with *R. albomaculatum*.

Rhaphium albomaculatum (Becker, 1891: 291)
= *Rhaphium montium* Vaillant, 1983: 273, **syn. n.**

(iv) Grichanov (2004) figures the first flagellomere in his original description of *Raphium viklundi* which is typical for the genus *Systemus* and, moreover, the figure of the male hypopygium may be identical with that of *Systemus bipartitus* Loew, 1859 as figured in Negrobov (2005).

Systemus viklundi (Grichanov, 2004: 268), **comb. n.**
? = *Systemus bipartitus* (Loew, 1850: 114)

The *Rhaphium albifrons* species group

Grichanov (2004) provided a key to the Swedish species of the *Rhaphium albifrons* group but gave no diagnosis. The species *R. albifrons* Zetterstedt, *R. albomaculatum* Becker, *R. fissum* Loew and *R. gruniniani* Negrobov are herewith combined in the *Rhaphium albifrons* species group, defined by the following combination of characters: hind coxa with a strong white lateral seta; fore basitarsus without comb of strong setae; frons usually with dense white or grey pruinosity; arista at most twice as long as or shorter than basal width of first flagellomere.

Key to males of the *Rhaphium albifrons* species group

The traditionally used characters, the length of the arista and the first flagellomere relative to its basal width, are somewhat difficult to use because the antennae are often flexed or shrunken in dry specimens. Also the colouration of the femora shows a certain variation. Therefore key to genitalia features is here provided, in addition to Negrobov's (1979) key (couplets 61–63):

1. Median lobe of cercus 4 times as long as lateral lobe; surstyli broadened apically *R. albomaculatum* (Becker)
- Median lobe of cercus about 2 times as long as lateral lobe; surstyli tapered apically 2
2. Surstyli pointed apically 3
- Surstyli stout and rounded apically *R. gruniniani* Negrobov
3. Surstyli basally broad, tapering from mid-point *R. fissum* Loew
- Surstyli slender from base to tip *R. albifrons* Zetterstedt

REFERENCES

- Becker, T. 1908. Dipteren der Kanarischen Inseln. — Mitteilungen des zoologischen Museums Berlin 4(1): 1–180.
- Becker, T. 1918. Dipterologische Studien. Dolichopodidae. A. Paläarktische Region. Zweiter Teil. — Nova Acta Academiae Caesareae Leopoldino Carolinae 103(3): 203–315.
- Grichanov, I.Ya. 2004. A list of Dolichopodidae from the Tyresta National Park, Sweden, with description of a new species of the genus *Rhaphium* Meigen (Diptera). — Zoosystematica Rossica 2003, 12(2): 267–269.
- Naglis, S. 2009. Checkliste der Langbeinfliegen der Schweiz (Diptera, Dolichopodidae). [Checklist of the long-legged flies of Switzerland (Diptera, Dolichopodidae)]. — Entomo Helvetica, 2: 17–22.
- Negrobov, O.P. 1979. 29. Dolichopodidae. In: Lindner, E.: Die Fliegen der palaearktischen Region. 4(5), Lief. 322, pp. 475–530. — Schweizerbart, Stuttgart.
- Negrobov, O.P. 2005. Species of the genus *Systemus* (Dolichopodidae, Diptera) from the Palaearctic Region. — Zoologicheskii Zhurnal 84(11): 1421–1424 (in Russian). [English translation: Entomological Review, 85(7): 826–830.]
- Pollet, M. 2004. Dolichopodidae. In: Pape, T. (ed.), Fauna Europaea: Diptera Brachycera. — Fauna Europaea version 1.3, <http://www.faunaeur.org>. (accessed September 2009)
- Vaillant, F. 1983. Sur les Insectes Diptères Dolichopodidae de la Sous-Famille des Rhaphiinae. Ministère de l'Education Nationale, Comité des Travaux Historiques et Scientifiques — Comptes Rendus du 108^e Congrès National des Sociétés Savantes. Section des Sciences, Fascicule 2, pp. 269–280.
- Yang, D., Zhu, Y., Wang, M. & Zhang, L. 2006. World Catalog of Dolichopodidae (Insecta: Diptera). — China Agricultural University Press, 704 pp.

(received July 7, 2009; accepted after revision September 24, 2009)