Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =

Bulletin de la Société Entomologique Suisse = Journal of the Swiss

Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 82 (2009)

Heft: 3-4

Artikel: Psyllids (Hemiptera, Psylloidea) captured in commercial apple and

stone fruit orchards in southwest Germany, eastern France and

northwest Switzerland

Autor: Jarausch, Barbara / Burckhardt, Daniel / Lauterer, Pavel

DOI: https://doi.org/10.5169/seals-402990

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

82: 205-215, 2009

Psyllids (Hemiptera, Psylloidea) captured in commercial apple and stone fruit orchards in southwest Germany, eastern France and northwest Switzerland

Barbara Jarausch¹, Daniel Burckhardt², Pavel Lauterer³ & Wolfgang Jarausch¹

- ¹ RLP AgroScience, AlPlanta-Institute for Plant Research, Breitenweg 71, D-67435 Neustadt, Germany. E-Mail: barbara.jarausch@agroscience.rlp.de.
- ² Naturhistorisches Museum, Augustinergasse 2, CH-4001 Basel, Switzerland. E-Mail: daniel.burckhardt@unibas.ch.

Psyllids are vectors of important phytoplasma diseases of apple and stone fruits. From 2001 to 2007 psyllid captures were carried out in commercial apple and stone fruit orchards in southwest Germany (D), Alsace (F) and northwest Switzerland (CH). Psyllids were also collected from wild *Prunus* species in the neighbourhood of commercial stone fruit orchards. In total, data were obtained from 94 sites in 14 districts in Germany (Baden-Württemberg, Rheinland-Pfalz), France (Bas-Rhin) and Switzerland (Aargau, Solothurn). Twenty five psyllid species were captured on apple. The confirmed phytoplasma vector *Cacopsylla picta* and *C. melanoneura*, a vector in northwest Italy, were the most abundant species followed by *C. mali* and *C. pulchra*. Twenty one psyllid species were found on the stone fruit species *Prunus amygdalus* (almond), *P. armeniaca* (apricot), *P. persica* (peach), *P. domestica* (European plum), *P. cerasifera* (cherry plum) and *P. spinosa* (blackthorn). The most abundant species was again a phytoplasma vector species: *C. pruni*. This species as well as *C. melanoneura* were captured on all tested *Prunus* species. A detailed list of the abundance and geographic distribution of the psyllid species in the studied regions is given. This list will give valuable information to advisors and growers less experienced in psyllid identification on which psyllid species may occur in their orchard when looking for phytoplasma vector species.

Keywords: *Malus*, *Prunus*, phytoplasma, apple proliferation, European stone fruit yellows, *Cacopsylla picta*, *Cacopsylla pruni*, *Cacopsylla melanoneura*, distribution, Central Europe.

INTRODUCTION

Apple proliferation (AP) and European stone fruit yellows (ESFY) are serious diseases of *Malus* and *Prunus*, respectively, leading to important economic losses in Central and Southern Europe. Both diseases are caused by phytoplasmas: AP by '*Candidatus* Phytoplasma mali' and ESFY by '*Candidatus* Phytoplasma prunorum' (Seemüller & Schneider 2004). These are transmitted by psyllids in a persistent manner, i.e. '*Ca*. P. mali' by *Cacopsylla picta* (Foerster) (Frisinghelli *et al*. 2000) and, in northwest Italy, by *C. melanoneura* (Foerster) (Tedeschi *et al*. 2002) and '*Ca*. P. prunorum' by *C. pruni* (Scopoli) (Carraro *et al*. 1998).

AP has been known in southwest Germany and neighbouring regions for decades. It re-emerges from time to time in serious epidemics (Bliefernicht & Krczal 1995). The last epidemic started in 2000 (Harzer 2000). Jarausch *et al.* (2003, 2004) identified *C. picta* as vector of '*Ca.* P. mali' in Germany. ESFY is a serious disor-

³ Moravian Museum, Department of Entomology, Hviezdoslavova 29a, CZ- 627 00 Brno-Slatina, Czech Republic. E-Mail: ento.laut@volny.cz.

Tab. 1. Numbers and types of orchards were psyllids were collected on apple (*Malus x domestica* Borkh.). Abbreviations: D = **Germany**: BW = Baden-Württemberg: FR = Freiburg/Breisgau, HD = Heidelberg, Rhein-Neckar-Kreis, KA = Karlsruhe, OG = Ortenaukreis (Offenburg); – RP = Rheinland-Pfalz: DÜW = Bad Dürkheim/Weinstrasse, MYK = Mayen-Koblenz, Andernach, MZ = Mainz, Mainz-Bingen, NW = Neustadt/Weinstrasse, SÜW = Südliche Weinstrasse (Landau); – F = **France**: BR = Bas-Rhin; – CH = **Switzerland**: AG = Aargau, SO = Solothurn.

Country	state	district	type of orcha	ırd			total
			commercial	organic farming	abandonned	scattered	
D	BW	HD	2			1	3
	BW	KA	1				1
	BW	OG	1			1	2
	BW	FR	1				1
	RP	MYK		1			1
	RP	MZ	4	1			5
	RP	DÜW	5	2	1	6	14
	RP	NW	5		2	5	12
	RP	SÜW	6				6
F	BR		2			1	3
СН	AG		1				1
	SO					1	1
total							50

der of apricots and Japanese plum in the Mediterranean Region and was only recently recognized as a problem in Switzerland (Ramel & Gugerli 2004) and Germany (Jarausch *et al.* 2007b). *C. pruni* was confirmed as vector of '*Ca.* P. prunorum' in both countries. However, nothing was known about the distribution of the vectors in commercial apple and stone fruit orchards in Germany, France and Switzerland. There was also a lack of information on the potential critical role played by wild *Prunus* species on the populations of *C. pruni* in orchards.

Two issues are crucial for the successful application of control measures against the vector species: 1 – the knowledge on abundance and geographical distribution of the vectors and 2 – the correct identification of the vectors by advisors and growers. The identification of psyllids is often demanding for the non-specialist. For this reason Burckhardt (2007) designed a simplified electronic identification key available on the internet. Knowing which psyllid species occur in a particular area greatly simplifies identifying specimens to species.

The present paper lists psyllid species found during vector monitoring in apple and stone fruit orchards in southwest Germany, eastern France and northwest Switzerland. This information reflects the current distribution of the vector species in this area.

MATERIAL AND METHODS

From 2001 to 2007 psyllids were monitored in 50 apple orchards in southwest Germany, eastern France and northwest Switzerland (Tab. 1) and in 44 stone fruit orchards in southwest Germany (Tab. 2). The orchards were situated in **Germany** (= D) in the following 2 states (Bundesland) and 11 districts (Stadt-/Landkreis, cited according to official car registration numbers): Baden-Württemberg (BW): Freiburg/Breisgau (= FR), Heidelberg, Rhein-Neckar-Kreis (= HD), Karlsruhe (= KA),

Tab. 2. Numbers and types of orchards in Germany (Rheinland-Pfalz) were psyllids were collected on various *Prunus* species. Abbreviations: DÜW = Bad Dürkheim/Weinstrasse, GER = Germersheim, MYK = Mayen-Koblenz, Andernach, MZ = Mainz, Mainz-Bingen, NW = Neustadt/Weinstrasse, RP = Rhein-Pfalz-Kreis, SÜW = Südliche Weinstrasse (Landau).

Prunus species	district	type of orchard			total
•		commercial		wild plants	
P. amygdalus Batsch	NW		3		3
P. armeniaca L.	MYK	3			3
	MZ	6			6
	NW	2			2
	RP	1			1
	GER	1			1
P. cerasifera Ehrh.	MZ			1	1
	DÜW			1	1
	NW			3	3
P. domestica L.	MYK	4			4
	NW	1			1
P. insititia Linnaeus	NW	1			1
P. persica (L.) Batsch	MYK	1			1
	NW	2			2
	RP	2			2
	GER	1			1
P. spinosa L.	MYK			2	2
	MZ			1	1
	DÜW			1	1
	NW			3	3
	SÜW			1	1
P. cerasifera and P. spinosa	MYK			1	1
	NW			1	1
Prunus rootstock suckers	NW	1			1
total				***************************************	44

Ortenaukreis (Offenburg) (= OG); – Rheinland-Pfalz (= RP): Bad Dürkheim/Weinstrasse (= DÜW), Germersheim (= GER), Mayen-Koblenz, Andernach (= MYK), Mainz, Mainz-Bingen (= MZ), Neustadt/Weinstrasse (= NW), Rhein-Pfalz-Kreis (= RP), Südliche Weinstrasse (Landau) (= SÜW); in **France** (= F) in the state (département) Bas-Rhin (= BR); and in **Switzerland** (= CH) in the states (cantons) Aargau (= AG) and Solothurn (= SO).

The psyllids were collected using the beating tray method described by Müther & Vogt (2002). One hundred branches were sampled per sampling date and orchard. Psyllid collections were done in reference orchards and wild locations near Neustadt at weekly intervals during 2001–2005. Psyllid captures in all other sites were done in the period from March to July when the vector species are present on their host plant. These captures were performed weekly for reference orchards of every district and irregularly in the other sites.

RESULTS

Faunal composition

Malus. In total 25 psyllid species of 8 genera were collected (Tab. 3). Three species, i.e. Cacopsylla mali, C. melanoneura and C. picta were represented by high numbers of individuals (> 1000). The remainder of species was represented by 100–1000 specimens (1 species), 10–100 specimens (8 species) and < 10 specimens (13 species). Most species develop on woody plants (22 species) and only 3 on herbs (Aphalara polygoni, Bactericera femoralis and Trioza urticae). The species developing on woody Rosaceae are best represented in terms of individuals (11'073) comprising 10 Cacopsylla spp.: C. affinis, C. breviantennata, C. crataegi, C. mali, C. melanoneura, C. picta, C. pruni, C. pyri, C. pyricola and C. pyrisuga. The second most important component is an assemblage of 4 species associated with Salicaceae (Bactericera albiventris, Cacopsylla brunneipennis, C. pulchra and C. saliceti), represented by 220 specimens. Fifteen species overwinter as adults on conifers.

Prunus. In total 21 psyllid species of 6 genera were collected (Tab. 4). One species, i.e. Cacopsylla pruni was represented by high numbers of individuals (> 1000). The remainder of species was represented by 100–1000 specimens (1 species), 10–100 specimens (7 species) and < 10 specimens (12 species). Nineteen species develop on woody plants and only 2 on herbs (Aphalara freji and Trioza urticae). The species developing on woody Rosaceae were best represented in terms of individuals (4737) comprising 10 Cacopsylla spp.: C. affinis, C. crataegi, C. mali, C. peregrina, C. melanoneura, C. picta, C. pruni, C. pyri, C. pyricola and C. pyrisuga. There were 5 species associated with Salicaceae (Bactericera albiventris, Cacopsylla ambigua, C. brunneipennis, C. pulchra and C. saliceti), represented by 101 specimens. Thirteen species overwinter as adults on conifers.

By far the largest number of adult *C. pruni* were collected on *Prunus spinosa* (> 1000 inididuals), followed by *P. armeniaca*, *P. cerasifera* and *P. domestica* (100–1000 individuals), *P. persica* (10–100 individuals) and *P. amygdalus* and *P. insititia* (1–10 individuals).

Distribution and abundance

Malus. The confirmed vector species of AP, C. picta, and the AP-vector in northwest Italy, C. melanoneura, were detected in all surveyed districts in southwest Germany, Alsace and northwest Switzerland. C. mali, whose host plant is apple, was found in 8 out of 12 districts. This species was very abundant in untreated orchards. The majority of the other species were captured occasionally in few districts only. Exceptions were two very common Trioza species: T. remota and T. urticae. The pear psyllid C. pyrisuga was also regularly captured (7 out of 12 districts) despite its low abundance. C. pulchra, developing on Salix spp., was found in important numbers in southwest Germany.

Prunus. The confirmed vector species of ESFY, *C. pruni*, was most abundant and was detected in 6 out of 7 surveyed districts of southwest Germany as well as on all examined *Prunus* species. *C. melanoneura* was the second most abundant species which was detected in all 7 districts and on almost all *Prunus* species. All other species were found more irregularly in lower numbers of specimens. Particular high numbers of *C. pruni* were captured on wild *P. spinosa* and on hedges where *P. spinosa* was present.

Tab. 3. Psyllids found in orchards on *Malus* x *domestica* Borkh. (= syn. *Malus pumila* Mill.). Abbreviations: D = **Germany**: BW = Baden-Württemberg: FR = Freiburg/Breisgau, HD = Heidelberg, Rhein-Neckar-Kreis, KA = Karlsruhe, OG = Ortenaukreis (Offenburg); – RP = Rheinland-Pfalz: DÜW = Bad Dürkheim/Weinstrasse, MYK = Mayen-Koblenz, Andernach, MZ = Mainz, Mainz-Bingen, NW = Neustadt/Weinstrasse, SÜW = Südliche Weinstrasse (Landau); – F = **France**: BR = Bas-Rhin; – CH = **Switzerland**: AG = Aargau, SO = Solothurn.

Psyllid species	number of specimens	country	state	district	number of sites
Aphalara polygoni (Foerster) = syn. A. rumicicola Klimaszewski	2	D	RP	DÜW	1
		D	BW	KA	1
Arytainilla spartiophila (Foerster)	3	D	RP	NW	1
Bactericera albiventris (Foerster)	15	D	RP	DÜW, NW, SÜW	7
		D	BW	HD	1
Bactericera femoralis (Foerster)	1	D	RP	DÜW	1
Cacopsylla affinis (Löw)	9	D	RP	DÜW, NW, SÜW	6
		D	BW	HD, OG	2
Cacopsylla breviantennata (Flor)	1	СН	AG		1
Cacopsylla brunneipennis (Zetterstedt)	5	D	RP	NW	1
Cacopsylla crataegi (Schrank)	11	D	RP	DÜW, SÜW	6
		D	BW	HD, KA	2
Cacopsylla mali (Schmidberger)	5187	D	RP	DÜW, MYK, MZ, NW, SÜW	15
		D	BW	HD, KA	2
		СН	SO	110,111	1
Cacopsylla melanoneura (Foerster)	4272	D	RP	DÜW, MYK, MZ, NW, SÜW	32
		D	BW	FR, HD, KA, OG	7
		F	BR		3
		СН	AG		1
		СН	SO		1
Cacopsylla picta (Foerster)	1506	D	RP	DÜW, MYK, MZ, NW, SÜW	33
		D	BW	FR, HD, KA, OG	7
		F	BR		3
		СН	AG		1
		СН	SO		1
Cacopsylla pruni (Scopoli)	22	D	RP	DÜW, NW, SÜW	11
		СН	AG		1
Cacopsylla pulchra (Zetterstedt)	166	D	RP	DÜW, NW, SÜW	12
		D	BW	HD	1
Cacopsylla pyri (Linnaeus)	41	D	RP	NW, SÜW	4
		D	BW	KA, OG	2
		СН	AG		1

Cacopsylla pyricola (Foerster)	4	D	BW	HD, KA	2
Cacopsylla pyrisuga (Foerster)	20	D	RP	DÜW, NW, SÜW	11
		D	BW	HD, OG	2
		СН	AG		1
		СН	SO		1
Cacopsylla rhamnicola (Scott)	3	D	RP	DÜW	1
		D	BW	OG	1
Cacopsylla saliceti (Foerster)	38	D	RP	DÜW, NW, SÜW	4
		F	BR		3
Psylla alni (Linnaeus)	1	D	RP	SÜW	1
Psyllopsis fraxini (Linnaeus)	1	F	BR		1
Psyllopsis fraxinicola (Foerster)	1	D	BW	KA	1
Rhinocola aceris (Linnaeus)	1	D	BW	KA	1
Trioza remota Foerster	67	D	RP	DÜW, MZ, NW, SÜW	13
		D	BW	HD, KA, OG	5
		СН	AG		1
		СН	SO		1
Trioza rhamni (Schrank)	2	D	AG		1
Trioza urticae (Linnaeus)	68	D	RP	DÜW, NW, SÜW	10
		D	BW	HD, KA, OG	3
		СН	AG		1
		СН	SO		1

DISCUSSION

In total 25 psyllid species of 8 genera were collected on apple and 21 psyllid species of 6 genera on *Prunus* spp., respectively. They all represent common and wide-spread species in Central Europe (Burckhardt 2004). By far the most individuals collected on apple belong to Cacopsylla mali, C. melanoneura and C. picta which develop only or also on apple. A comparable result was found on *Prunus* spp. were the vast majority of individuals were C. pruni which develops only on this host genus. This result was expected. Fifteen species collected on apple and thirteen species collected on *Prunus* spp. overwinter as adults on conifers. Adults of these species migrate in summer/autumn from their host plants to conifers and in the following spring back to their hosts over long distances (Čermák & Lauterer 2008). In spring, before reaching their host plants, they may be found accidentally also on other plants. The polyvoltine species C. pyri and C. pyricola which overwinter on their host, Pyrus spp., have short distance migratory phases and accordingly the two species have been found predominantly in stone fruit orchards where pear orchards were in the surroundings. The findings of the vast majority of species are accidental, suggesting that additional similar sampling effort would add more species.

Tab. 4. Psyllids collected on different *Prunus* species in and around orchards in Germany (Rheinland-Pfalz). Abbreviations: DÜW = Bad Dürkheim/Weinstrasse, GER = Germersheim, MYK = Mayen-Koblenz, Andernach, MZ = Mainz, Mainz-Bingen, NW = Neustadt/Weinstrasse, RP = Rhein-Pfalz-Kreis, SÜW = Südliche Weinstrasse (Landau).

Prunus species	psyllid species	number of specimens	district	number of sites
P. amygdalus Batsch	Cacopsylla melanoneura (Foerster)	2	NW	1
	Cacopsylla pruni (Scopoli)	10	NW	4
	Psylla buxi (Linnaeus)	1	NW	1
P. armeniaca L.	Cacopsylla affinis (Löw)	1	NW	1
	Cacopsylla crataegi (Schrank)	2	MZ, NW	1
	Cacopsylla mali (Schmidberger)	5	MZ	1
	Cacopsylla melanoneura (Foerster)	28	GER, MZ, NW, RP	5
	Cacopsylla peregrina (Foerster)	8	GER, MZ	3
	Cacopsylla picta (Foerster)	1	MZ	1
	Cacopsylla pruni (Scopoli)	539	DÜW, GER, MYK, MZ, NW	11
	Cacopsylla pulchra (Zetterstedt)	10	GER, NW	3
	Cacopsylla pyri (Linnaeus)	26	GER	1
	Cacopsylla pyricola (Foerster)	4	GER, NW	2
	Cacopsylla saliceti (Foerster)	3	NW	1
	Trioza remota Foerster	10	GER, MZ, NW	4
P. cerasifera Ehrh.	Bactericera albiventris (Foerster)	1	NW	1
	Cacopsylla affinis (Löw)	1	NW	1
	Cacopsylla melanoneura (Foerster)	211	DÜW, MZ, NW	6
	Cacopsylla pruni (Scopoli)	310	DÜW, MZ, NW	6
	Cacopsylla pulchra (Zetterstedt)	11	NW	3
	Cacopsylla pyri (Linnaeus)	5	NW	1
	Cacopsylla rhamnicola (Scott)	4	NW	1
	Cacopsylla saliceti (Foerster)	1	NW	1
	Trioza remota Foerster	1	NW	1
	Trioza urticae (Linnaeus)	4	NW	2
P. domestica L.	Cacopsylla melanoneura (Foerster)	7	NW	1
	Cacopsylla pruni (Scopoli)	348	MYK, NW	5
	Cacopsylla pulchra (Zetterstedt)	5	NW	1
	Trioza remota Foerster	1	NW	1
P. insititia Linnaeus	Cacopsylla pruni (Scopoli)	8	NW	1
P. persica (L.) Batsch	Bactericera albiventris (Foerster)	1	GER	1
	Cacopsylla affinis (Löw)	1	NW	1
	Cacopsylla melanoneura (Foerster)	33	NW, RP	3

	Cacopsylla pruni (Scopoli)	54	DÜW,	6
			GER,	
			MYK,	
			NW	
	Cacopsylla pulchra (Zetterstedt)	14	GER, NW	3
	Cacopsylla pyri (Linnaeus)	41	GER	2
	Cacopsylla rhamnicola (Scott)	2	GER, NW	2
	Cacopsylla saliceti (Foerster)	7	NW, RP	2
	Trioza remota Foerster	10	GER, NW	2
P. spinosa L.	Bactericera albiventris (Foerster)	1	MZ	1
11500000	Cacopsylla affinis (Löw)	1	NW	1
	Cacopsylla crataegi (Schrank)	3	DÜW,	3
	eucopsynu erunegi (semunk)	3	MZ,	3
			SÜW	
	Cacopsylla melanoneura (Foerster)	318	DÜW,	6
	cacopsylla metanoneura (1 ocisici)	310	MZ, NW,	U
			SÜW	
	Cacopsylla picta (Foerster)	1	MZ	1
	Cacopsylla pruni (Scopoli)	1217	DÜW,	8
	cacopsyna pram (Geopon)	1217	MYK,	O
			NW,	
			SÜW	
	Cacopsylla pulchra (Zetterstedt)	12	NW	2
	Cacopsylla pyricola (Foerster)	2	NW	1
	Cacopsylla pyrisuga (Foerster)	1	NW	1
	Cacopsylla saliceti (Foerster)	1	NW	1
	Rhinocola aceris (Linnaeus)	1	NW	1
	Trioza urticae (Linnaeus)	3	NW	2
	Trioza arricae (Elimacus)		14 44	
sites with P. spinosa	Aphalara freji Burckhardt &	1	MYK	1
and <i>P. cerasifera</i>	Lauterer			
	Cacopsylla affinis (Löw)	3	NW	1
	Cacopsylla ambigua (Foerster)	1	MYK	1
	Cacopsylla brunneipennis	2	MYK,	2
	(Zetterstedt)		NW	
	Cacopsylla crataegi (Schrank)	2	NW	1
	Cacopsylla mali (Schmidberger)	7	MYK,	2
	(NW	
	Cacopsylla melanoneura (Foerster)	55	MYK,	2
			NW	
	Cacopsylla picta (Foerster)	1	NW	1
	Cacopsylla pruni (Scopoli)	1109	NW	2
	Cacopsylla pulchra (Zetterstedt)	31	MYK,	2
		90% (5)	NW	
	Cacopsylla pyri (Linnaeus)	2	NW	1
	Cacopsylla pyrisuga (Foerster)	3	NW	1
	Cacopsylla rhamnicola (Schott)	4	NW	1

	Trioza urticae (Linnaeus)	15	NW	1
rootstock suckers of different <i>Prunus</i> domestica and other <i>Prunus</i> hybrids	Cacopsylla melanoneura (Foerster)	11	NW	1
•	Cacopsylla pruni (Scopoli)	356	NW	1

The psyllid captures were focused on the detection of the known vector species of AP and ESFY in the different regions. The population dynamics of *C. picta* and *C. melanoneura* on apple as well as those of *C. pruni* on *Prunus* were studied in the years 2001–2005 by regular insect captures during the entire year. These data have been published by Jarausch (2003) and Mayer *et al.* (2009) and demonstrate that overwintered adults of *C. melanoneura* are the first to remigrate from conifers to the orchards in February to beginning of March, followed by overwintered adults of *C. picta* which can be found from the middle of March onwards. The data for *C. melanoneura* were confirmed by Mayer & Gross (2007). Emerging adult *C. melanoneura* and *C. picta* were observed from April to June or May to July, respectively. From the end of May onwards emerging adults of *C. mali* can be found (Jarausch 2003) and may be confused with emerging adult *C. picta*. The population dynamics of *C. pruni* on *Prunus* are similar to those described for *C. picta* (Jarausch *et al.* 2007b).

The psyllid captures in the other regions and districts were restricted to the period (March to July) when the vector species are present on their respective host plant. Other psyllid species with a different life cycle may, therefore, be underrepresented in the captures. The data of psyllid collections made during the entire year in reference orchards near Neustadt indicate that no other psyllid species are dominant on *Malus* and *Prunus*. An exception is the situation, when the dominant psyllid species have been eradicated by insecticides and other psyllid species from the surroundings colonize the fruit trees for a short period of time (P. Lauterer, unpublished data). Thus, with the obvious exception of *C. mali* developing on apple, the phytoplasma vector species are the predominant species in apple and stone fruit orchards. *C. mali* is not abundant in treated commercial apple orchards but may occur in high population densities in untreated orchards. The larvae can cause direct damage to apple flowers (Burckhardt 1994). Young adults of *C. picta* and emerging adults of *C. mali* can be found during the same period on apple and, therefore, may be mis-identified by inexperienced persons.

The presented data are of high practical relevance because *C. picta* and *C. pruni* were found to be infected by the respective phytoplasma in different regions of the study. About 10 % of the field-captured overwintered adults of *C. picta* were tested positive by PCR for '*Ca.* P. mali' and successful transmission trials suggest that a high portion of these PCR-positive specimens were indeed infectious (Jarausch *et al.* 2007b). '*Ca.* P. prunorum' infected individuals of *C. pruni* were identified in all studied regions and a natural infection rate of the overwintered adults ranging from 1–3 % was determined (Jarausch *et al.* 2007a, b). Successful transmission trials demonstrated again the infectivity of these individuals (Jarausch

et al. 2007a). Data about the host plant preference of *C. pruni* with respect to different *Prunus* species have been presented by Jarausch et al. (2007b) and confirm the data presented here: about 10-fold higher population densities were found on wild *P. spinosa* and *P. cerasifera* than on cultivated *P. armeniaca* or *P. persica*. In this regard, root stock suckers of *P. cerasifera* or *P. domestica* genotype are of particular concern because they are better colonised by *C. pruni* than the grafted cultivar of either apricot or peach. They play, therefore, an important role in the disease spread.

C. melanoneura has been identified as the most widespread species which was captured in all districts. The species can be found on many different woody plant species during its migration phases but has not been found on non-host plants during its reproduction period. Other work has shown that the population of C. melanoneura in Germany, Alsace and northern Switzerland has no capacity to transmit 'Ca. P. mali' and is, therefore, regarded as non-vector (Mayer et al. 2009). These authors also showed that the highest population density of C. melanoneura in the area of the present study is found on wild Crataegus monogyna. Thus, despite its high abundance, this species seems to represent no risk for the spread of AP in the studied area.

ACKNOWLEDGMENTS

We thank Annette Fuchs, Isabelle Lampe, Isabel Mühlenz, Uwe Harzer, Arno Fried, Elfie Schell, Hubertus Gernoth, Hubert Schneider, Hervé Bentz, Othmar Eicher, Josef Brägger, Werner Dahlbender, Günther Hensel, Wiebke Hein and Matthias Jönck for collecting psyllids. The survey was funded by Rhein-Pfalz-Wohnen GmbH (FELR), the European Community (INTERREG IIIA) and by the German Federal Ministry of Economics and Technology (ProInno II).

REFERENCES

- Bliefernicht, K. & Krczal, G. 1995. Epidemiological studies on apple proliferation disease in Southern Germany. Acta Horticulturae 386: 444–447.
- Burckhardt, D. 1994. Psylloid pests of temperate and subtropical crop and ornamental plants (Hemiptera, Psylloidea): A review. Entomology (Trends in Agricultural Science) 2: 173–186.
- Burckhardt, D. 2004. Hemiptera, Psylloidea. http://www.faunaeur.org.
- Burckhardt, D. 2007. Bestimmungsschlüssel für Psylliden auf Rosaceen in Mitteleuropa. http://www.psyllidkey.com.
- Carraro, L., Osler, R., Loi, N., Ermacora, P. & Refatti, E. 1998. Transmission of European stone fruit yellows phytoplasma by *Cacopsylla pruni*. Journal of Plant Pathology 80: 233–239.
- Čermák, V. & Lauterer, P. 2008. Overwintering of psyllids in South Moravia (Czech Republic) with respect of the vectors of the Apple Proliferation cluster phytoplasmas. Bulletin of Insectology 61: 147–148.
- Frisinghelli, C., Delaiti, L., Grando, M.S., Forti, D. & Vindimian, M.E. 2000. *Cacopsylla costalis* (Flor, 1861), as a vector of apple proliferation in Trentino. Journal of Phytopathology 148: 425–431.
- Harzer, U. 2000. Apfeltriebsucht tritt verstärkt auf. Obstbau 10: 568–569.
- Jarausch, B. 2003. Welche Rolle spielen Blattsaugerarten bei der Übertragung von Apfeltriebsucht-Phytoplasmen in deutschen Apfelanlagen? — Obstbau 4: 205–206.
- Jarausch, B., Fuchs, A., Mühlenz, I., Lampe, I., Harzer, U. & Jarausch, W. 2007a. Research on European stone fruit yellows (ESFY) in Germany. Bulletin of Insectology 60: 389–390.
- Jarausch, B., Mühlenz, I., Fuchs, A., Lampe, I., Harzer, U. & Jarausch, W. 2007b. Untersuchungen zur Europäischen Steinobstvergilbung (ESFY) in Deutschland. — Gesunde Pflanzen 59: 183–192.
- Jarausch, B., Schwind, N., Jarausch, W. & Krczal, G. 2004. Overwintering adults and springtime generation of *Cacopsylla picta* (synonym *C. costalis*) can transmit apple proliferation phytoplasmas.

 Acta Horticulturae 657: 409–413.

- Jarausch, B., Schwind, N., Jarausch, W., Krezal, G., Seemüller, E. & Dickler, E. 2003. First report of Cacopsylla picta as a vector for apple proliferation phytoplasma in Germany. — Plant Disease 87: 101.
- Mayer, C.J. & Gross, J. 2007. Different host plant odours influence migration behaviour of *Cacopsylla melanoneura* (Förster), an insect vector of the apple proliferation phytoplasma. IOBC WPRS Bulletin 30: 177–184.
- Mayer, C.J., Jarausch, B., Jarausch, W., Jelkmann, W., Vilcinskas, A. & Gross, J. 2009. *Cacopsylla melanoneura* has no relevance as vector of apple proliferation in Germany. Phytopathology 99: 729–738.
- Müther, J. & Vogt, H. 2003. Sampling methods in orchards trials: a comparison between beating and invertory sampling. *In*: Vogt, H., Heimbach, U. & Vinuela, E. (eds): Pesticides and beneficial organisms. Proceedings of the meeting at Avignon, France, 8–11 October, 2002. Dijon: INRA, Bulletin OILB srop, IOBC wprs bulletin 26: 67–72.
- Ramel, M.E. & Gugerli, P. 2004. Epidemiological survey of European stone fruit yellows phytoplasma in two orchards in western Switzerland. Acta Horticulturae 657: 459–463.
- Seemüller, E. & Schneider, B. 2004. 'Candidatus Phytoplasma mali', 'Candidatus Phytoplasma pyri' and 'Candidatus Phytoplasma prunorum', the causal agents of apple proliferation, pear decline and European stone fruit yellows, respectively. International Journal of Systematic and Evolutionary Microbiology 54: 1217–1226.
- Tedeschi, R., Bosco, D. & Alma, A. 2002: Population dynamics of *Cacopsylla melanoneura* (Homoptera: Psyllidae), a vector of apple proliferation phytoplasma in northwestern Italy. Journal of Economical Entomology 95: 544–551.

(received September 16, 2009; accepted October 29, 2009)