

**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft =  
Bulletin de la Société Entomologique Suisse = Journal of the Swiss  
Entomological Society

**Herausgeber:** Schweizerische Entomologische Gesellschaft

**Band:** 84 (2011)

**Heft:** 3-4

**Artikel:** Osellaeus bonvouloirii (Ch. Brisout, 1880) : a summary of its  
references, distribution and new insights into biology and phenology  
(Coleoptera, Apionidae)

**Autor:** Germann, Christoph / Szallies, Alexander

**DOI:** <https://doi.org/10.5169/seals-403031>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 30.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

*Osellaeus bonvouloirii* (Ch. Brisout, 1880) – a summary of its references, distribution and new insights into biology and phenology (Coleoptera, Apionidae)

CHRISTOPH GERMANN<sup>1</sup> & ALEXANDER SZALLIES<sup>2</sup>

<sup>1</sup> Natur-Museum Luzern, Kasernenplatz 6, 6003 Luzern and Naturhistorisches Museum der Bürgergemeinde Bern, Bernastrasse 15, 3005 Bern; Email: [germann.christoph@gmail.com](mailto:germann.christoph@gmail.com)

<sup>2</sup> Institut Umwelt und Natürliche Ressourcen, Zürcher Hochschule für Angewandte Wissenschaften ZHAW Campus Wädenswil, Grüental, 8820 Wädenswil; Email: [szal@zhaw.ch](mailto:szal@zhaw.ch)

We provide new observations on the biology and phenology of *Osellaeus bonvouloirii* (Ch. Brisout, 1880) *sensu lato*. Data on its distribution is summarized and discussed, and literature references are compiled.

Keywords: Coleoptera, Curculionoidea, *Osellaeus bonvouloirii*, *O. bonvouloirii baldensis*, *O. bonvouloirii occidentalis*, Alps, France, Italy, Switzerland.

#### INTRODUCTION

*Osellaeus bonvouloirii sensu lato* (s. l.) is distributed in the subalpine and alpine zones from 1400–3450 m a.s.l. in the central and western Alps from France to Switzerland and Italy. As distributional records are dispersed in the literature and original new data on distribution, taxonomy and phenology were gained, we here present a synopsis for this remarkable Alpine element.

#### MATERIAL & METHODS

Data for the distribution map is taken from references cited further down and from the following collections (including abbreviations): Naturhistorisches Museum Basel (NMBA); Naturhistorisches Museum der Bürgergemeinde Bern (NMBE); collection Eidgenössische Technische Hochschule Zürich (ETHZ); collection Alexander Szallies, Reutlingen (cAS); collection Christoph Germann, Thun (cCG).

Specimens of *O. b. bonvouloirii* were sifted from mosses and plants (mainly *Saxifraga* spp.). The specimens were extracted from the litter following Stüben *et al.* (2000).

Breeding experiments were carried out under room temperature conditions in a glass terrarium with sandy ground, where specimens of *O. b. bonvouloirii* were kept in plastic boxes (diameter: 50 mm, height: 20 mm) with absorbent paper on the ground, together with their host plants. These were exchanged every second day. Ten specimens were placed together in each of the four boxes.

## RESULTS &amp; DISCUSSION

*Taxa****Osellaeus bonvouloirii bonvouloirii* (Ch. Brisout, 1880)**

*First description:* *Apion bonvouloirii* (Brisout de Barneville 1880: 232).

*Locus typicus:* Switzerland, Bern, Briener Rothorn (see remark in Germann 2010a: 3, and/or Janetschek 1956: 503–504).

*References:* Desbrochers 1893; Schilsky 1906; Wagner 1910; Wagner 1914; Schatzmayr 1922; Hoffmann 1929a, b, 1958; Luigioni 1929; Solari 1933; Portevin 1935; Sainte-Claire Deville 1938; Janetschek 1956; Péricart 1960, 1974; Osella 1966, 1972, 1977; Focarile 1976, 1982, 1987; Dieckmann 1977; Tempère 1978; Bellò et al. 1980; Bajtenov 1980; Lohse 1981; Tempère & Péricart 1989; Alonso-Zarazaga 1990; Ehret 1990, 1992, 1997; Abbazzi & Osella 1992; Abbazzi *et al.* 1994; Behne 1994; Scherler 1995; Alonso-Zarazaga & Lyal 1999; Pedroni 2000; Colonnelli 2003; Pelletier 2005; Abbazzi & Maggini 2009, Germann 2010a, b; Alonso-Zarazaga 2011, Goggi 2011.

*Nomen dubium* (mostly considered as synonym to the nominal species): *Apion bonvouloirii georgeli* Hoffmann, 1929.

Two valid subspecies are presently known:

*Osellaeus bonvouloirii baldensis* (Bellò, Meregalli & Osella, 1980).

*First description:* *Apion bonvouloirii baldensis* (Bellò *et al.* 1980: 519–525).

*Locus typicus:* Italy, Veneto, Monte Baldo (Telegrafo).

*Osellaeus bonvouloirii occidentalis* Germann, 2010

*First description:* *Osellaeus bonvouloirii occidentalis* (Germann 2010a: 3–10)

*Locus typicus:* France, Vercors, Col de la Bataille.

*Distribution:* Distribution maps of *O. bonvouloirii* were previously provided by Janetschek (1956), Osella (1972), Focarile (1986, 1987) and Bellò *et al.* (1980). Records used to draw the distribution map given herein were taken from the selected references listed in the following. France: Péricart (1960), Germann (2010a); Italy: Osella (1966), Bellò *et al.* (1980), Pedroni (2000), Germann (2010a), Goggi (2011); Switzerland: Germann (2010a). We here provide an up-to-date overview on the distribution of *O. bonvouloirii* s. l. (Fig. 1).

The highly doubtful record of *O. b. georgeli* from Eymoutiers, not yet verified by the examination of type material (thus categorized here as *nomen dubium*) was not included.

Among the material examined, further records from Switzerland were made. These are listed below:

- 1 specimen (female) TI, Gridone, 2000 m, 28.10.2010. Remarks: black, without bluish shimmer (cAS).
- 1 specimen (sex not specified) GR, Avers-Cresta, 8.1939, leg. A. Linder (vidit P. Scherler), not found in the ETHZ.




Fig. 1. A) Map showing records of *Osellaeus bonvouloirii* (Ch. Brisout, 1880) s. l. (including an overview map B) with records of *D. alpestris* Stierlin, 1878 modified after Germann (2011), which strikingly resembles the distribution pattern of *O. bonvouloirii* s. l.

- 38 specimens (21 males, 17 females) FR, Kaiseregg, 3.11.2010, from *Saxifraga exarata* Vill. s.l. leg. A. Szallies (cAS, NMBE)
- 142 specimens (54 females and 65 males, and 23 specimens where sexes were not specified) FR, Kaiseregg, 1.4.2011, from *Saxifraga exarata* and *S. oppositifolia* L. leg. A. Szallies & C. Germann (cAS, cCG).
- 2 specimens (1 male and 1 female) NW, Brisen, 2350 m, 8.4.2011, leg. A. Szallies (cAS)

Janetschek (1956) regarded the wingless and hence quite immobile *O. bonvouloirii* as a typical glaciation relict element, supposedly having survived at least the last glaciation on mountain peaks and massifs de refuge. But another persuasive hypothesis can be put forward, complicating Janetschek's straightforward hypothesis, and not being necessarily mutually exclusive. *O. bonvouloirii* might have gained much of its recent distribution just recently in the much warmer post-glacial period following the last glaciation. Thus the hypothesized glacial refugia on nunataks of the western Alps for instance might only be most recent refugia in the wake of climate cooling. How much of *O. bonvouloirii*'s recent distribution thus might be regarded «protocratic» (see Schmölzer 1999), after the last glaciation, certainly is worth considering. Very recently Lohse *et al.* (2011, see also Schneeweiss & Schönswetter 2011)

showed, based on the *Trechus pertyi* Heer, 1837 species group (Carabidae) in the Orobian Alps, that both, survival on nunataks and recolonisation from refuge areas might have been involved in species' biogeographical history, although acting at different times.

Our updated and summarized records available from literature and collection data (Fig. 1) provide a much more complete picture of the present distribution of *O. bonvouloirii*. Seemingly, we provide support for Janetschek's hypothesis, as all records strictly follow the arc of the Alps, and the finds are restricted to ice free mountaintops at high altitudes. Furthermore, with the subspecies *baldensis* at the eastern border, and *occidentalis* at the western one, morphologically different populations are extant. These might indeed represent the suspected «relicts» of an ancient wider distribution in pre-glaciation time Janetschek (1956) hypothesized. But these might as well be the consequences of more ancient glaciations and furthermore, much of the dispersion of any subspecies, in particular the nominate species, might be most recent and thus protocratic.

The overlap of the distribution of *O. bonvouloirii* with the one of *Dichotrachelus alpestris* Stierlin, 1878, a strictly western alpine member of Curculionidae, based on the modified map from Germann (2011) is striking (Fig. 1). Although their host plants are different (*D. alpestris* lives in moss cushions, whereas *O. bonvouloirii* lives on *Saxifraga* spp.), their ecological and/or microhabitat demands are very similar, and they are likely to share the same biogeographical history during the ice age glaciations.

### Phenology

Freshly emerged specimens of *O. b. baldensis* in September (Bellò *et al.* 1980), and a fresh (immature) specimen of *O. b. bonvouloirii* from 13 August 2001 from Rochers de Naye, coll. P. Scherler (NMBE) both indicated that imagines of *O. bonvouloirii* emerge in late summer and beginning of autumn. The record of 38 specimens in November 2010 in one particular location, and the subsequently collected 142 specimens on 1st April 2011 at the same location definitely confirms that the imagines of *O. bonvouloirii* overwinter.

Péricart (1974) reported on a very short appearance time of *O. bonvouloirii* in the French Alps (Briançonnais und Queyras) with a peak in the middle of July and disappearance of the imagines already at the beginning of August «...la durée d'apparition de l'imago est très brève.» and «...l'abondance maximale a lieu vers la mi-juillet, et l'insecte disparaît après les premiers jours d'août.» Unfortunately Péricart (1974) did not indicate how many specimens his observations were based on. However, regarding the presented data here, his generalized conclusion cannot be valid.

We compiled 417 records of specimens available from literature and collection data and plotted them against the year's months (Fig. 2). Records peak in April, September and November, not being common during usual sampling times of most entomologists in the Alpine zone (which is June to August). All three peaks are mainly based on single collection events, and therefore show that previous assumptions on the rarity of *O. bonvouloirii* are mostly due to the rarefaction of specimens in summer time, when the overwintered specimens have propagated and died off and the new generation has not yet emerged.




Fig. 2. Plot of the collection dates (months 1–12) of the 417 specimens of *Osellaeus bonvouloirii* (Ch. Brisout, 1880) s. l. included here (data was taken from collections and listed references).

### Breeding experiments

40 specimens of *O. b. bonvouloirii* collected on 1st April 2011 were fed in captivity kept in a terrarium. Because indigenous *Saxifraga* spp. were not available ad libitum, we used instead a *Saxifraga x arendsii* -hybrid (cult.) (a crossbreeding from *S. hypnoides* L., *S. exarata* s. l., and *S. rosacea* Moench.), which is commonly sold in garden centres. All plant organs were offered, and controlled daily regarding feeding traces and possible oviposition. On 3rd of April, the first egg was observed in a fruit capsule. Its colour was bright yellow when freshly deposited. Its size (measured from totally 10 eggs) was about 0.3 mm, its form varied from circular to oval. Until April 12th, 12 deposited eggs could be observed: 3 in fruit capsules, 1 in crown leaf, 1 in flower bud, 1 in flower stem, 5 in leaves, 1 in a leaf basis (Fig. 3). The eggs were deposited in the following manner: first a hole was bored into the substrate, and then a cavity was hollowed out wherein the egg was deposited. Afterwards the cavity was sealed with faeces. About ten days after oviposition, the colour of the eggs changed to greyish-blackish. On April 14th the first two larvae (from the two first eggs laid in fruit capsules), of about the same length as the egg, emerged, which fed immediately on plant tissue, and immature corns provided (Fig. 3E). One of the larvae was kept alive and resetted every two days in fresh fruit capsules of *Saxifraga x arendsii* and also *S. oppositifolia* (that was now available during a certain time). The larva mostly fed in a mining manner, and grew (Fig. 4), and till beginning of Mai, its length was about 1 mm. On 10th of Mai the larva unfortunately died, because its substrate desiccated, due to absence of the first author.




Fig. 3. Oviposition sites of *Osellaeus b. bonvouloirii* (Ch. Brisout, 1880) with eggs deposited in different plant organs of *Saxifraga x arendsii* -hybrid. The arrows indicate the oviposition sites: A) in leaf basis; B) in fruit capsule (freshly deposited egg at the arrow's point); C) in leaves (the freshly deposited egg is visible through the plant tissue, and the hole is sealed with dark faeces); D) in flower stem; E) feeding first larval instar in fruit capsule (Photos: C. Germann).

A hypothesis of the generalized life cycle of *O. bonvouloirii* s. l. is proposed:

- Oviposition occurs in spring (April to May) after overwintering. Another (egg-) maturation feeding of the adults and copulating follow (directly observed). Substrates for oviposition are leaves, nodes, stems, fruits and crown leaves of *Saxifraga* spp. (presented laboratory data).
- Larval substrates are the same where eggs were deposited, the larvae feed/mine within the respective plant tissue (presented laboratory data).
- Larvae develop in late spring and during summer. Probably – depending on potential harsh climatic conditions – the mean developmental time of 40 days within Apionidae (Dieckmann 1977) might be exceeded. Overwintered adults die off (rarefaction of the species; likely the reason for low specimen numbers of *O. bonvouloirii* in collections).
- Pupal development (most probably within the feeding substrate, and not in the ground) is completed in August and September, adults emerge, feed – observations by Bellò *et al.* (1980) – and overwinter (as stated here).




Fig. 4. A) Traces of a mining larva of *Osellaeus b. bonvouloirii* (Ch. Brisout, 1880) in a flower bottom of a *Saxifraga arendsii* -hybrid (view from the outside, see arrow); B and C) older (probably second) larval instar at 6th of Mai (Photos: C. Germann).

### Host-plants

As possible host-plant of *O. bonvouloirii*, *Saxifraga oppositifolia* is predominantly reported. Also *O. b. occidentalis* was found on this plant (Germann 2010b). Osella (1972) reported *Saxifraga biflora* All. as a potential host plant, since specimens were found nearby under stones. Péricart (1974) was the first who reported *S. caesia* as another possible host-plant for *O. bonvouloirii* in the Briançonnais «... aussi sur une espèce du groupe de *caesia* L.». Scherler (1995, and unpublished records in NMBE) reported finds from both plants. *O. b. baldensis* was found on *S. caesia*; freshly emerged adults were feeding on the leaves and stems, and also emerged from collected plant cushions (Bellò *et al.* 1980).

Solari (1933) reported on adults sifted under *Saxifraga oppositifolia* showing deformations on the elytra. He concluded these deformations to be due to movements of the ground, wherein he supposed the embedded pupal stage. According to Dieckmann (1977), nymphosis to take place in the ground would be rather exceptional within Apionidae. Because he did not find feeding traces or larval stages of *O. bonvouloirii*, Solari (1933) assumed that larvae might feed ectophagously on the plant roots, something we would dismiss regarding our here presented results.

We found about 50 specimens (of totally 142) of *O. b. bonvouloirii* on the Kaiseregg in a scree slope where exclusively *Saxifraga exarata* grew. Feeding tests showed that *S. exarata* is very likely another host plant for *O. b. bonvouloirii*, apart from another finding location at the same mountain (just 100 m away) where *O. b. bonvouloirii* was collected near *S. oppositifolia*. Furthermore, as *O. b. bonvouloirii* also fed (and laid eggs) on a *Saxifraga x arendsii* -hybrid, other *Saxifraga* spp. can also be expected as host plants. However, further study is needed to clear this unambiguously.

## ACKNOWLEDGEMENTS

For access to the collections we thank Andreas Müller, Franziska Schmid (both ETHZ), Eva Sprecher, Michel Brancucci (both NMBA) and Charles Huber (NMBE). We thank Lutz Behne (Deutsches Entomologisches Institut Müncheberg, Senckenberg) for literature provided, and Henryk Luka (Forschungsinstitut für biologischen Landbau (FiBL), Frick, and Universität Basel, Departement Umweltwissenschaften, NLU-Biogeographie) for checking the manuscript. We acknowledge financial and other support from the Bundesamt für Umwelt (BAFU), and in particular from Francis Cordillot to the ZHAW and the co-author.

## REFERENCES

- Abbazzi, P., Colonnelli, E., Masutti, L. & Osella, P. 1994. Coleoptera Polyphaga XVI (Curculionidae). *In*: Minelli, A., Ruffo, S. & La Posta, S. (eds). Checklist delle specie della fauna italiana 61. — Calderini Bologna. 68 pp.
- Abbazzi, P. & Maggini, L. 2009. Elenco sistematico-faunistico dei Curculionidae italiani, Scolytidae e Platypodidae esclusi (Insecta, Coleoptera). — *Aldrovandia* 5: 29–216.
- Abbazzi, P. & Osella, G. 1992. Elenco sistematico-faunistico degli Anthribidae, Rhinomaceridae, Attelabidae, Apionidae, Brentidae, Curculionidae italiani (Insecta, Coleoptera, Curculionidae). — *Redia* (Firenze) 75 (2): 267–414.
- Alonso-Zarazaga, M.A. 1990. Revision of the supraspecific taxa in the Palaearctic Apionidae Schoenherr, 1823 (Coleoptera, Curculionidae). 2. Subfamily Apioninae Schoenherr, 1823: introduction, keys and descriptions. — *Graellsia* 46: 19–156.
- Alonso-Zarazaga, M.A. 2011. Apionidae p. 79. *In*: Löbl, I. & Smetana, A. (eds) Catalogue of Palaearctic Coleoptera, Vol. 7. — Stenstrup, Apollo Books. 373 pp.
- Alonso-Zarazaga, M.A. & Lyal, C.H.C. 1999. A World Catalogue of Families and Genera of Curculionidae (Insecta: Coleoptera) (Excepting Scolytidae and Platypodidae). — *Entomopraxis*, Barcelona. 315 pp.
- Bajtenov, M.S. 1980. Notizen über Arten der Untergattung *Synapion* Schilsky der Gattung *Apion* Herbst (Coleoptera, Curculionidae). — *Reichenbachia*, Dresden 18 (5): 49–50.
- Behne, L. 1994. Superfamilie Curculionidae. *In*: Lohse, G.A & Lucht, W.H. (eds). Die Käfer Mitteleuropas. 3. Supplementband mit Katalogteil. — Goecke & Evers, Krefeld. Band 14: 181–246.
- Bellò, C., Meregalli, M. & Osella, G. 1980. Una nuova sottospecie di *Apion bonvouloiri* Bris. del Monte Baldo (Coleoptera: Curculionidae: Apionidae). — *Bolletino del Museo Civico di Storia Naturale*, Verona 7: 519–525.
- Brisout de Barneville, Ch. 1880. Descriptions de Coléoptères nouveaux d'Europe. — *Annales de la Société entomologique de France*, p. 232.
- Colonnelli, E. 2003. A revised checklist of Italian Curculionidae (Coleoptera). — *Zootaxa* 337: 142 pp.
- Desbrochers, J. 1893. Révision des espèces de Curculionides appartenant à la tribu des Gymnetridae d'Europe et circa. — *Le Frelon* 3 (1–2): 37–68.
- Dieckmann, L. 1977. Beiträge zur Insektenfauna der DDR: Coleoptera – Curculionidae (Apionidae). — *Beiträge zur Entomologie*, Berlin 27: 7–143.
- Ehret, J.-M. 1990. Les Apions de France. Clés d'identification commentées (Coleoptera, Curculionidae, Apioninae). — *Bulletin mensuel de la Société linnéenne de Lyon* 59 (7): 209–292.
- Ehret, J.-M. 1992. Les Apions de France. (Coleoptera Curculionidae) : Notes complémentaires et corrigenda. — *Bulletin mensuel de la Société linnéenne de Lyon* 61 (6): 180–192.
- Ehret, J.-M. 1997. Essai de classification des Apions paléarctiques (Coleoptera Curculionidae: Apioninae) en fonction des familles végétales d'Angiospermes Dicotylédones parasitées. — *Nouvelle Revue d'Entomologie* (N.S.), Paris 13 (3): 191–221.

- Focarile, A. 1976. Sulla Coleotterofauna alticola della conca del Breuil (Valtournanche) e osservazioni sul popolamento pioniero delle zone di recente abbandono glaciale. — *Revue Valdôtaine d'Histoire Naturelle* (Aosta) 30: 126–168.
- Focarile, A. 1982. La coleotterofauna geobia del Monte Generoso (Ticino, Svizzera) nei suoi aspetti ecologici, cenotici e zoogeografici. — *Bollettino della Società Ticinese di Scienze Naturali* 70: 15–62.
- Focarile, A. 1987. *Ecologie et biogéographie des coléoptères de haute altitude en Vallée d'Aoste*. — *Arti Grafiche Duc*, Aosta. 90 pp.
- Germann, C. 2010a. *Osellaeus bonvouloiri occidentalis* ssp. nov. from the Vercors (France) – an alpine element at its distributional limits (Coleoptera, Apionidae). — *Revue suisse de Zoologie*, 117 (1): 3–10.
- Germann, C. 2010b. Die Rüsselkäfer der Schweiz – Checkliste (Coleoptera, Curculionoidea) mit Verbreitungsangaben nach biogeografischen Regionen. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 83: 41–118.
- Germann, C. 2011. Review of the *Dichotrachelus alpestris* Stierlin, 1878 species group with evidence for a species complex of *D. augusti* F. Solari, 1946, and *D. sondereggeri* sp. nov. from Switzerland (Coleoptera, Curculionidae). — *Contributions to Natural History* 17: 1–21.
- Goggi, G. 2011. Indagine faunistica sui Coleotteri Scarabeidi e Curculionidi (Coleoptera: Scarabaeoidea, Curculionoidea) della Valsessina (Lecco, Lombardia). — *Giornale Italiano di Entomologia* 56 (12): 355–371.
- Hoffmann, A. 1929a. Liste d'insectes Rhynchophores capturés dans le département de la Haute-Vienne, accompagnée d'observations diverses. — *Miscellanea Entomologica* 32: 3–12.
- Hoffmann, A. 1929b. Une nouvelle variété du genre *Apion*: (*Perapion*) *Bonvouloiri* Ch. Bris., var. *Georgeli* n. var. (Col. Curculionidae). — *Miscellanea Entomologica* 32: 43–44.
- Hoffmann, A. 1958. Faune de France, No. 62. Coléoptères Curculionides. — *Editions Paul Lechevalier*, Paris. 3ième partie: 1210–1839.
- Janetschek, H. 1956. Das Problem der inneralpinen Eiszeitüberdauerung durch Tiere. — *Österreichische Zoologische Zeitschrift*, Wien 6: 421–506.
- Lohse, G.A. 1981. 5. Unterfamilie: Apionidae 127–183. In: Freude, H., Harde, K.W. & Lohse, G.A. (eds), *Die Käfer Mitteleuropas*. Band 10. — Goecke & Evers, Krefeld. 310 pp.
- Lohse, K.L., Nicholls, J.A. & Stone, G.N. 2011. Inferring the colonization of a mountain range – refugia vs. nunatak survival in high alpine ground beetles. — *Molecular Ecology* 20: 394–408.
- Luigioni, P. 1929. I Coleotteri d'Italia. Catalogo sinonimico-topografico-bibliografico. — *Memorie della Pontificia Accademia delle Scienze Nuovi Lincei, Serie III* 13: 1–1160.
- Osella, G. 1966. Alcuni interessanti reperti di Curculionidi italiani ed osservazioni sulla loro corologia. — *Memorie del Museo Civico di Storia Naturale, Verona* 14: 389–399.
- Osella, G. 1972. Notulae Curculionidologiche. II. (Coleoptera, Curculionidae). — *Bollettino dell'Associazione Romana di Entomologia* 27 (3–4): 53–62.
- Osella, G. 1977. La Curculionidofauna dei pascoli d'alta quota d'Alpi ed Apennino. — *Animalia*, 4 (3): 223–276.
- Pedroni, G. 2000. Secondo contributo alla conoscenza della Curculionidofauna della Valle di La Thuile (Valle d'Aosta, Italia settentrionale) (Insecta, Coleoptera, Apionidae, Curculionidae). — *Il naturalista valtellinese* 11: 27–38.
- Pelletier, J. 2005. Catalogue des Curculionoidea de France (Coleoptera). — *Biocosme Mésogéen*, Nice 21 (3): 75–147.
- Péricart, J. 1960. Notes systématiques relatives à divers Coléoptères Curculionides. — *Bulletin de la Société Entomologique de France* 65: 161–170.
- Péricart, J. 1974. Notes sur divers Curculionidae français avec la description d'une espèce et d'une sous-espèce nouvelles (Coleoptera). — *Nouvelle Revue d'Entomologie* 4 (1): 55–70.
- Portevin, G. 1935. *Histoire Naturelle des Coléoptères de France*. Tome IV Polyphaga : Rhynchophora. — *Paul Lechevalier*, Paris, 500 pp.
- Sainte-Claire Deville, J. 1938. *Catalogue raisonné des Coléoptères de France*. — *Firmin-Didot et Cie Mesnil*, Eure. 466 pp.
- Schatzmayr, A. 1922. Gli Apionini Italiani. — *Memorie della Società Entomologica Italiana* 1: 158–225.
- Scherler, P. 1995. Répartition actuellement connue en Suisse de quelques Coléoptères Curculionides d'altitude. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 68: 179–187.
- Schilsky, J. In: Küster, H.C. & Kraatz, G. 1906 (eds). *Die Käfer Europas* 42: 1–42.
- Schmölzer, K. 1999. Prä- und interglaziale Elemente in der Acrofauna der Alpen. — *Carinthia* II 109: 573–602.
- Schneeweiss, G.M. & Schönswetter, P. 2011. Perspective. A re-appraisal of nunatak survival in arctic-alpine phylogeography. — *Molecular Ecology* 20: 190–192.

- Solari, F. 1933. Notizie Biologiche sull' *Apion bonvouloiri* Bris. e sull' *Dichotrachelus sulcipennis* Stierl. (Col. Curc.). — Bollettino della Società Entomologica Italiana 65: 159–161.
- Stüben, P.E., Bahr, F. & Bayer, Ch. 2000. Cryptorhynchinae-Exkursion in die Ost- und Zentralpyrenäen (Coleoptera: Cryptorhynchinae) — Weevil News: No.1: 10 pp.
- Tempère, G. 1978. Catalogue des Coléoptères Curculionidae de France. Essai de mise à jour critique. Sixième partie. — Entomops 46: 213–232.
- Tempère, G. & Péricart, J. 1989. Faune de France 74. Coléoptères Curculionidae 4ième Partie. — Fédération française des sociétés de sciences naturelles, Paris. 534 pp.
- Wagner, H. 1910. Curculionidae, Apioninae. In: Junk, W. & Schenkling, S. (eds), 1910–1939. — Coleopterorum Catalogus. Vol. 28. Curculionidae II. Pars 6: 1–81.
- Wagner, H. 1914. Beiträge zur Kenntnis der Gattung *Apion* Herbst. V. — Münchener Koleopterologische Zeitschrift 4: 12–32.

(received May 18, 2011; accepted November 28, 2011)