

**Zeitschrift:** Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

**Herausgeber:** Schweizerische Entomologische Gesellschaft

**Band:** 85 (2012)

**Heft:** 3-4

**Artikel:** Nebria (Patrobonebria) megalops sp. n. from Yunnan (China)  
(Coleoptera, Carabidae)

**Autor:** Huber, Charles / Geiser, Michael

**DOI:** <https://doi.org/10.5169/seals-403045>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 05.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

*Nebria (Patrobonebria) megalops* sp. n. from Yunnan (China)  
(Coleoptera, Carabidae)

CHARLES HUBER<sup>1</sup> & MICHAEL GEISER<sup>2</sup>

<sup>1</sup> Naturhistorisches Museum der Burgergemeinde Bern, Bernastrasse 15, CH-3005 Bern, Switzerland;  
charles.huber@nmbe.ch

<sup>2</sup> Michael Geiser, Naturhistorisches Museum, Augustinergasse 2, CH-4001, Basel, Switzerland and  
University of Basel, Department of Environmental Sciences, Biogeography, St. Johanns-Vor-  
stadt 10, CH-4056, Basel, Switzerland; michael.geiser@stud.unibas.ch

*Nebria (Patrobonebria) megalops* sp. n. from Cang Shan near Dali/Yunnan (China) is described. The new species is related to *N. (Patrobonebria) desgodinsi* Oberthür, 1883, and is the species with the easternmost distribution in the Himalayan range of the subgenus.

Keywords: Coleoptera, Carabidae, *Nebria*, *Patrobonebria*, taxonomy, new species, Yunnan, China.

## INTRODUCTION

The subgenus *Patrobonebria* Bänninger, 1923 is a small group of only six species of the genus *Nebria* Latreille, 1802 comprising *N. desgodinsi* Oberthür, 1883, *N. himalayica* Bates, 1889, *N. elegans* Andrewes, 1925, *N. capillosa* Ledoux & Roux, 1992, *N. pertinax* Huber & Schmidt, 2009, and *N. assidua* Huber & Schmidt, 2009. All species are restricted to the Himalayan range sensu stricto covering an area from Afghanistan to Sikkim.

Huber & Schmidt (2009) gave new records of *N. desgodinsi* Oberthür, 1883, which is known up to now to be the *Patrobonebria* species with the easternmost distribution (Eastern Nepal, Sikkim). Newly collected specimens by the junior author on the Cang Shan near Dali/Yunnan (China) turned out to be a new *Patrobonebria* species outside of the known distribution area of the subgenus. The new species is described below.

## MATERIAL AND METHODS

The type series was collected at the southern slope of Cang Shan, a chain of the Yunlin Mountains on the southern fringes of the Hengduan Mountains system, which for its part forms the eastern extent of the Himalayan range sensu lato. Although the eastern slope of Cang Shan near Dali city has been visited by naturalists since the 19th century, hardly any entomologists have ever visited the southern slope.

### Abbreviations and material depository:

NMB: Natural History Museum Basel, Switzerland

NMBE: Natural History Museum Bern, Switzerland

IZAS: Institute of Zoology, Chinese Academy of Sciences, Beijing, China



Fig 1: Head of *Nebria (Patrobonebria) megalops* sp. n. Scale bar: 1 mm.

The photographs were made with a digital camera Leica DFC 425, and the images were processed by the automated multifocus software (Imagic Image Access, Version 10).

#### DESCRIPTION

##### *Nebria (Patrobonebria) megalops* sp. n.

*Material examined.* Holotype male: China, Yunnan prov., Dali, Cang Shan, Da Po Qing vill., 2000–2200 m, 25° 34' N, 100° 08' E, 15. 4. 2011, Michael Geiser leg. (NMB).— Paratypes: 3 males, 4 females, same data as holotype (NMB, NMBe, IZAS).

*Diagnosis.* By reason of the collar impression of the head, the bisetose penultimate labial palpomere, and the dilated protarsomeres 1–2 in the male, *N. megalops* sp. n. belongs of the subgenus *Patrobonebria* defined by Bänninger (1923).

*Description.* Body length: 13.5–15 mm. Colour black, mandibulae und appendages of the head brown. Head usually with two small brown spots on the vertex. Antennae brown, the four basal antennomeres lightened brown. Legs black, except tibia and tarsus lightened brown.

Head large, with a distinct transverse collar impression of the neck behind the eyes (fig 1). Mandibles conspicuously long. Labrum with anterior margin distinctly trisinuate, medially deeply notched, bearing 6 setae. Apical margin of clypeus slightly concave. Supraorbital impression distinct, obliquely wrinkled. Head near supraorbital seta longitudinally wrinkled. Vertex and collar impression distinctly



Fig 2: Pronotum of *Nebria (Patrobonebria) megalops* sp. n. lip: lateral impression of the pronotum; blip: basolateral impression of the pronotum. Scale bar: 1 mm.

punctate. Hemispheric eyes very prominent. Temples long, regularly rounded towards the neck. Antennae long and slender, extending to the middle of the elytra. Antennal scape subcylindrical, basally narrowed, shorter than the eye's diameter, with 1 dorsal seta (rarely bisetose). 4th antennomere only with an apical collar of long setae and only few additional short setae. Ligula blunt with 2 long setae apically. Maxillary stipes basolaterally with 5–7 robust setae. Penultimate labial palpomere bisetose. Mentum with bifid medial tooth. Submentum with a row of 4–8 setae. Microsculpture of the head isodiametric.

Pronotum cordate (fig 2), transverse, widest at apical fifth; ratio width/length of the pronotum 1.39 (1.34–1.42). Lateral margin convex, with a distinct concave lateral impression at the insertion site of the midlateral seta deforming the curve of the lateral margin. Pronotum narrowed towards the anterior angles, strongly and almost in a straight line narrowed towards the posterior angles, concavely and sharply angled before the posterior angles. Posterior angles acute, slightly protruding outwards. Posterior angles distinctly narrower than the anterior ones (0.78 (0.76–0.81)). Lateral groove narrow, narrowest at the insertion site of the midlateral seta, apically distinctly broadened. Anterior angles large, of triangular shape, protruding. Basal margin bisinuate. Pronotal disc convex, transversally wrinkled. Basal fovea deep, not reaching the basal margin. Anterior and posterior transverse impressions deep, median longitudinal impression weak. Basal fovea, the anterior and the posterior transverse impressions and the narrow lateral groove tightly and coarsely punctate.




Fig 3. Median lobe of *Nebria (Patrobonebria) megalops* sp. n. Scale bar: 1 mm.

Apical and basal margination absent. 1 midlateral and 1 basolateral seta present. Microsculpture of the pronotum isodiametric. Pronotal disc faintly and sparsely punctate. Proepisternum punctate. Prosternal process broad, triangular, longitudinally bulged, unmargined.

Elytral silhouette parallel-elongate. Lateral margin at the basal fourth with a faint concave impression, subapically faintly sinuate. Elytral apex rounded. Basal margination curved, joined at an obtuse angle with the lateral margination. Humeral carina faint. Striae distinct, distinctly punctate. Striae 1–3 reaching the apex, the other striae obliterated just before the apex. Intervals faintly convex, interval 3 with 6–8 setae. Scutellar seta present. Microsculpture isodiametric. Mesepisterna tightly punctate. Metepisterna 2.5 times as long as wide, coarsely and tightly punctate. Metacoxa with 3–4 basal and 1 apical setae. Hindwings fully developed. 2nd sternum (sensu Ledoux & Roux 2005) laterally rough, coarsely punctate. 3rd abdominal sternum medially asetose. Sterna 4–6 each with 2–3 posterior paramedial setae. Anal sternum with 1 paramedial seta in the male and 2 in the female. All sterna with faint impressions laterally.

Legs long and slender. All tarsi without a dorsal pubescence. Protarsus of the male with tarsomeres 1–2 dilated, ventrally with pads of adhesive setae. Metatarsomere 4 with a short ventroapical tooth, bearing long setae. Metatarsomere 5 slender, ventrally with 2 rows of only 1–2 short setae. Metatarsomere 5 shorter than the metatarsomeres 3+4. Metatarsomeres 2–4 laterally compressed.

Median lobe (fig 3): Basal part of the median lobe large. Mid-shaft slender, narrowed directly after the basal part, slightly curved to the apex. Apex slender, acute, faintly deflected towards the left.

*Etymology:* The specific epithet refers to the distinctly protruding hemispheric eyes.

*Affiliation.* Due to the parallel sided elytra and the faintly punctate vertex, *N. megalops* sp. n. belongs to the *N. desgodinsi* group (*N. desgodinsi* Oberthür, 1883 and *N. capillosa* Ledoux & Roux, 1992). *N. megalops* sp. n. is closely related to *N.*

*desgodinsi* from which it is separated by the distinct concave impression of the pronotum at the insertion site of the midlateral seta, by the narrow lateral groove of the pronotum, and by the blackish legs. *N. capillosa* differs from *N. megalops* sp. n. and from *N. desgodinsi* by the polypilose apex of antennomere 4 and by the slender pronotum.

*Ecology.* *N. megalops* sp. n. is a hydrophilic, riparian species.

*Collecting circumstances.* The vicinity of Da Po Qing village, at slightly over 2000 m a. s. l., is mainly covered by dry *Pinus* forests and irrigated, small-scale agricultural fields. All specimens of the new species were found under stones directly beneath a waterfall (see fig 4), accompanied by other Carabidae species (*Bembidion eurydice* Andrewes, 1926; *Onycholabis* sp.) and large numbers of *Dianous* sp. (Staphylinidae). Further searching along the course of the same stream remained unsuccessful, which may indicate that *N. megalops* sp. n. is restricted to these extremely wet microhabitats.

*Distribution.* *N. megalops* sp. n. is known from the *locus typicus* only.

The new species is the species with the south-easternmost distribution of the subgenus *Patrobonebria*. There is a large distributional gap of more than a thousand kilometers between the area of *N. desgodinsi* in Eastern Nepal and Sikkim and *N. megalops* sp. n. in Yunnan. Due to the fully developed hind wings, a wide distribution area of *N. megalops* sp. n. is to be expected, as it is the case in *N. himalayica*, *N. elegans* and *N. capillosa* (Ledoux & Roux 2005).

*Remarks.* The lateral concave impression of the pronotum is very pronounced in *N. megalops* sp. n. (Fig. 2), less pronounced in *N. himalayica*, *N. capillosa* and *N. elegans* (Ledoux & Roux 2005), and only covertly present in *N. desgodinsi* and *N. assidua* (see fig. 2 in Huber & Schmidt 2009). This deformation of the marginal curve is a unique and synapomorphic character of this species group. The lateral pronotal deformation is absent only in *N. pertinax*.

Ledoux & Roux (2005, p. 616) mentioned a unique and immature (and therefore hardly determinable) specimen similar to *N. desgodinsi* from Daxue Shan/Gongga Shan (Sichuan), which is north of the area of *N. megalops* sp. n. and which is situated right in the middle of the large distributional gap along the mountain ranges between these two species. This single record indicates that further populations or even taxa of *Patrobonebria* are to be expected in the south-eastern extensions of the Himalayas sensu lato in the Chinese provinces of Sichuan and Yunnan.

The record of *N. megalops* sp. n. seriously enlarges the distribution area of the subgenus *Patrobonebria* and proves the occurrence of the subgenus in the whole Himalayan range sensu lato from Afghanistan to Yunnan/China.



Fig 4: Habitat of *Nebria (Patrobonebria) megalops* sp. n. near Da Po Qing village near Dali/Yunnan (China). Photo: M. Geiser.

#### KEY

The steps 2 and 4 of the determination key in Huber & Schmidt (2009) have to be modified:

- 2 Legs and appendices of the head yellow, distinctly brighter than the body . . . 3
- Legs and appendices of the head brown to black, scarcely brighter than the body . . . 4
- 4 Outline slender. Shoulder less distinct, elytra ovoid-elongate. Kashmir to Kumaon Himalaya . . . . . *N. elegans* Andrewes, 1925
- Outline robust. Shoulder distinctly produced, elytra parallel or subparallel . . . 4a
- 4a Pronotum flat. Lateral groove of the pronotum large, lateral margin evenly rounded. Afghanistan to Central Nepal . . . . . *N. himalayica* Bates, 1889
- Pronotum strongly convex. Lateral groove of the pronotum narrow, lateral margin with a distinct concave impression at the insertion site of the mid-lateral seta. Cang Shan/Yunnan . . . . . *N. megalops* sp. n.

#### ACKNOWLEDGEMENTS

We gratefully acknowledge the help of Dr. Yu-Xia Yang (Hebei University, Baoding) with organizing the trip to Yunnan and enabling the discovery of this new species, the help of Prof. Zi-Zhong Yang (Dali University) and Prof. Xing-Ke Yang (IZAS) with the necessary permits. We are very grateful to Dr. Christoph Germann (NMBE) for the help with the photographs, to Eva Feltkamp (Frankfurt D) with the distribution map, and to Elsa Obrecht (NMBE) with the manuscript.




Fig 5: Locus typicus (Cang Shan) of *Nebria (Patrobonebria) megalops* sp. n. near Dali and near Erhai lake/Yunnan (China). Map design: Eva Feltkamp.

#### ZUSAMMENFASSUNG

*Nebria (Patrobonebria) megalops* sp. n. aus dem Cang Shan bei Dali/Yunnan (China) wird beschrieben. Die neue Art steht *N. desgodinsi* Oberthür, 1883 nahe und ist diejenige *Patrobonebria*-Art mit der östlichsten Verbreitung im Himalaya.

#### REFERENCES

- Bänninger, M. 1923. Versuch einer Bestimmungstabelle der zentral- und ostasiatischen *Nebria*-Arten ohne gelbe Flügeldeckenzeichnung, nebst Bemerkungen über einige andere Formen (6. Beitrag zur Kenntnis der Carabinae) – Koleopterologische Rundschau 10: 129–142.
- Huber, C. & Schmidt, J. 2009. Contribution to the knowledge of *Nebria* Latreille, 1802 (*Patrobonebria* Bänninger, 1923) from the Nepal Himalaya (Coleoptera, Carabidae, Nebriinae) – Biodiversität und Naturausstattung im Himalaya 3: 115–120.
- Ledoux, G. & Roux, P. 1992. *Nebria (Patrobonebria) capillosa* nouvelle espèce himalayenne (Col. Nebriidae) – Nouvelle Revue d'Entomologie (N.S.) 9: 142.
- Ledoux, G. & Roux, P. 2005. *Nebria*. – Chirat, Saint-Just-la-Pendue, 976 pp.

(received September 20, 2012; accepted November 6, 2012; published December 31, 2012)

