

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 86 (2013)

Heft: 3-4

Artikel: Entomophthorales (Fungi, Entomophthoromycota) attacking Coleoptera
with a key for their identification

Autor: Keller, Siegfried

DOI: <https://doi.org/10.5169/seals-403074>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 03.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Entomophthorales (Fungi, Entomophthoromycota) attacking Coleoptera with a key for their identification

SIEGFRIED KELLER

Rheinweg 14, CH-8264 Eschenz; siegfriedke@bluewin.ch

A key to 30 species of entomophthoralean fungi is provided. These 30 species are listed in an alphabetical order and characterized based on their original descriptions and amended with new data. A few incompletely described but probably new species are included as well as a few unclassified species mentioned as pathogens of Coleoptera. A list of the coleopteran families containing hosts of Entomophthorales is given to help with the identification.

Key words: Insect pathogenic fungi, Entomophthorales, species list, identification key, Coleoptera.

INTRODUCTION

The systematics of the arthropod-pathogenic fungi previously placed in the order Entomophthorales has dramatically changed since DNA-based data are applied (Humber 2012). A new phylum, Entomophthoromycota, has been erected. The family Neozygitaceae became separated from the order Entomophthorales and placed in the new class Neozygitomycetes, order Neozygitales. The other arthropod-pathogenic species remained in the order Entomophthorales, placed in the newly erected class Entomophthoromycetes (Humber 2012).

The last census of the arthropod-pathogenic Entomophthoromycota revealed 236 species (Keller 2008). Since then a few further species have been added. There is no key for the identification of all these species, but there are keys to identify *Neozygites* spp. and *Entomophthora* spp. (Keller 1997, 2002) as well as keys to identify species attacking aphids and brachycerous Diptera (Keller 2006, 2007a). Keys focusing on host taxa have shown to be better suited for practical purposes and help to improve the collaboration between insect mycologists and entomologists. The following key for the identification of entomophthoralean fungi attacking Coleoptera is a further step in this direction and gives interested scientists and especially coleopterologists the possibility to identify these fungi.

METHODS

The dichotomous key is based on original descriptions except *Conidiobolus* spp. (originally described from detritus) and the widespread *Zoophthora radicans* whose original description is very poor and has as type host *Pieris brassicae* (Lepidoptera). The key is followed by a thorough description of the species listed in alphabetical order. The bold numbers in brackets refer to the number of the species in this list.

An important microscopic criterium to separate *Conidiobolus* spp. (family Ancylistaceae) from Entomophthoraceae is the staining ability of the nuclei with lactophenol-aceto-orcein (LPAO).

Generally, species are described from fresh material or from material stored for a short term in ethanol. Occasionally, species are described from air-dried (herbarium) material, so-called exsiccatae. In this case we must be aware that air-drying leads to a shrinking of the conidia and probably also of other structures except the resting spores. Since most species of Entomophthorales are very host specific, the knowledge of the host insect is very important and represents a significant role in the key.

RESULTS

Most species attacking Coleoptera produce conidia as well as resting spores, either as zygosporangia or as azygosporangia. Some species produce both spore types in the same host individual, others produce only one spore type in an individual host insect. A few species are known only in their conidial state and others only or predominantly in their resting spore state like species of *Tarichium* but also *Zoophthora rhagonycharum* and *Z. crassitunicata*. This fact is considered in the first step of the key. A branch leads to species known only or primarily in their resting spore stage while the other branch leads to species known only or predominantly in their conidial stage.

The key based on host genera as well as the key based on fungus characteristics refer to a species list which is arranged in alphabetical order and numbered consecutively.

KEY BASED ON HOST FAMILIES

In the following paragraph the coleopteran families which contains species attacked by Entomophthorales are listed alphabetically. The number in brackets after the fungus name refers to the species list given after the identification key.

Cantharidae:	<i>Entomophthora</i> sp. (6), <i>Eryniopsis lampyridarum</i> (8), <i>Pandora lipai</i> (11), <i>Zoophthora crassitunicata</i> (22), <i>Zoophthora larvivora</i> (24), <i>Zoophthora rhagonycharum</i> (27), <i>Zoophthora</i> sp. 1 (29).
Carabidae:	<i>Erynia nebriae</i> (7), <i>Furia zabri</i> (9), <i>Tarichium jaczewskii</i> (18).
Cerambycidae:	<i>Batkoa apiculata</i> (1).
Chrysomelidae:	<i>Batkoa apiculata</i> (1), <i>Pandora suturalis</i> (14), <i>Tarichium</i> sp. (unclassified)
Curculionidae:	<i>Conidiobolus coronatus</i> (3), <i>Conidiobolus osmodes</i> (4), <i>Pandora phyllobii</i> (13), <i>Tarichium cleoni</i> (15), <i>Tarichium hylobii</i> (17), <i>Tarichium phytonomi</i> (19), <i>Tarichium punctatum</i> (20), <i>Zoophthora phytonomi</i> (25).
Elateridae:	<i>Zoophthora anglica</i> (21), <i>Zoophthora elateridiphaga</i> (23).
Lagriidae:	<i>Entomophaga lagriae</i> (5).
Melolonthidae:	<i>Pandora brahminae</i> (10).
Ptilodactylidae:	<i>Batkoa major</i> (2).

Scirtidae:	<i>Zoophthora</i> sp. 2 (30).
Staphylinidae:	<i>Pandora philonthi</i> (12), <i>Zoophthora tachypori</i> (28).

KEY TO FUNGUS SPECIES

- 1 Only resting spores known or present. Conidia unknown or rare. 24
- 1* Conidia present. Resting spores may be present as well 2
- 2 Conidiophores simple. Conidia multinucleate 3
- 2* Conidiophores branched. Conidia mononucleate 9
- 3 Body of primary conidia spherical to subspherical, papilla prominent. Nuclei smaller than 3 μm , stain not or only faintly in LPAO. Rhizoids absent. Hyphal bodies complex. Good growth in artificial media . . . genus *Conidiobolus* 4
- 3* Primary conidia spherical to pyriform or with pointed apex. Nuclei larger than 3 μm , stain deeply in LPAO. Rhizoids present or absent. Hyphal bodies spherical to elliptical or irregularly rounded 5
- 4 Primary conidia on average 50.5–53 x 38–40 μm , varying largely in size, L/D = 1.30–1.34; papilla abruptly protruding, prominent, elongate. Resting spores are azygospores, on average 30.5–31.5 μm , villose, spherical *C. coronatus* (3)
- 4* Primary conidia 25–37 x 22–30 μm ; papilla broad, obtusely tapered. Resting spores are zygosporangia, spherical to ellipsoid with a diameter of 13–37 μm *C. osmodes* (4)
- 5 Primary conidia with pointed apex and truncate papilla *Entomophthora* sp. (6)
- 5* Primary conidia spherical to pyriform, prominent conical papilla. Rhizoids present or absent. Secondary conidia like primary or long-cylindrical, and born vertically on capillary conidiophores 6
- 6 Body of primary conidia nearly spherical, papilla demarcated, pointed or rounded genus *Batkoa* 7
- 6* Primary conidia pyriform to ovoid, papilla smoothly joining conidial body or with indistinct papilla 8
- 7 Primary conidia 30–37 x 28–30 μm , average 35 x 30 μm , papilla pointed *B. apiculata* (1)
- 7* Primary conidia 55–60 x 38–45 μm *B. major* (2)
- 8 Primary conidia 35–43 x 24–30 μm , ovoid or broadly turbinate with 12–18 nuclei, papilla narrow conical. Secondary conidia like primary *Entomophaga lagriae* (5)
- 8* Conidia 20–37 x 14–30 μm , average 35 x 15 μm , less than 10 nuclei. Secondary conidia like the primary or long-cylindrical and born vertically on capillary conidiophores *Eryniopsis lampyridarum* (8)
- 9 Rhizoids compound (pseudorhizomorph). Primary conidia elongate, cylindrical, subcylindrical or ellipsoid. Two types of secondary conidia: like primary or capilliconidia genus *Zoophthora* 16

9*	Rhizoids simple (monohyphal). Primary conidia ovoid to pyriform. Two types of secondary conidia: like primary or subspherical to ovoid	10
10	Pathogen of Carabidae	11
10*	Pathogen of other Coleoptera	12
11	Primary conidia ovoid, on average 25 x 14 μm . Zygosporangia spherical, formed mostly on the surface of the host. Pathogen of larval <i>Zabrus</i>	<i>Furia zabri</i> (9)
11*	Conidia ellipsoidal to fusiform, asymmetrical, 28–37 x 10–13 μm . Resting spores hyaline to pale brown, developing outside the body of the host, spherical, 35–50 μm diameter. Pathogen of <i>Nebria</i>	<i>Erynia nebriae</i> (7)
12	Pathogen of Melolonthidae. Primary conidia 18–22 x 11–15 μm	<i>Pandora brahminae</i> (10)
12*	Pathogen of other Coleoptera	13
13	Pathogen of Cantharidae. Primary conidia 19–23 x 9–11 μm . . .	<i>P. lipai</i> (11)
13*	Pathogen of other Coleoptera	14
14	Pathogen of Staphylinidae. Primary conidia 21–26 x 11–14 μm	<i>P. philonthi</i> (12)
14*	Pathogen of other Coleoptera	15
15	Pathogen of Curculionidae. Primary conidia narrow ovoid, 19.5–23 x 10–12.5 μm , or ellipsoid, 23–31 x 11.5–14.5 μm	<i>P. phyllobii</i> (13)
15*	Pathogen of Chrysomelidae. Primary conidia ovoid or ellipsoid, average 17.5 x 10.6 μm	<i>P. suturalis</i> (14)
16	Primary conidia on average shorter than 24 μm	17
16*	Primary conidia on average longer than 24 μm	20
17	Primary conidia on average shorter than 18 μm . . .	<i>Zoophthora</i> sp. 1 (29)
17*	Primary conidia on average 18–24 μm	18
18	Pathogen of adult Staphylinidae. Primary conidia 19–22 x 5.5–6.5 μm . Capilliconidia 20–25 x 5–5.5 μm	<i>Z. tachypori</i> (28)
18*	Pathogen of other or unspecific Coleoptera	19
19	Capilliconidia 17–22 x 4.5–6 μm	<i>Z. radicans</i> (26)
19*	Capilliconidia 26–27 x 7 μm . Pathogen of Scirtidae	<i>Zoophthora</i> sp. 2 (30)
20	Pathogen of Curculionidae. Primary conidia 24–28 x 7–10 μm	<i>Z. phytonomi</i> (25)
20*	Pathogen of other Coleoptera	21
21	Pathogen of larval Catharidae. Primary conidia 40–42 x 8–9 μm	<i>Z. larvivora</i> (24)
21*	Pathogen of adult Cantharidae and Elateridae	22
22	Pathogen of Cantharidae. Primary conidia 31–32 x 10 μm , resting spores spherical, thick walled 41–44 μm	<i>Z. crassitunicata</i> (22)
22*	Pathogen of Elateridae	23

- 23 Primary conidia on average 27.6 x 9.8 μm and an L/D-ratio of 2.5. Resting spores spherical, 27.6–34.5 μm with an average of 31.1 μm . Capilliconidia unknown *Z. anglica* (21)
- 23* Primary conidia on average 27–32 x 8–12 μm , L/D ratio 2.51–3.52. Capilliconidia 31–36 x 6.5–8 μm , L/D = 4.53–4.86. Resting spores spherical, hyaline 25–46 μm with an average of 31–38 μm *Z. elateridiphaga* (23)
- 24 Resting spores spherical to subspherical, 35–62 μm , on average between 40 and 49 μm , with thick or double-layered walls, densely verrucose 25
- 24* Average diameter of resting spores smaller than 40 μm 26
- 25 Resting spores on average 45–49 μm , spherical, double-layered wall, inner layer more or less hyaline, outer layer brown. Young resting spores contain 8–24 nuclei. Pathogen of adult *Rhagozycha* spp. (Cantharidae) *Z. rhagozycharum* (27)
or *Tarichium coleopterorum* (16)
- 25* Resting spores on average 41–45 μm dark brown, spherical, double walled, inner wall hyaline, episporium brown, about as thick as inner spore wall. Mature resting spores usually binucleate. Pathogen of adult Cantharidae, probably *Malthodes* spp. *Z. crassitunicata* (22)
or *T. coleopterorum* (16)
- 26 Pathogen of larval Carabidae. Resting spores spherical, 28–46 μm in diameter, episporium dark brown *T. jaczewskii* (18)
- 26* Pathogen of Curculionidae 27
- 27 Pathogen of *Hypera* (*Phytonomus*) spp. Resting spores spherical, 26–38 μm , brown *T. phytonomi* (19), *T. punctatum* (20)
- 27* Pathogen of other Curculionidae 28
- 28 Resting spores 32–40 μm , episporium with 3.5–6 μm long spines. Pathogen of *Bothynoderes* (*Cleonus*) *punctiventris* *T. cleoni* (15)
- 28* Resting spores 28–31 μm , episporium with bulges and ridges. Pathogen of larvae and prepupae of *Hylobius abietis* *T. hylobii* (17)

LIST OF FUNGAL SPECIES

1. *Batkoa apiculata* (Thaxter) Humber (1989)

Hyphal bodies nearly spherical. Conidiophores simple, sometimes with a tendency to become digitate, originating directly from the hyphal bodies. Primary conidia 30–37 x 28–30 μm , average 35 x 30 μm , nearly spherical with prominent papillate base terminating in a short, sharp and abrupt point. Secondary conidia like primary. Resting spores hyaline, spherical, 30–45 μm diameter, zygospores or azygospores, formed laterally or terminally from hyphae. Rhizoids long and conspicuous, terminating in an irregular disc-like expansion.

Hosts: Originally described from larval and adult Lepidoptera, numerous genera of Diptera (small flies and gnats), and adult leafhoppers (Homoptera) collected in the USA (Thaxter 1888). The following hosts among Coleoptera are mentioned: *Haltica quercetorum* (Chrysomelidae) collected in Poland (Bałazy 1993) and *Rosalia alpina* (Cerambycidae) collected in Switzerland (Keller, 2008).

2. *Batkoa major* (Thaxter) Humber (1989)

Hyphal bodies nearly spherical to elliptical. Conidiophores simple sometimes with a tendency to become digitate. Primary conidia nearly spherical, 55–60 x 38–45 μm ; papilla smaller in proportion to the body of the conidium than in *B. apiculata*.

Host. Adult *Ptilodactyla serricollis* (Say) (Ptilodactylidae) collected in the USA.

3. *Conidiobolus coronatus* (Costantin) Batko (1964a)

Hyphal bodies irregular. – Conidiophores not or only slightly enlarged terminally. Nuclei not or only exceptionally staining in LPAO, very small, 2.4 μm (2.0–3.0 μm). – Primary conidia 50.5–53 x 38–40 μm (34–74 x 24–58 μm), L/D = 1.30–1.34, varying largely in size, conidial body spherical to subspherical; papilla abruptly protruding, prominent, elongate, sometimes without cytoplasmic content. – Secondary conidia like primary, other types not observed. – Resting spores 30.5–31.5 μm (16–42 μm), villose, spherical; developing terminally from hyphae. – Cystidia and rhizoids absent.

Hosts: The species is unspecific and can be isolated from soil, detritus and dead insects. It is known as primary pathogen of aphids (Homoptera). The data given above originate from *Ceutorhynchus napi* (Curculionidae) collected as larvae from soil before pupation (Keller 1987).

4. *Conidiobolus osmodes* Drechsler (1954)

Primary conidia from cultures isolated from *Hypera* (*Phytonomus*) *variabilis* Herbst (= *H. postica* (Gyll.)). Larvae are described by Ben-Ze'ev & Kenneth (1980) as follows: 21–42 μm long and 16–34 μm broad, average 29.2–34.3 x 23.1–27.2 μm , globose to ovoid; papilla broad, obtusely tapered, 2.5 x 7–10 μm . Secondary conidia like primary but smaller. Zygosporangia globose to ellipsoid with a diameter of 28–44 μm , average 34 μm . Resting spores from *H. variabilis* measured 27–44 μm , average 34.0 μm . According to King (1977) the primary conidia from culture measure 25–37 x 22–30 μm . The zygosporangia are usually formed in axial alignments with conjugating segments, globose to ellipsoidal with a diameter of 13–37 μm surrounded by smooth walls or by walls with weakly marked small ridges, wall thickness 2–6.5 μm . Cultures produce a characteristic odour of benzene hexachloride.

Hosts: The species was isolated from detritus and from dead insects. It is known as primary pathogen of *Hypera* (*Phytonomus*) spp. (Curculionidae) (Ben-Ze'ev & Kenneth 1980).

Remarks: Ben-Ze'ev & Kenneth (1980) consider *Tarichium punctatum* Garbowski and *T. phytonomi* Jaczewski as identical with *C. osmodes*.

5. *Entomophaga lagriae* Bałazy (1993)

Hyphal bodies 7–15 nucleate, subspherical with a diameter of 28–35 μm or irregularly ovoid to ellipsoid up to 80 x 15–20 μm , germinate with one or two hyphae 9–10 μm thick. Conidiophores unbranched, conidiogenous cells claviform 65–102 x 16–21 μm . Primary conidia 35–43 x 24–30 μm , ovoid or broadly turbinate with regularly semispherically rounded top parts, with 12–18 nuclei, papilla narrow con-

ical or somewhat obtuse, 9–11.5 μm wide. Secondary conidia like primary. Resting spores unknown.

Host: Lagria hirta L. (Lagriidae) collected in Switzerland.

6. *Entomophthora* sp. (Eilenberg 1987)

Primary conidia (17–) 21.6–24.4 (–29) \times (13–) 17.1–19.9 (–24) μm , L/D = 1.23–1.28, with 8–17 nuclei. Secondary conidia (16–) 15.9 (–21) \times (16–) 19.3 (–24) μm , L/D = 1.21. Resting spores probably azygospores (39–) 44.5 (–50) \times (31–) 40.4 (–40) μm . Rhizoids absent.

Host: Adult Rhagonycha fulva (Scopoli) (type host) and *Cantharis livida* L. (Cantharidae), collected from June to August on grass and lower vegetation of a hedge in Denmark.

7. *Erynia nebriae* (Raunkiaer) Humber & Ben Ze'ev (1981)

Conidiophores branched, septate, hyaline, club-like enlarged, 11–15 μm diameter. Conidia ellipsoidal to fusiform, asymmetrical, often slightly curved, 28–37 μm long, 10–13 μm broad, hyaline, smooth. Resting spores hyaline to pale brown, developing outside the body of the host, spherical, 35–50 μm diameter.

Host: Nebria brevicollis (F.) (Carabidae) collected in Sealand, Denmark.

Remarks: According to the index of fungi (<http://www.indexfungorum.org>) *Erynia nebriae* is an invalid name. Consequently, the name *Entomophthora nebriae* given by Raunkiaer (1893) is still valid, although there is no doubt that the species does not belong to *Entomophthora*. It is certainly a member of the subfamily Erynioideae but the data given in the original description do not allow to attribute the species to a genus. Therefore, we do not validate the name but consider *Erynia nebriae* as more appropriate.

Below in the paragraph «unclassified fungi», we mention a fungus as *Erynia* sp. found on the same host species, unfortunately with unprecise data. The two species are probably identical.

8. *Eryniopsis lampyridarum* (Thaxter) Humber (1984a)

Conidia 20–37 \times 14–30 μm , average 35 \times 15 μm , regular ovoid, slightly tapering towards the apex, with an abrupt slightly papillate base, content granular without larger oil globules. Secondary conidia like the primary or more commonly long-cylindrical, rounded at either end and born vertically on capillary conidiophores. Resting spores unknown. Host fixed to leaves by its mandibles.

Hosts: Adult Chauliognathus pensylvanicus (De Geer) (Cantharidae) collected in the USA.

9. *Furia zabri* (Rozsypal ex Ben-Ze'ev & Kenneth) Humber (1989)

Hyphal bodies are about 11 μm thick and richly branched with a larger number of nuclei. The hemolymph contains fragmented mycelium, spherical and lobular shapes are present. Cystidia up to 300 μm long, the flask-shaped base has a diameter of 40–50 μm . Conidiophores branched. Primary conidia ovoid, 25 \times 14 μm , maximum 29 \times 18 μm . Zygospores spherical, formed mostly on the surface of the host. Rhizoids present.

Host: Larvae of *Zabrus tenebrioides* Goeze (Carabidae) collected in Moravia, Czech Republic.

10. *Pandora brahminae* (Bose & Metha) Humber (1989)

Conidiophores branched. Rhizoids monohyphal with a diameter of 14–24 μm , endings with disc-like holdfast. Primary conidia ovoid, 18–22 x 11–15 μm , tapering to a papillate base. Secondary conidia like primary. Resting spores zygosporous, formed inside the host at the same time as the conidia, hyaline to light brown, 22–39 μm , average 29 μm , thick walled; epispodium spiny, cannot be separated by pressure.

Host: Adult *Brahmina* sp. (Melolonthidae), collected in Uttar Pradesh, India. Dead beetles fixed with mouthparts and rhizoids to leaves of chestnut trees.

11. *Pandora lipai* (Bałazy, Eilenberg & Papierok) Keller in Keller & Petrini (2005)

Hyphal bodies irregularly elongate, hyphae-like, 50–70 x 7–12 μm , sometimes branched. Conidiophores 7–10 μm thick, richly branched. Primary conidia narrow ovoid to somewhat clavate, (18.5–) 19.5–23 (–24) x (8.5–) 9–10.5 (–11) μm , papilla slightly convex, 3.8–5 μm wide. Secondary conidia ovoid, shorter but of similar thickness as primary ones, 14.5–17.5 x 8.5–10.5 μm , strongly narrowing towards the base closed by semispherical papilla 2.5–3.5 μm wide. Cystidia numerous, oblong conical, 150–250 μm long, 10–17 μm thick at the level of the conidiogenous layer, 4.5–7 μm thick at the blunt tips. Rhizoids monohyphal terminated with wide, flat disc-like holdfast with uneven margins. Resting spores known only from cultures; regularly globose, diameter 20–33 μm and 2.3–3.0 μm thick walls; epispore minutely rough, brown.

Hosts: *Cantharis livida* L., *C. rustica* Fallen, *Malthinus flaveolus* (Herbst), and *Rhagonycha lignosa* (Müller) (Cantharidae) collected between end of May and mid June in Poland and Denmark.

Remarks: *Pandora lipai* is possibly a species complex or a species that shows a wide variation in conidial dimensions. Keller (2007b, 2012) found two groups of fungi attributed to *P. lipai* in a mixed host population consisting of *Ancistronycha* (*Cantharis*) *abdominalis* (F.) and *A. (C.) erichsonii* Bach. One fungus group had primary conidia measuring (22–) 23.9–25.1 (–28) x (11–) 12.3–13.6 (–15) μm (L/D = 1.80–2.05) and the other one had primary conidia measuring (9.5–) 17.6–19.0 (–21) x (9.5–) 10.1–10.9 (–12) μm (L/D = 1.74). The dimensions of the conidia of the two fungus groups clearly differ from each other but they both slightly overlap with those given in the original description. Further investigations are needed to clarify the situation.

12. *Pandora philonthi* (Bałazy) Keller in Keller & Petrini (2005)

Hyphal bodies irregularly elongate, 40–250 x 15–25 μm . Conidiophores 7–9.5 μm thick, richly branched, grow in intersegmental junctions, mouth parts and between articles of legs and antennae. Terminal conidiogenous cells mostly flask-shaped about 25 x 13 μm . Primary conidia ellipsoid (19–) 21–26 (–28) x (10.5–) 11–14 (–14.5) μm , rarely ovoid, slightly but abruptly narrowed at the bases, forming very short overpapillar neck; papillae of rather distinctly arcuate outline or semiglobose,

5–6.5 μm wide. Secondary conidia similar, mostly ovoid, smaller. Few cystidia scattered in area of intensive conidiation, rather short, up to 120 μm long and about 10–14 μm thick at the base, unmarkedly tapering to obtuse tip. Rhizoids monohyphal, ribbon-like, 8–10 μm wide, terminated with irregularly rounded reticulate holdfasts, 20–40 μm in diameter, grow mostly in thoracic part of the host's body. Resting spores unknown.

Host: *Philonthus* sp. (Staphylinidae), collected in Poland in forests and on agricultural land.

13. *Pandora phyllobii* (Bałazy) Keller in Keller & Petrini (2005)

Hyphal bodies irregularly elongate, often with branches and outgrowths, about 20–130 x 7–21 μm . Conidiophores 6–9.5 μm thick typically branched in apical part. Conidiogenous cells cylindrical or clavate, 22–28 x 10–12.5 μm . Primary conidia narrow ovoid, 19.5–23 x 10–12.5 μm , or ellipsoid, 23–31 x 11.5–14.5 μm , basal part conically attenuated towards small papilla, about 4 μm wide. Secondary conidia short ovoid or almost subglobose with small papillae strongly protruding. Rhizoids monohyphal with terminal disc-like holdfast with unevenly sinuous margins. Cystidia 180–400 μm long, 9–11 μm wide at the base, slightly attenuate upwards. Resting spores unknown.

Host: Adult *Phyllobius* sp. (Curculionidae), collected in Poland in June.

14. *Pandora suturalis* (Ben-Ze'ev) Humber (1989)

Hyphal bodies short hyphal segments, up to 120 μm long, usually not branched, plurinucleate. Conidiophores digitately branched with 3–7 conidiogenous cells. Primary conidia ovoid or ellipsoid, most of them of subpapillate type, 14.3–24.6 x 8.0–16.6 μm , average 17.5 x 10.6 μm , L/D ratio on average 1.67; usually with 2–3 vacuoles; papillae usually less than half of the conidial diameter. Secondary conidia pyriform or ovoid-pyriform, 13.2–15.5 x 9.7–11.5 μm , average 14.3 x 11.0 μm , L/D ratio on average 1.3. Cystidia 15–20 μm at the base and about 150–200 μm long, tapering. Rhizoids abundant, emerging in fascicles that appeared as pseudo-rhizomorphs, but diverged into individual, monohyphal threads, up to 35 μm wide, sometimes branching in the middle, with dichotomically lobed holdfasts.

Host: Adult *Lochmaea suturalis* (Thomson) (Chrysomelidae), collected in Scotland near Edinburgh.

Remarks: Data taken from exsiccata.

15. *Tarichium cleoni* (Wize) Lakon (1915)

The following description is taken from Keller *et al.* (2009). Resting spores 32.2–40.1 (28–47) μm , spherical to subspherical, red in mass and light brown in microscopic preparations, with regularly arranged 3.5–6.0 μm long spines. The endospores have a diameter of 21.2–23.6 (18–28) μm , the spore wall measures on average 3.5–4.0 μm . Endospores and epispores are tightly connected. In the light microscope the two walls appear clearly separated. The diameter of the inner wall measures 16.8–17.9 (14–21) μm .

Host: *Bothynoderes* [= *Asproparthenis*] (*Cleonus*) *punctiventris* (Germar) (Curculionidae) collected in the Ukraine.

16. *Tarichium coleopterorum* (Petch) Balazy (1993)

Rhizoids pseudorhizomorph emerging from two points on the lower surface of the body of the insect. The resting spores were dark brown, thick-walled (up to 6 μm), densely verrucose, globose, 35–50 μm diameter, or broadly oval, 48–52 x 44–46 μm , or sometimes pyriform, 56 x 44 μm .

Host: Unidentified Coleoptera collected in England.

Remarks: The spherical resting spores resemble those of *Zoophthora crassitunicata* and *Z. rhagonycharum*.

17. *Tarichium hylobii* Keller, Weiser & Wegensteiner (2009)

Mature resting spores measure 28.1–31.1 (24–36) μm , spherical to subspherical, grey in mass and light brown in microscopic preparations. In the light microscope the silhouette appears sinuous. The electron microscope shows irregularly undulating bulges and ridges. In the light microscope the separation between endospore and epispore is not clear. The diameter of the inner wall measures 19.4–20.2 (16–25) μm .

Host: Larvae or prepupae of *Hylobius abietis* (L.) (Curculionidae). The material originated from a closed pupal cell collected near Prague, Czech Republic.

18. *Tarichium jaczewskii* Zaprometov (1928)

Resting spores zygo- or azygospores, spherical, 28–46 μm in diameter, with almost smooth or verruculous two-layered epispore 4.5–7 μm thick, dark brown in colour. The infected larvae become black, dry, mummified and brittle.

Host: Larvae of *Zabrus gibbosus* (Zimm.) (Carabidae) collected at Samarkand in 1926 and at Katta-Kurgan in 1928 (former USSR). *T. jaczewskii* is possibly the resting spore state of *Furia zabri* (Ben-Ze'ev & Kenneth 1982).

19. *Tarichium phytonomi* Jaczewski in Zaprometov (1928)

The following data are taken from MacLeod & Müller-Kögler (1970). Resting spores zygo- or azygospores, spherical, 32–36 μm in diameter. Epispore dark brown, 2.6–3.9 μm thick with a few irregularly scattered warts that are 1.3–2.6 μm high.

Host: Larvae of *Hypera (Phytonomus) variabilis* [sic] (Curculionidae) collected in Taschkent (former USSR).

Remarks: As MacLeod & Müller-Kögler (1970) stated, the name is illegitimate as a later homonym of *Zoophthora phytonomi* (Arthur) Batko (1964b). Ben-Ze'ev & Kenneth (1980) considered *T. phytonomi* as the resting spore state of *Conidiobolus osmodes* and as conspecific with *T. punctatum*.

20. *Tarichium punctatum* Garbowski (1927)

The following data are taken from MacLeod & Müller-Kögler (1970). Resting spores spherical, 26–37.5 μm in diameter, smoky dark in colour. Spore wall 2–3 μm thick with tiny spots that give it a punctate appearance.

Host: Larvae of *Hypera (Phytonomus) variabilis* [sic] (Curculionidae) collected in Poland. Ben-Ze'ev & Kenneth (1980) considered *T. punctatum* as the resting spore state of *Conidiobolus osmodes* and as conspecific with *T. phytonomi*.

21. *Zoophthora anglica* (Petch) Humber (1989)

Petch (1943) described this species based on fungi found on the following hosts: *Plateumeris* sp. (Chrysomelidae), *Agriotes sputator* (L.) (Elateridae), *Lochmaea suturalis* (Thomson) (Chrysomelidae), and *Cantharis* sp. (Cantharidae) collected in England. He described the fungus as follows: Primary conidia narrow oval or sub-fusoid, sometimes slightly bent, with a broad truncate papilla, 22–27 x 11–13 μm , secondary conidia of the same shape but shorter, 18–21 x 10–11 μm . Rhizoids are produced from the ventral surface and the sides. The conidiophores in fully developed examples extend in a continuous sheet from the sides of the body to the substratum, but in general they do not spread over the elytra or only partially along their outer edges.

Ben-Ze'ev (1986) re-examined the air-dried material and found that it contained more than one fungus species and that the description is a mixture of different fungi. He separated it according to the host species. The fungus from *Agriotes sputator* had primary conidia measuring 16.7–33.4 x 8.0–13.8 μm , with an average of 27.6 x 9.8 μm and an L/D-ratio of 2.5. Most nuclei were ellipsoidal and measured 4.6 x 6.8 μm . The spherical resting spores measured 27.6–34.5 μm with an average of 31.1 μm . The spore wall was 1.2–2.3 μm thick. Secondary conidia, neither type Ia nor capilliconidia were present. He found the fungus to be identical with *Z. elateridiphaga* and considered the latter as a synonym of *Z. anglica*.

The two fungi are indeed closely related. Nevertheless, there are differences between them, which were discussed by Keller (1991). They justify to consider the two species as distinct.

Hosts: Adult *Agriotes sputator* (L.) (Elateridae) collected in England.

22. *Zoophthora crassitunicata* Keller (1980)

Rhizoids pseudorhizomorph. Conidiophores sparingly branched, oligonucleate. Primary conidia (25–) 31.4–32.1 (–36) x (8.5–) 9.8 (–12) μm , L/D = 3.2–3.3, subcylindrical to slightly fusiform, papilla distinct, conical. Capilliconidia 33–39 x 8–9 μm , fusiform curved to banana-shaped. Resting spores (35–) 41.2–44.7 (–56) μm , spherical, double walled, inner wall hyaline, epispodium brown, about as thick as inner spore wall, densely and regularly covered with minute knobs, predominantly binucleate, but up to 8 nuclei observed, formed inside the host.

Host and symptoms: Adult cantharid beetle, probably *Malthodes* sp. (Cantharidae). Wings of the host insect closed when the fungus sporulated.

Remarks: The dimensions of the thick-walled resting spores match those of *Tarichium coleopterorum*. The surface structure as well is similar, «densely verrucose» (*T. coleopterorum*) and «densely and regularly covered with minute knobs» (*Z. crassitunicata*). A difference concerns the shape of the resting spores. Those of *Z. crassitunicata* were all spherical while some of *T. coleopterorum* were oval or pyriform. In spite of the large similarities some doubts concerning a conspecificity of the two species remain.

E. crassitunicata was hitherto known only from the type locality (Rickenbach, Switzerland, canton Zurich). Recently, it was found on a cantharid beetle (probably *Malthodes* sp.) at Fischingen (Switzerland, canton Thurgau).

23. *Zoophthora elateridiphaga* (Turian) Ben-Ze'ev & Kenneth (1980)

Rhizoids pseudorhizomorph on ventral side of thorax between the legs, holdfasts disc-like, or unspecialised and sometimes united to form a layer. Hyphal bodies elongate with (5–) 9–12 (–22) nuclei; nuclei measuring on average 5.1–5.7 μm . Conidiophores branched, terminally slightly enlarged. Primary conidia (24–) 27.5–31.6 (–36) \times (7–) 8.8–11.4 (–13) μm , L/D ratio 2.51–3.52, cylindrical, papilla distinct, conical. Secondary conidia similar to primary, (21–) 25–25.7 (–30) \times (7–) 8.6–9.9 (–12) μm , L/D = 2.5–3.0, or capilliconidia (27–) 31.4–35.5 (–42) \times (6–) 6.7–7.5 (–9) μm , L/D = 4.53–4.86, fusiform to banana-like; length of capillary tube (67–) 86–90 (–109) μm . Resting spores formed inside the host, (25–) 31.2–37.7 (–46) μm , spherical, hyaline, smooth; zygospores or azygospores, young resting spores with (14–) 18–21 (–29) nuclei measuring 4.9–5.0 μm . Fully developed but broken resting spores contained (6–) 10 (–14) nuclei.

Hosts: Adult *Agriotes sputator* (L.) (type host), *A. lineatus* (L.) (Elateridae) collected in Switzerland. Recently, Grabenweger (pers. comm.) reported heavy epizootics in laboratory rearings of *A. lineatus*, *A. obscurus* and *A. sputator* at Reckenholz, Zurich.

24. *Zoophthora larvivora* Bałazy (1993)

Mycelium inside host's body consists of almost equally narrow hyphae 7–11 μm thick, rarely as separate hyphal segments up to 40 μm long. Conidiophores 6–8 μm thick, digitately branched, terminal conidiogenous branches cylindric or indistinctly clavate 32–47 \times 7.5–11.5 μm . Primary conidia cylindric with obtuse ends or with distal parts weakly, steadily narrowing towards blunt apex, (35–) 40–42 (–49.5) \times (7.5–) 8.5–9 (–10.7) μm , L/D ratio commonly exceeding 4.5. Secondary conidia, resting spores and cystidia unknown. Few rhizoids, pseudorhizomorph.

Host: A single larva of an unidentified Cantharidae collected in Poland.

25. *Zoophthora phytonomi* (Arthur) Batko (1964b)

Hyphal bodies branched, non-septate, colourless, 9–12 μm in diameter. Rhizoids on the ventral surface. Hymenium over the whole surface except the head. Conidiophores branched at the base, as thick as mycelium. Primary conidia oblong, colourless, 24–28 \times 7–10 μm . Resting spores not observed.

Host: Larval *Hypera punctata* (F.) (syn.: *Brachyptera zoilus* (Scop.)) (type host) collected in the USA. Dead insects fixed on top of plants. Further hosts are *H. postica* (Gyll.) (syn.: *H. variabilis*) (Curculionidae).

26. *Zoophthora radicans sensu lato* (Brefeld) Batko (1964a)

Hyphal bodies hyphae-like. Conidiophores on average 6–8 μm thick, branched with nuclei measuring 4.7 (4–5.5) μm , conidiogenous cells terminally enlarged to 7–11 μm . Primary conidia 15.5–22.5 \times 6.3–8.4 μm (13–28 \times 5–10 μm), L/D = 2.41–2.98, subcylindrical to subfusiform straight or slightly bent; papilla usually rounded; diameter of nuclei 4.0–4.6 (3–5.5) μm . Secondary conidia similar to primary, 11.9–13.2 \times 6.9–7.1 μm (10–16 \times 6–9 μm), L/D = 1.67–1.88, or capilliconidia 17.3–21.8 \times 4.6–5.9 μm (15–24 \times 4–7.5 μm), L/D = 3.23–4.18, fusiform, slightly bent or straight, apically rounded; length of capillary tube 42.5–58.8 μm (30–81

μm). Resting spores 24.5–27.2 μm (20–33 μm), spherical, hyaline, smooth. Rhizoids abundant, emerging ventrally or latero-ventrally, pseudorhizomorph or monohyphal with disc-like ending. Cystidia not observed. The data given above are taken from Keller (1991). According to Bałazy (1993), the primary conidia measure (17.5–) 19.5–22.5 (–27.5) \times (4.7–) 5.5–6 (–6.5) μm with an L/D ratio of 3.5–4.5.

Hosts: Many species of the insect orders Plecoptera, Heteroptera, Homoptera, Diptera, Hymenoptera and Lepidoptera. Type host is *Pieris brassicae* (L.) (Pieridae, Lepidoptera).

Remarks: The original description is very poor and a redescription has never been done. Therefore, it is very difficult to separate morphologically similar fungi from the type species. Consequently, *Zoophthora radicans* seems to be an unspecific species, but should be considered as a species complex. Bałazy (1993) has tried to split the complex based on tiny morphological differences but mainly by their hosts. Genetic markers should be used to prove the correctness of this separation.

So far, there are no confirmed reports of *Z. radicans* attacking Coleoptera. However, Lakon (1919) mentioned a report of Bail who observed an epizootic among *Nebria brevicollis* (L.) (Carabidae) at Jäschkental (now Jaškowa Dolina near Gdansk, Poland) caused by *Entomophthora sphaerosperma* which is a synonym of *Z. radicans*. As listed below in the paragraph «unclassified fungi», this fungus could also be identical with *Erynia* sp. reported from the same host species.

27. *Zoophthora rhagonycharum* (Bałazy) Keller (2007)

Rhizoids pseudorhizomorph, numerous, mainly emerging from the ventral side of the abdomen, endings unspecialised or with disc-like holdfast. Hyphal bodies in living hosts filamentous usually unbranched, with 7–16 nuclei with a diameter of 6–9.5 μm . Conidia unknown. Resting spores (33–) 45.4–48.8 (–62) μm , spherical, double-layered wall; inner layer more or less hyaline, outer layer (episporium) brown with minute knobs at the surface. Young resting spores contain 8–24 nuclei with a diameter of 5–6 μm .

Host: Adult *Rhagonycha lignosa* (Müller) (type host) collected in Poland and *R. fulva* (Scopoli) (Cantharidae) collected in Switzerland.

Remarks: The average dimensions of the resting spores are within the range of the extreme dimensions of those of *Tarichium coleopterorum* although at the upper limit. Nevertheless, an identity of the two species cannot be excluded.

Fitton (1982) observed a fungus occurring epizootically in populations of *R. fulva* at Silwood Park in England. The fungus was identified at the Commonwealth Mycological Institute as *Entomophthora coleopterorum*. However, the host and the symptoms described by Fitton make it very probable that the observed fungus is *Z. rhagonycharum*. At least, his observations are a strong indication that *Tarichium* (*Entomophthora*) *coleopterorum* and *Z. rhagonycharum* might be identical.

28. *Zoophthora tachypori* Bałazy (1993)

Hyphal bodies elongate, 25–80 \times 9–15 μm , often with short outgrowths and branchlets. Conidiophores 6.5–12 μm thick, richly digitately branched; ultimate sporogenous branches about 20–24 μm long, cylindrical or slightly widened near the top, 6–7 μm thick. Primary conidia cylindrical, seldom somewhat subfusiform, (18–) 19.5–21.5 (–22) \times 5.5–6 (–6.5) μm . Secondary conidia like primary or capilliconi-




Plate 1. *Zoophthora* sp. 2 ex *Elodes minuta*. — 1: Mycelium on beetle (nat. size 5 mm); — 2: Primary conidia with typical, separated outer wall; — 3: Branched conidiophores; — 4: Primary conidia, a detached capilliconidium (left upper corner) and two developing type I secondary conidia; — 5: Primary conidia and a developing type I secondary conidium (left upper corner). — Bars in figs 2–5 represent 20 μm . (Photos: R. Wegensteiner).

dia, 20–25 \times 5–5.5 μm , oblong fusiform. Capillary tube 40–50 μm long. Cystidia conical, 30–80 (–120) μm long and 8–12 μm thick at the conidial layer. Pseudorhizomorphs growing from thoracic parts, monohyphal rhizoids produde from abdominal sternites. Resting spores unknown.

Host: Adult *Tachyporus* spp. (Staphylinidae) collected in Poland.

29. *Zoophthora* sp. 1 (Keller, 2012)

Rhizoids compound with common disc-like holdfast, lateral parts sometimes splitting into thinner compound rhizoids which are further divided into monohyphal holdfasts. Hyphal bodies hyphae-like, branched or unbranched with a diameter of (5–) 6.5 (–8) μm ($n = 25$). Conidiophores branched with a diameter of 5–6 μm . Primary conidia (13–) 15.6 (–17) \times (5–) 5.9 (–6) μm , $L/D = 2.65$ ($n = 25$), narrow fusiform to ellipsoidal, bitunicate, papilla blunt, separated by an indistinct bulge from the body of the conidium. Cystidia slender, tapering, basal part faintly stained with a diameter of 5–6 μm , terminal parts slightly enlarged and darker stained. Secondary conidia and resting spores not present.

Host: Unidentified species of Cantharidae (Coleoptera), probably *Malthodes* sp. collected at Fischingen, canton Thurgau, Switzerland.

Remarks. The fungus was collected in early July on the underside of a leaf of *Corylus avellana* L. about 1.5 m above ground on the shore of the brook Murg at an altitude of about 650 m. The fungus is considered as a new species but the material is too incomplete to give a formal description.

30. *Zoophthora* sp. 2 (Plate 1, figs. 1–5)

The wings of the host were slightly opened, when the fungus sporulated (fig. 1). Rhizoids compound. Conidiophores branched (fig. 3). Primary conidia (20–) 22.8–23.4 (–26) x (9–) 10–10.5 (–12) μm , L/D = 2.17–2.34, subcylindric to slightly ellipsoidal, bitunicate; papilla indistinctly demarcated from the conidial body, conical with rounded end (figs. 2, 4 and 5). Secondary conidia like primary but shorter (fig. 5), or capilliconidia 26–27 x 6.5–7.5 μm (n = 3), L/D = 3.8, elongate fusiform and slightly bent, banana-like, usually with a single large vacuole (fig. 4).

Host: *Elodes minuta* L. (Scirtidae).

Location: Berglsteinersee, 714 m altitude, 3.8 km NE of Rattenberg, Kramsach, Tirol, Austria. The material was collected on June 19, 2011 by C. Tkaczuk and identified by the author. The fungus is probably an undescribed species.

Unclassified fungi

Entomophthora telaria Giard (1888)

This entomophthoralean fungus was recorded as pathogen of *Rhagonycha melanura* F. (Cantharidae) in northern France near Fécamp. Giard did not give any data of the fungus and described only the symptoms. According to him the fungus is characterized by a large membrane which covers the body of the host and fixes it to the substrate. The hyphae emerging from the host's body seemed to be united by a rubber-like secretion which looks like a cloth. The infected beetles were fixed to the underside of leaves of *Galeopsis tetrahit* L. The cadavers were always placed parallel to the median vein with the head towards the stem.

From the symptoms described we can suppose that this fungus is close to *Pandora lipai* but this assumption needs verification.

Entomophthora carpentieri Giard (1888)

This fungus was found by Mr. Carpentier near Amiens in northern France infecting *Agriotes sputator* L. (Elateridae). Giard described the symptoms very precisely and in the author's opinion there is no doubt that the fungus is identical with *Zoophthora elateridiphaga* (Turian) Ben-Ze'ev & Kenneth. Giard especially noticed the strong compound rhizoids unknown till then among entomophthoralean fungi. Therefore, he proposed *Lophorhiza* as a new genus for this fungus. However, the name was apparently overlooked or forgotten and has never been used by other authors. The genus name *Zoophthora* now generally accepted for species with compound rhizoids was proposed by Batko (1964a) who was obviously not aware of the existence of an older genus name.




Plate 2. *Erynia* sp. ex *Nebria brevicollis*. — 1: Front view of the infected beetle with fully sporulating fungus; — 2: Lateral view of the same beetle (same magnification); — 3: Macrophotograph showing structures considered as cystidia towering above the sporulating fungus layer (unknown magnification). (Photos: U. Arnscheidt).

Erynia sp. (Plate 2, figs. 1–3)

A fungus collected on 17.11.2011 on two specimens of adult *Nebria* cf. *brevicollis* in Germany by Mr. Udo Arnscheidt was provisionally identified as *Erynia* sp. by the author. The identification is based on photos and measurements sent to the author by Mr. Arnscheidt. The photos showed extremely swollen cadavers, when the fungus sporulated, the wings were spread (figs. 1–2). Close-up photos showed structures which were interpreted as cystidia (fig. 3). These were powerful, often with an enlarged ending. Such strong cystidia are typical for *Erynia* spp. That's why the fungus was provisionally attributed to this genus. Unfortunately, the measurements of the conidia did not differentiate between primary conidia and conidia of higher orders and probably included also other structures. The average dimension of conidia considered as primary ones was calculated (based on photos) to be about 23–25 x 13 μm with an L/D ratio of 1.8–2.0. The largest conidium measured 34 x 16.5 μm , the length is within the range of the primary conidia of *Erynia nebriae* whose conidia measure 28–37 x 10–13 μm with an L/D ratio distinctly above 2. Considering the identity of the host and the similarities of the conidia an identity of the two fungi cannot be excluded.

Erynia sp. was found in a forest called Hoersterholz, near Bochum, Germany, altitude 120 m, coordinates: 51°26'24.60 N; 7°8'52.14 E. The two sporulating ground beetles were on the soil in a forest dominated by *Fagus sylvatica* L. The two cadavers were found at a distance of 50 cm from each other.

***Tarichium* sp.**

This fungus was found in the state of New York, USA, as pathogen of adult *Diabrotica barberi* Smith & Lawrence (Chrysomelidae) (Naranjo & Steinkraus, 1988). The infected beetles had a swollen abdomen and raised elytra. The resting spores produced inside the body of the host were smooth-walled and spherical with a mean diameter of $27.2 \pm 2.8 \mu\text{m}$ and contained 8–14 nuclei with a mean diameter of $5.8 \pm 2.0 \mu\text{m}$. Cystidia and rhizoids were absent. Size and structure of the nuclei allowed to attribute the species to the family Entomophthoraceae.

DISCUSSION

Most of the species listed above are well described. Only a few of them are doubtful or lack a formal description due to missing details or for other reasons. Doubtful species are *Tarichium coleopterorum* (Petch) Bałazy, *T. jaczewskii* Zaprometov, *T. phytonomi* Jaczewski in Zaprometov and *T. punctatum* Garbowski.

According to the Index Fungorum (<http://www.speciesfungorum.org>) *T. coleopterorum* is an invalid name. Petch (1932) described the species as *Entomophthora coleopterorum* which is still the valid name. Nevertheless, *T. coleopterorum* refers to the fact that the fungus is known only in its resting spore state. Later on Petch (1944) found a conidial state on other host species (larval Coleoptera and *Sitones flavescens* Marsch. (= *Sitona obsoletus* (Gmelin))) and attributed it to *E. coleopterorum*, although he was in doubt of the conspecificity of the two fungi. These conidia measured 32–44 x 8–14 μm . However, the correct specific attribution of the conidial state remains unsolved (Ben-Ze'ev 1986). As discussed under *Z. rhagonycharum*, there are indications that *T. coleopterorum* might be identical with this species.

A possible synonymy of *T. jaczewskii* Zaprometov with *Furia zabri* has been discussed by Ben-Ze'ev & Kenneth (1982), however, an identity of the two species has not been proven. On the other hand Ben-Ze'ev & Kenneth (1980) gave good evidence, that *T. phytonomi* Jaczewski in Zaprometov and *T. punctatum* Garbowski are not only identical but also the resting spore stage of *Conidiobolus osmodes*. Due to remaining doubts they did not formally synonymize the three species.

Zoophthora sp. 1 (nr. 29) and *Zoophthora* sp. 2 (nr. 30) are incompletely known but supposed to be new species. *Pandora lipai* (nr. 11) is possibly a species complex. These three species need further investigations to clarify the situation. *Entomophthora* sp. (nr. 6) is well characterized and represents without doubt a new species. However, a formal description is still missing.

Although Coleoptera is the order with the highest number of described species, comparably low is the number of Entomophthorales known to attack these insects. Only eleven families are known as hosts of Entomophthorales, Cantharidae and Curculionidae are the preferred ones. Reasons for the relative scarcity of these fungi in Coleoptera are the unawareness of entomologists, the often inconspicuous symptoms of infections and characteristics of the Coleoptera like the strongly sclerotized body, which is supposed to act as a barrier to fungus infections. Therefore it is not surprising that larvae and soft-skinned adults like Cantharidae are the preferred hosts. Nevertheless, Entomophthorales can infect strongly sclerotized beetles like adult Elateridae and are even able to cause epizootics among them (Keller, 1994).

Entomophthorales and their role in the population dynamics of insects are still too less known. The key provided in this paper hopefully will animate coleopterists, other entomologists and insect mycologists to keep an eye out for these interesting fungi and thus contribute to a better knowledge of the diversity of Entomophthorales and their ecological importance.

ZUSAMMENFASSUNG

Käfer sind die artenreichste Insektenordnung. Trotzdem sind im Vergleich zu anderen Insektenordnungen nur wenige Arten von Entomophthorales bekannt, die Käfer infizieren. Gründe dafür sind unter anderem die mangelnde Bekanntheit dieser Pilze bei Entomologen, die oftmals undeutlichen Symptome von Infektionen sowie käferspezifische Merkmale wie das ausgeprägte und wahrscheinlich vor Infektionen schützende Exoskelett.

Die vorliegende Arbeit erfasst 30 Arten von Entomophthorales, darin eingeschlossen drei formal nicht beschriebene. Der dichotome Schlüssel erlaubt die Bestimmung dieser 30 Arten vorwiegend anhand von Primärkonidien und Wirtsarten. Aber auch dem Umstand, dass relative viele Pilzarten nur in der Dauersporenform bekannt sind, wird Rechnung getragen. Dem Bestimmungsschlüssel geht eine Zusammenstellung voraus, bei der die einzelnen Pilzarten elf Käferfamilien zugeordnet sind. Dies erleichtert in vielen Fällen die Bestimmung.

ACKNOWLEDGEMENTS

The author thanks Petr Zabransky and the late Michel Brancucci for identifying *Elodes minuta*, Cezary Tkaczuk for providing the material collected at Berglsteinersee and Rudolf Wegensteiner for providing measurements and photos of *Zoophthora* sp. 2, and Jørgen Eilenberg for providing the original description of *Entomophthora nebriae*.

REFERENCES

- Bałazy, S. 1993. Flora of Poland. Fungi (Mycota), vol. 24, Entomophthorales. — Polish Academy of Sciences, 356 pp.
- Batko, A. 1964a. On the new genera: *Zoophthora* gen. nov., *Triplosporium* (Thaxter) gen. nov. and *Entomophaga* gen. nov. (Phycomycetes: Entomophthoraceae). — Bulletin de l'académie polonaise des sciences Cl. II. Série des sciences biologiques, Vol. XII, No. 7, 323–326.
- Batko, A. 1964b. Some new combinations in the fungus family Entomophthoraceae (Phycomycetes) — Bulletin de l'académie polonaise des sciences Cl. II. Série des sciences biologiques, Vol. XII, No. 7, 403–406.
- Batko, A. 1966. On the subgenera of the fungus genus *Zoophthora* Batko 1964 (Entomophthoraceae) — Acta Mycologica 2: 15–21.
- Ben-Ze'ev, I. 1986. Notes on Entomophthorales (Zygomycotina) collected by T. Petch: *Erynia anglica* comb. nov. and *Erynia coleopterorum*. — Mycotaxon 25: 1–10.
- Ben-Ze'ev, I. & Kenneth, R.G. 1980. *Zoophthora phytonomi* and *Conidiobolus osmodes* (Zygomycetes, Entomophthoraceae), two pathogens of *Hypera* species (Col., Curculionidae) coincidental in time and place. — Entomophaga 25: 171–186.
- Ben-Ze'ev, I. & Kenneth, R.G. 1982. Features-criteria of taxonomic value in the Entomophthorales: I. A revision of the Batkoan classification. — Mycotaxon 14: 393–455.
- Drechsler, C. 1954. Two species of *Conidiobolus* with minutely ridged zygospores. — American Journal of Botany 41: 567–575.
- Eilenberg, J., Bresciani, J., & Martin, J. 1987. *Entomophthora* species with *E. muscae*-like primary spores on two new insect orders, Coleoptera and Hymenoptera. — Nordic Journal of Botany 7: 577–584.
- Fitton, M.G. 1982. A fungus attacking the soldier beetle *Rhagonycha fulva* (Coleoptera: Cantharidae) in southern England. — Entomologist's Gazette 33: 215–219.
- Giard, A. 1888. Fragments biologiques. XI. Sur quelques Entomophthorées. — Bulletin Scientifique de la France et de la Belgique 19: 298–309.
- Humber, R.A. 1984. *Eryniopsis*, a new genus of the Entomophthoraceae (Entomophthorales). — Mycotaxon 21: 257–264.
- Humber, R.A. 1989. Synopsis of a revised classification for the Entomophthorales (Zygomycotina). — Mycotaxon 34: 441–460.

- Humber, R.A. 2012. Entomophthoromycota: a new phylum and reclassification for entomophthoroid fungi. — *Mycotaxon* 120: 477–492.
- Humber, R.A. & Ben Ze'ev, I. 1981. *Erynia* (Zygomycetes: Entomophthorales): Emendations, synonymy, and transfers. — *Mycotaxon* 13: 506–516.
- Keller, S. 1980. Two new species of the genus *Zoophthora* Batko (Zygomycetes: Entomophthoraceae): *Z. lanceolata* and *Z. crassitunicata*. — *Sydowia* 33: 167–173.
- Keller, S. 1987. Arthropod-pathogenic Entomophthorales of Switzerland. I. *Conidiobolus*, *Entomophaga* and *Entomophthora*. — *Sydowia* 40: 122–167.
- Keller, S. 1991. Arthropod-pathogenic Entomophthorales of Switzerland. II. *Erynia*, *Eryniopsis*, *Neozygites*, *Zoophthora* and *Tarichium*. — *Sydowia* 43: 39–122.
- Keller, S. 1994. The fungus *Zoophthora elateridiphaga* as an important mortality factor of the click beetle *Agriotus sputator*. — *Journal of Invertebrate Pathology* 63: 90–91.
- Keller, S. 1997. The genus *Neozygites* (Zygomycetes, Entomophthorales) with special reference to species found in tropical regions. — *Sydowia* 49(2): 118–146.
- Keller, S. 2002. The genus *Entomophthora* (Zygomycetes, Entomophthorales) with a description of five new species. — *Sydowia* 54(2): 157–197.
- Keller, S. 2006. Species of Entomophthorales attacking aphids with description of two new species. — *Sydowia* 58(1): 38–74.
- Keller, S. 2007a. Arthropod-pathogenic Entomophthorales: Biology, Ecology, Identification. — COST action 842, ISBN 978-92-898-0037-2, pp 124–126.
- Keller, S. 2007b. Arthropod-pathogenic Entomophthorales of Switzerland. III. First addition. — *Sydowia* 59: 75–113.
- Keller, S. 2008. The arthropod-pathogenic Entomophthorales from Switzerland – is central Europe the centre of their global species-richness? — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 81: 39–51.
- Keller, S. 2012. Arthropod-pathogenic Entomophthorales from Switzerland. IV. Second addition. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 85: 115–130.
- Keller, S. & Petrini, O. 2005. Keys to the identification of the arthropod-pathogenic genera of the families Entomophthoraceae and Neozygitaceae (Zygomycetes), with descriptions of three new subfamilies and a new genus. — *Sydowia* 57: 23–53.
- Keller, S., Weiser, J. & Wegensteiner, R. (2009). *Tarichium hylobii* sp. nov., a pathogen of *Hylobius abietis*. — *Sydowia* 61 (2): 249–254.
- King, D.S. 1977. Systematics of *Conidiobolus* (Entomophthorales) using numerical taxonomy. III. Descriptions of recognized species. — *Canadian Journal of Botany* 55: 718–729.
- Lakon, G. 1915. Zur Systematik der Entomophthorengattung *Tarichium*. — *Zeitschrift für Pflanzenkrankheiten* 25: 257–272.
- Lakon, G. 1919. Die Insektenfeinde aus der Familie der Entomophthoraceen. — *Zeitschrift für angewandte Entomologie* 5: 161–216.
- MacLeod, D.M. & Müller-Kögler, E. 1970. Insect pathogenic species originally described from their resting spores mostly as *Tarichium* species (Entomophthorales, Entomophthoraceae). — *Mycologia* 62: 33–66.
- Naranjo, S.E. & Steinkraus, D.C. 1988. Discovery of an entomophthoralean fungus (Zygomycetes: Entomophthorales) infecting adult northern corn rootworm, *Diabrotica barberi* (Coleoptera: Chrysomelidae). — *Journal of Invertebrate Pathology* 51: 298–300.
- Petch, T. 1932. A list of the entomogenous fungi of Great Britain. — *Transactions of the British Mycological Society* 17: 170–178.
- Petch, T. 1944. Notes on entomogenous fungi. — *Transactions of the British Mycological Society* 26: 81–93.
- Raunkiaer, C. 1893. Et par nye Snyltesvampe. Fundne og beskrevne. — *Botanisk Tidsskrift* 18: 108 (1893) [1892–93].
- Thaxter, R. 1888. The Entomophthoraceae of the United States. — *Memoirs of the Boston Society of Natural History* 4: 133–201.

(received August 25, 2013; accepted November 1, 2013; published December 31, 2013)

