

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 87 (2014)

Heft: 3-4

Artikel: Window gnats (Diptera : Anisopodidae) from beer traps in various countries across Europe

Autor: Dvoák, Libor

DOI: <https://doi.org/10.5169/seals-403091>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Window gnats (Diptera: Anisopodidae) from beer traps in various countries across Europe

LIBOR DVOŘÁK

Municipal Museum Mariánské Lázně, Goethovo náměstí 11, CZ-353 01 Mariánské Lázně, Czech Republic; lib.dvorak@seznam.cz

Results of catches of Anisopodidae trapped by beer traps in Europe during summer of 2006 and 2007 are presented. During this survey, 2234 specimens belonging to six species were caught. The most commonly found species was *Sylvicola punctatus* (Fabricius, 1787) followed by *S. cinctus* (Fabricius, 1787). *Sylvicola fenestralis* (Scopoli, 1763), *S. fuscatus* (Fabricius, 1775), *S. stackelbergi* Krivosheina & Menzel, 1998 and *S. zetterstedti* (Edwards, 1923) were caught in small numbers. The most interesting records are three males of *S. zetterstedti*, which are the third and the fourth known locality records from the Czech Republic, and one female of *S. stackelbergi*, the first record from Austria.

Key words: *Sylvicola*, bait traps, new record, distribution, Central Europe.

INTRODUCTION

Anisopodidae is a small family of dipteran insects with nine species of the genus *Sylvicola* Harris, 1776 present in continental Europe (De Jong 2004). Adults of all species of the genus *Sylvicola* feed on nectar and similar liquids (Kovář & Barták 2001) and are regularly attracted by traps baited with beer, syrup, mead, etc. This method is very effective for faunistic surveys of Anisopodidae, as exemplified in Estonia by Kurina (2006) who obtained most of the material of his study by using light and bait traps and by Dvořák (2014) who evaluated results of beer trapping reaching 3796 specimens.

In this contribution, the results of trapping Anisopodidae using beer traps in various European countries (Austria, Czech Republic, Estonia, Germany, Italy, Poland, Sweden, Switzerland, United Kingdom, see Fig. 1) in 2006 and 2007 are presented.

MATERIAL AND METHODS

Bigger PET bottles filled with 0.5 litre of beer were hung 1.5–2 metres above the ground on a branch of a tree or shrub (Dvořák 2007, Dvořák *et al.* 2008). In both years, the traps were exposed ca from mid July to mid August and material was taken out only once during the whole trapping period. The project was designed to study the usefulness of this method for faunistic research of various insect groups such as social wasps (Dvořák 2007, Dvořák *et al.* 2008), fly families Anisopodidae, Drosophilidae, Heleomyzidae, Lauxaniidae, and Scathophagidae, or Mecoptera and Dermaptera (most of this data was not previously published).

Localities are described in the following order: locality code, nearest village or town, coordinates in WGS 84, mapping quadrant code (for the localities in the



Fig. 1. Collecting localities in 2006 and 2007.

Czech Republic), a description of the locality including biotope and date of trap exposure. Localities from 2006 published by Dvořák (2007) are designated as «06» in the text, localities from 2007 published by Dvořák *et al.* (2008) are designated as «07» in the text. Unpublished localities are designated here by their respective years.

The studied localities are shown in Fig. 1.

LIST OF LOCALITIES

AUT 07-08, Austria, Styria, middle Enns valley, Aigen/Ennstal, 47°31'2.6" N 14°7'53.3" E, extensive meadow near a lake, 19.7.–22.8.2007, S. Heinrich.

AUT 07-09, Austria, Styria, middle Enns valley, Aigen/Ennstal, 47°31'33.2" N 14°6'53.8" E, meadow in the floodplain of Enns river, 649 m, 19.7.–22.8.2007, S. Heinrich.

- AUT 07-11, Austria, Enns valley, Reichraming, 47°53'51.1" N 14°28'4.9" E, south-facing cattle pasture, 20.7.–23.8.2007, S. Heinrich.
- AUT 07-12, Austria, Enns valley, Reichraming, 47°53'55" N 14°29'45.9" E, south-facing cattle pasture, 20.7.–23.8.2007, S. Heinrich.
- CZE 06-04, Czech Republic, Borkovice, 6753, 49°13'46" N, 14°37'35" E, Borkovická blata, peat-bog, 14.7.–15.8.2006, J. Máca.
- CZE 06-14, Czech Republic, Ostrava-Zábřeh, 6275, oak-elm forest in Polanský les forest, 14.7.–14.8.2006, M. Roháčová.
- CZE 06-33, Czech Republic, Hradec Králové, Nový Hradec Králové, 5861, 50°10'44" N, 15°52'11" E, town park forests, oak-hornbeam forest, 14.7.–14.8.2006, Š. + M. Mikátovi.
- CZE 06-37, Czech Republic, Štěmčehy-Dašov, 6860, 49°11'50" N, 15°42'56" E, secondary spruce forest, 11.7.–12.8.2006, V. Křivan.
- CZE 07-01, Czech Republic, Benešova Hora, 6848, 49°07'32.7" N, 13°42'15.3" E, pasture E of a village, 7.7.–7.8.2007, L. Dvořák.
- CZE 07-04, Czech Republic, Rovina, 6846, 49°09'31" N, 13°24'32.5" E, wet meadow in W part of an ancient village Rovina, 895 m, 10.7.–14.8.2007, L. Dvořák.
- CZE 07-05, Czech Republic, Nová Hůrka, 6845, 49°08'45.9" N, 13°19'11.6" E, mire W of a village, 895 m, 11.7.–15.8.2007, L. Dvořák.
- CZE 07-07, Czech Republic, Spálenec, 7049, 48°55'38" N, 13°58'00.9" E, wet meadow NW of a settlement, 11.7.–15.8.2007, L. Dvořák.
- CZE 07-08, Czech Republic, Sviadnov, 6375, 49°40'53.9" N, 18°19'29.1" E, crop field, 10.7.–9.8.2007, V. Ďurinová.
- CZE 07-22, Czech Republic, Prameny, 5942, 50°2'29.2" N, 12°44'18.1" E, Mokřady pod Vlčkem Nature Reserve, wet meadow, 10.7.–9.8.2007, P. Tájek.
- CZE 07-30, Czech Republic, Krásno, 5842, 50°6'22.5" N, 12°45'31.01" E, Krásenské rašelinš tě mire, dry exploited peat-bog W of a village, 714 m, 19.6.–10.7.2007, P. Tájek.
- CZE 07-31, Czech Republic, Bečov nad Teplou, 5942, 50°5'0.4" N, 12°49'23.63" E, Bečovské stráně slopes, dry meadow W of a town, 11.7.–9.8.2007, P. Tájek.
- CZE 07-35, Czech Republic, Lazy, 5941, 50°3'1.23" N, 12°38'5" E, submontane meadow W of a village, 802 m, 10.7.–9.8.2007, P. Tájek.
- CZE 07-37, Czech Republic, Mnichov, 5942, 50°3'23.01" N, 12°46'6.34" E, meadow under Pluhův bor site N of a village, 724 m, 10.7.–9.8.2007, P. Tájek.
- CZE 07-42, Czech Republic, Hamrníky, 6042, 49°57'16.7" N, 12°41'27.6" E, gardener's colony, 10.7.–10.8.2007, P. Tájek.
- EST 06-02, Estonia, Järvamaa, Koeru, 58°59'25" N, 26°0'35.3" E, spruce forest with several ash and maple trees, 19.7.–22.8.2006, K. Püssa.
- EST 06-03, Estonia, Järvamaa, Väike-Maarja, 59°10'31.9" N, 26°17'54.9" E, spruce forest with several ash and maple trees, 19.7.–22.8.2006, K. Püssa.
- EST 06-04, Estonia, Järvamaa, Väike-Maarja, 59°8'51.8" N, 26°18'34.3" E, spruce forest with several ash and maple trees, 19.7.–22.8.2006, K. Püssa.
- GER 06-02, Germany, Oberbayern, Traunstein, Reit im Winkl, spruce-beech forest, 900 m, 10.7.–10.8.2006, K. Wieser.
- GER 06-04, Germany, Oberbayern, Traunstein, Reit im Winkl, light spruce forest on stony debris, 1380 m a.s.l., 10.7.–10.8.2006, K. Wieser.
- GER 06-05, Germany, Lackenberg, 49°6'16" N, 13°18'37.9" E, sunny spruce forest, 10.7.–10.8.2006, J. Müller.
- GER 06-07, Germany, Mittelsteighütte, 49°05'43.4" N 13°14'59.2" E, beech-fir virgin forest, 10.7.–10.8.2006, J. Müller.
- GER 06-09, Germany, Lower Saxony, Barterode, Kattenbühl, 51°24'30.786" N, 9°39'30.878" E, pine forest, 10.7.–14.8.2006, M. Irskens.
- GER 06-12, Germany, Hesse, Roszbach, Pfaffenberg, 51°18'31.955" N, 9°47'13.325" E, beech forest, 10.7.–14.8.2006, M. Irskens.
- ITA 07-01, Italy, Livenza (Pordenone), 46°1'17.8" N 12°28'42.8" E, plane-maple forest, 25.7.–7.9.2007, L. Patrizia.
- POL 07-01, Poland, Malopolska Upland, Piekary, Góra Grodzisko, 50°00'27" N, 19°48'20" E, seminatural broadleaf forest, 12.7.–14.8.2007, H. Szentgyörgyi.
- SWE 06-01, Sweden, Uppsala, Lapland, Abisko, 68°21'10.7" N 18°49'4.9" E, birch forest, 12.7.–15.8.2006, O. Bohuslavová.
- SWE 1, Sweden, Uppsala, Forkarby, Skörkulla, 59°54'15" N, 17°33'47" E, spruce forest, 13.7.–12.8.2007, E. Sjödin.
- SWE 5, Sweden, Uppsala, Hagby, Myrängen, 59°48'36" N, 17°23'18" E, moraine spruce-pine forest, 13.7.–12.8.2007, E. Sjödin.

- SWE 7, Sweden, Uppsala, Forkarby, Marsta, 59°56'2" N, 17°34'52" E, mixed forest, 13.7.–12.8.2007, E. Sjödin.
- SWI 06-01, Switzerland, Jura, Delémont, 47°24'69" N, 7°19'39" E, mixed forest, 10.7.–15.8.2006, Ch. Péré.
- UK 06-10, United Kingdom, Berkshire, Newbury, Lambourn Downs, 51°29'19" N, 1°26'28" W, oak, sycamore, young plantation, 11.7.–10.8.2006, E.G. Chambers.
- UK 06-11, United Kingdom, Berkshire, Newbury, Lambourn Downs, 51°28'43" N, 1°25'36" W, beech, oak, ash, mature wood, 11.7.–10.8.2006, E.G. Chambers.
- UK 06-12, United Kingdom, Berkshire, Newbury, Lambourn Downs, 51°27'52" N, 1°27'46" W, sycamore, oak, ash, beech, mature wood, 11.7.–10.8.2006, E.G. Chambers.
- UK 06-13, United Kingdom, Berkshire, Newbury, Lambourn Downs, 51°27'36" N, 1°26'44" W, beech, elder, ash, field maple, mature wood, 11.7.–10.8.2006, E.G. Chambers.
- UK 06-14, United Kingdom, Berkshire, Newbury, Lambourn Downs, 51°27'32" N, 1°25'06" W, beech, ash, sycamore, mature wood, 11.7.–10.8.2006, E.G. Chambers.
- UK 06-15, United Kingdom, Berkshire, Newbury, Lambourn Downs, 51°29'19" N, 1°25'31" W, oak, ash, sycamore, mature wood, 11.7.–10.8.2006, E.G. Chambers.
- UK 06-17, United Kingdom, Berkshire, Reading, Chilterns, 51°31'24" N, 1°05'35" W, beech, oak, mature wood, 11.7.–11.8.2006, E.G. Chambers.
- UK 06-18, United Kingdom, Berkshire, Reading, Chilterns, 51°31'08" N, 1°05'20" W, silver birch, oak, silver fir, mature wood, 11.7.–11.8.2006, E.G. Chambers.
- UK 06-19, United Kingdom, Berkshire, Reading, Chilterns, 51°30'39" N, 1°06'49" W, ash, beech, yew, mature wood, 12.7.–11.8.2006, E.G. Chambers.
- UK 06-20, United Kingdom, Berkshire, Reading, Chilterns, 51°30'48" N, 1°05'15" W, beech, mature wood, 12.7.–11.8.2006, E.G. Chambers.
- UK 06-21, United Kingdom, Berkshire, Reading, Chilterns, 51°30'19" N, 1°05'16" W, beech, sycamore, holly, mature wood, 12.7.–11.8.2006, E.G. Chambers.

The material was identified based on the papers of Haenni (1997), Krivosheina & Menzel (1998), and Michelsen (1999). The nomenclature follows De Jong (2004). Voucher specimens (pinned or preserved in small vials with ethanol) are deposited in the collection of the Municipal museum Mariánské Lázně, Czech Republic.

LIST OF SPECIES

Altogether 2234 specimens of Anisopodidae belonging to six species of the genus *Sylvicola* have been trapped using 46 traps and identified.

Sylvicola cinctus (Fabricius, 1787)

AUT 07-11, 1 ♂, 2 ♀♀. CZE 06-04, 1 ♂. CZE 06-14, 2 ♀♀. CZE 06-33, 3 ♀♀. CZE 06-37, 1 ♀. CZE 07-04, 7 ♂♂, 17 ♀♀. CZE 07-05, 21 ♂♂, 41 ♀♀. CZE 07-22, 1 ♀. CZE 07-31, 1 ♀. CZE 07-35, 2 ♂♂, 1 ♀. CZE 07-37, 7 ♀♀. CZE 07-42, 3 ♀♀. EST 06-02, 3 ♀♀. EST 06-03, 1 ♂. EST 06-04, 2 ♂♂, 2 ♀♀. GER 06-02, 1 ♀. GER 06-04, 4 ♀♀. GER 06-05, 7 ♀♀. GER 06-07, 1 ♀. GER 06-09, 1 ♀. GER 06-12, 1 ♀. ITA 07-01, 1 ♂, 4 ♀♀. POL 07-01, 1 ♀. SWE 06-01, 2 ♂♂. SWE 1, 1 ♂. SWE 5, 3 ♀♀. SWE 7, 3 ♂♂, 22 ♀♀. UK 06-10, 1 ♂, 4 ♀♀. UK 06-11, 14 ♀♀. UK 06-12, 1 ♂, 5 ♀♀. UK 06-13, 1 ♀. UK 06-14, 2 ♀♀. UK 06-15, 2 ♀♀. UK 06-17, 7 ♂♂, 60 ♀♀. UK 06-18, 1 ♂, 22 ♀♀. UK 06-19, 1 ♂, 7 ♀♀. UK 06-20, 63 ♀♀. UK 06-21, 5 ♀♀.

Overall this is the most common species known from almost all of the European countries. This was the second most common species found during this research. It is one of two most commonly found species in forests, while it seems rather rare in open stands.

***Sylvicola fenestralis* (Scopoli, 1763)**

UK 06-13, 1 ♀.

Species distributed in almost all of Europe, but it is rarely encountered.

***Sylvicola fuscatus* (Fabricius, 1775)**

AUT 07-09, 18 ♂♂, 9 ♀♀. AUT 07-11, 1 ♂. AUT 07-12, 1 ♂. CZE 06-33, 2 ♀♀. CZE 06-37, 1 ♀. CZE 07-35, 1 ♂, 1 ♀. CZE 07-37, 1 ♀. CZE 07-42, 1 ♂, 7 ♀♀. EST 06-04, 12 ♀♀. POL 07-01, 2 ♂♂, 2 ♀♀. SWI 06-01, 1 ♀.

This is a common species known from northern, central, and western Europe. It is a surprisingly rare species in both forest and open stands.

***Sylvicola punctatus* (Fabricius, 1787)**

AUT 07-09, 134 ♂♂, 111 ♀♀. AUT 07-08, 10 ♂♂, 16 ♀♀. AUT 07-11, 10 ♂♂, 6 ♀♀. AUT 07-12, 16 ♂♂, 18 ♀♀. CZE 06-04, 11 ♂♂, 9 ♀♀. CZE 06-33, 1 ♀. CZE 06-37, 7 ♀♀. CZE 07-01, 76 ♂♂, 114 ♀♀. CZE 07-04, 6 ♂♂, 10 ♀♀. CZE 07-05, 1 ♀. CZE 07-07, 3 ♂♂, 8 ♀♀. CZE 07-08, 1 ♀. CZE 07-22, 100 ♂♂, 61 ♀♀. CZE 07-30, 141 ♂♂, 232 ♀♀. CZE 07-31, 20 ♂♂, 20 ♀♀. CZE 07-35, 41 ♂♂, 269 ♀♀. CZE 07-37, 28 ♂♂, 34 ♀♀. EST 06-03, 4 ♀♀. EST 06-04, 4 ♂♂, 2 ♀♀. GER 06-02, 1 ♂. GER 06-04, 1 ♂. GER 06-09, 1 ♀. ITA 07-01, 1 ♀. SWE 1, 6 ♀♀. SWE 5, 2 ♀♀. SWE 7, 14 ♂♂, 48 ♀♀. SWI 06-01, 1 ♂, 1 ♀. UK 06-10, 12 ♂♂, 28 ♀♀. UK 06-11, 10 ♀♀. UK 06-12, 13 ♂♂, 17 ♀♀. UK 06-13, 4 ♀♀. UK 06-14, 5 ♂♂, 17 ♀♀. UK 06-15, 2 ♂♂, 8 ♀♀. UK 06-17, 3 ♂♂, 19 ♀♀. UK 06-18, 2 ♂♂, 2 ♀♀. UK 06-19, 1 ♂, 13 ♀♀. UK 06-20, 19 ♀♀. UK 06-21, 2 ♀♀.

Overall this is a common species known from almost all European countries except in the southeast. It was the most collected species found within this research, being predominant in forests and very commonly found (more than 90%) in open stands.

***Sylvicola stackelbergi* Krivosheina & Menzel, 1998**

AUT 07-09, 1 ♀. EST 06-02, 2 ♂♂. EST 06-03, 1 ♂, 1 ♀. EST 06-04, 13 ♂♂, 3 ♀♀. SWE 06-01, 1 ♀. SWE 5, 1 ♂, 2 ♀♀.

The most interesting finding in this paper is the capture of one female of *S. stackelbergi* from Austria (Figs 2–4). *S. stackelbergi* was described from NW Russia (Krivosheina & Menzel 1998) and was later found in Finland, Sweden, the Netherlands, and Estonia (Krivosheina & Menzel 2002, Kurina 2006, Haarto 2011). The only published record from central Europe is from Slovakia (Ševčík 2011). This is the second record from central Europe and it is a new species for Austria.

***Sylvicola zetterstedti* (Edwards, 1923)**

CZE 07-04, 2 ♂♂. CZE 07-05, 1 ♂.

This species was published as a new member of the fauna of the Czech Republic by Dvořák (2014), who has summarized the at present known species' distribution in Europe. All previous records of *S. zetterstedti* from the Czech Republic originate from the vicinity of Mariánské Lázně, New records are from the Bohe-



Fig. 2. *Sylvicola stackelbergi* from Austria, top view. Photo: Zbyněk Kejval.



Fig. 3. Frons of *Sylvicola stackelbergi* from Austria. Photo: Zbyněk Kejval.



Fig. 4. *Sylvicola stackelbergi* from Austria, genital fork. Photo: Zbyněk Kejval.




Fig. 5. Known distribution of *Sylvicola zetterstedti* in the Czech Republic. Circles: localities published by Dvořák (2014). Stars: localities presented in this paper. Square: unpublished localities found in 2014.

mian Forest (= the Šumava Mts.); one new locality of *S. zetterstedti* was discovered in the vicinity of Mariánské Lázně, (see Fig. 5). The new localities as well as published records clearly show that *S. zetterstedti* inhabits forests or shrubby damp localities in the Czech Republic.

Sylvicola sp.

CZE 07-35, 1 ♀.

Unidentified female with genital plate different from all known species. It is similar to *S. cinctus* and *S. zetterstedti*.

ACKNOWLEDGEMENTS

I wish to thank everyone who installed and recovered the traps and Zbyněk Kejval (Museum of the Chodsko region, Domažlice, Czech Republic) for preparing the Figures 2–4. Anthony Daglio (Orlando, Florida, USA) revised the English.

REFERENCES

- De Jong, H. 2004. Fauna Europaea: Anisopodidae. *In*: Beuk, P. & Pape, T. (eds): Fauna Europaea: Diptera: Nematocera. — Fauna Europaea, version 1.0, <http://www.faunaeur.org> (accessed 23.10.2014).
- Dvořák, L. 2007. Social wasps (Hymenoptera: Vespidae) trapped with beer in European forest ecosystems. — *Acta Musei Moraviae, Scientiae biologicae* 92: 181–204.
- Dvořák, L., Castro, L. & Roberts, S.P.M. 2008. Social wasps (Hymenoptera: Vespidae) trapped with beer bait in European open ecosystems. — *Acta Musei Moraviae, Scientiae biologicae*, 93: 105–130.

- Dvořák, L. 2014. Window gnats (Diptera: Anisopodidae) from beer traps in the vicinity of Mariánské Lázně with the first records of *Sylvicola zetterstedti* (Edwards, 1923) from the Czech Republic. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 87: 41–48.
- Haarto, A. 2011: Suomen ikkunasääsket - Finnish window gnats (Diptera, Anisopodidae). — *w-album* 10: 3–13.
- Haenni, J.-P. 1997. Anisopodidae (Diptera) de la faune de Suisse, avec la description d'une espèce nouvelle. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 70: 177–186.
- Kovář, I. & Barták, M. 2001. Anisopodidae, pp. 101–104. *In*: Barták, M. & Vaňhara, J. (eds): *Diptera in an industrially affected region (north-western Bohemia, Břlina and Duchcov Environs) II*. — *Folia Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis, Biologia* 105: 1–274.
- Krivosheina, N.P. & Menzel, F. 1998. The Palaearctic species of the genus *Sylvicola* Harris, 1776 (Diptera, Anisopodidae). — *Beiträge zur Entomologie* 48 (1): 201–217.
- Krivosheina, N.P. & Menzel, F. 2002. [The morphology of females of the xylo-mycetophilous species of the genus *Sylvicola* Harris (Diptera, Anisopodidae)]. — *Entomologicheskoe obozrenie*, 81(2): 424–438 (in Russian, English abstract).
- Kurina, O. 2006. A review of Estonian wood gnats (Diptera: Anisopodidae). — *Sahlbergia* 11: 18–22.
- Michelsen, V. 1999. Wood gnats of the genus *Sylvicola* (Diptera, Anisopodidae): taxonomic status, family assignment, and review of nominal species described by J.C. Fabricius. — *Tijdschrift voor entomologie* 142: 69–75.
- Ševčík, J. 2011. Anisopodidae and Dixidae (Diptera) of the Gemer region, with a new record for Slovakia. — *Časopis Slezského Muzea Opava (A)*, 60: 181–184, 2011.

(received July 7, 2014; accepted October 23, 2014; published December 31, 2014)