

Zeitschrift: Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the Swiss
Entomological Society

Herausgeber: Schweizerische Entomologische Gesellschaft

Band: 88 (2015)

Heft: 1-2

Artikel: A new stonefly from Greece, *Leuctra pavesii* sp. n. (Plecoptera :
Leuctridae)

Autor: Vinçon, Gilles

DOI: <https://doi.org/10.5169/seals-514995>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A new stonefly from Greece, *Leuctra pavesii* sp. n. (Plecoptera: Leuctridae)

GILLES VINÇON

55 Bd Joseph Vallier, F 38100 Grenoble, France; vincon@kls-logistic.fr

Leuctra pavesii sp. n. from the Cephalonia Island (Greece) is described and compared with closely related species: *L. hippopoides* Kačanski & Zwick, 1970, *L. pseudohippopus* Raušer, 1965 and *L. hippopus* Kempny, 1899.

Keywords: Plecoptera, Stoneflies, *Leuctra pavesii* sp. n., new species, Greece, Cephalonia.

MATERIAL AND METHODS

This study deals with stoneflies collected in 2001 on the Cephalonia Island by Dr. Pavesi. They were previously stored in the collection of Carlalberto Ravizza, Milan, Italy, and given to me for study in 2012.

The specimens are preserved in 70 % ethanol. The holotype and 1 male, 1 female (paratypes) are deposited in the collection of the Museum of Zoology, Lausanne (MZL) (Switzerland). Other paratypes are stored in the Gilles Vinçon Collection.

RESULTS AND DISCUSSION

Leuctra pavesii sp. n.

(Figs 1–2)

Material examined. **Holotype** male: Greece, Cephalonia Island, 2 km east of Nikolaos Mount, 140 m, 20.03.2001, deposited in the MZL (Pavesi leg.). 1 ♂, 1 ♀ **paratypes** same locality and date, deposited in the MZL. **Other paratypes:** same locality and date: 1 ♂, 1 ♀ (Pavesi leg.). **Other material:** Cephalonia, Koulourata Mount, 180 m a.s.l., 15.03.2001, 2 ♂♂, 2 ♀♀ (Pavesi leg.).

Description (Figs 1a–e). Medium sized *Leuctra* species: ♂ body length 5.1–5.5 mm, ♀ 6.5–8.2 mm. Fore wing length ♂ 6.3–6.5 mm, ♀ 7.2–8.5 mm. General color brown. Head dark brown. Antennae brownish, with short erect bristles at the tip of each segment; bristles as long as antennal segment width. Pronotum brown with dark pattern. Legs light brown with tarsal segments darker. Body, wings and legs covered with short hairs.

Male abdomen (Figs 1a–b, 1d–e): terga I–VI simple, terga VII–X modified. Tergum VII with median bell-shaped membranous field; anterior margin interrupted on nearly one third of segment's width (Fig. 1a). Tergum VIII: antecosta interrupted medially on nearly one fourth of segment's width. From the inner corners of




Fig. 1. *Leuctra pavesii* sp. n.: male abdomen in dorsal view (a), lateral view (b), paraprocts in ventral view (d), lateral view (e). Female abdomen in ventral view (c).

the antecosta, two long sclerotized expansions extend backwards, nearly reaching two thirds of segment's length (Fig. 1a). Expansions parallel, hardly prominent in lateral view (Fig. 1b), ending in a blunt tip and separated by a median membranous field. Tergum IX: anterior margin divided for more than half segment's width; wide trapezoidal sclerite in middle of median membranous field. Tergum X anteriorly bilobed and with wide rounded posterior notch where the epiproct is placed. Cerci very long, covered with long erect setae. Epiproct rounded with short stalk (Fig. 1a). Style of paraprocts with rounded base flanked by well visible lateral lobe, and with long thin inner expansion (Fig. 1d); expansions sickle-shaped ending in a sharp




Fig. 2. *Leuctra hippopoides*: male abdomen in dorsal view (a), paraprocts in lateral view (b). — *Leuctra pseudohippopoides*: male abdomen in dorsal view (c), paraprocts in lateral view (d).

point in lateral view (Fig. 1e). Specilla long, straight, blade-shaped, much longer than styles, progressively getting narrower toward sharp tip (Fig. 1d–e). Sternum IX without vesicle (Fig. 1b).

Female abdomen (Fig. 1c): sternum VII with wide sub-rectangular median sclerite; posterior edge slightly rounded, partly projecting on anterior part of sternum VIII (Fig. 1c). Sternum VIII: subgenital plate nearly semicircular with two long lateral lobes and a median small triangular expansion between them. Each lobe about two times longer than wide, ending in a rounded tip slightly bent inward, and widely separated by a median membranous field (Fig. 1c). Plate brownish, less pigmented than dark sclerotized lobes.

Affinities. In the *L. hippopus* group, *L. pavesii* sp. n. is characterised by the two parallel sclerotized expansions on tergite VIII, also characterizing other species of the same group: *L. andalusiaca* Aubert, 1962 (Iberian Peninsula), *L. artvinensis* Vinçon & Sivec, 2001 (eastern Anatolia), *L. collaris* Martynov, 1928 (Anatolia, Caucasus and Elburz), *L. elisabethae* Ravizza, 1985 (Italian Peninsula), *L. fraterna* Morton, 1930 (Corsica), *L. gallica* Aubert, 1953 (Iberian Peninsula and France), *L. hippopoides* (western and southern Balkan Peninsula), *L. hippopus* (Europe, Anatolia and Lebanon) and *L. pseudohippopus* (eastern Balkan Peninsula). It is more closely related to the three last ones. It differs from *L. hippopoides* and *L. pseudohippopus* in the male by the shape of the tergal ornamentations, especially the shape of the median plate on tergite IX and the shape of the epiproct (Figs 1a, 2a, 2c), by the shape of the specilla, nearly straight in lateral view in *L. pavesii* (Fig. 1e) and with curved tip in the two other species (Figs 2b, 2d) and by the lack of ventral

vesicle, and in the female by the triangular sclerotized expansion between the two lobes of the subgenital plate in *L. pavesii* (Fig. 1c), that is absent in *L. hippopoides* (Kačanski & Zwick 1970, Fig. 3c), and in *L. pseudohippopus* (Raušer 1965, Fig. 12). It separates from *L. hippopus* in the male by: – trapezoidal sclerite of tergite IX lacking the longitudinal median depression that is present in *L. hippopus* (Ravizza & Vinçon 1998, Fig. 17a), – different shape of the epiproct, – lack of ventral vesicle and – in the female, presence of a triangular sclerotized expansion between the 2 lobes (Fig. 1c).

Geographical distribution and ecology. *L. pavesii* sp. n. seems to be a micro-endemic species only known from the Greek island of Cephalonia where it occurs in mountain brooks (140–180 m a.s.l.). The flight period coincides with early spring (II–IV). *Brachyptera beali beali* (Navás, 1923) and *Protonemura mattheyi* (Aubert, 1956) have been collected on the same island.

Etymology. This species is dedicated to Dr. Maurizio Pavesi, first collector of this species.

ACKNOWLEDGEMENTS

I thank Dr. John Brittain and Dr. Jean-Paul Reding for having kindly revised the English text and Dr. Carlalberto Ravizza and Dr. Maurizio Pavesi for the gift of the material.

REFERENCES

- Kačanski, D. & Zwick, P. 1970. Neue und wenig bekannte Plecopteren aus Jugoslawien. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft 43 (1): 1–16.
Ravizza, C. & Vinçon, G., 1998. Les Leuctridae (Plecoptera, Leuctridae) des Alpes. — Mitteilungen der Schweizerischen Entomologischen Gesellschaft, 71: 285–342.
Raušer, J. 1965. Plecoptera bulgarica - II. — Acta faunistica entomologica Musei Nationalis Pragae, 10 (92): 125–138.

(accepted February 10, 2015; published June 30, 2015)